

CHEMISTRY

A European Journal

A Journal of



Accepted Article

Title: Total Synthesis of Skylamycins A-C

Authors: Richard James Payne, Andrew Giltrap, Jake Haeckl, Kenji Kurita, and Roger Linington

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Chem. Eur. J.* 10.1002/chem.201704277

Link to VoR: <http://dx.doi.org/10.1002/chem.201704277>

Supported by
ACES

WILEY-VCH

Total Synthesis of Skyllamycins A-C

Andrew M. Giltrap,^[a] F. P. Jake Haeckl,^[b] Kenji L. Kurita,^[b] Roger G. Linington^[b] and Richard J. Payne^{[a]*}

Dedicated to Professor Chris Abell on the occasion of his 60th Birthday

Abstract: The skyllamycins are a family of highly functionalized non-ribosomal cyclic depsipeptide natural products which contain the extremely rare α -OH-glycine functionality. Herein we report the first total synthesis of skyllamycins A-C, together with the biofilm inhibitory activity of the natural products. Linear peptide precursors for each natural product were prepared through an efficient solid-phase route incorporating a number of synthetic modified amino acids. A novel macrocyclization step between a C-terminal amide and a N-terminal glyoxylamide moiety served as a key transformation to install the unique α -OH-glycine unit and generate the natural products in the final step of the synthesis.

Skyllamycins A (1), B (2), and C (3) are a family of non-ribosomal cyclic depsipeptide natural products isolated from *Streptomyces* sp.^[1] (Figure 1). They contain a number of unusual structural features: three β -OH amino acids (β -OH-Phe, β -OH-O-Me-Tyr, β -OH-Leu) (blue), an N-terminal cinnamoyl residue (red), a β -Me-aspartic acid (β -Me-Asp) (green) and the extremely rare α -OH-Gly residue (purple). The biosynthetic origin of these modifications has been recently investigated.^[1b] A single P450 monooxygenase (P450_{sky}) was found to be responsible for catalyzing the β -hydroxylation of Phe, Leu and O-Me-Tyr, producing the (3S)-diastereomer in each case,^[2] however the exact timing at which the α -OH-Gly residue is installed remains unknown. Indeed, α -OH-Gly residues are extremely rare and to date this unusual moiety has only been found in one other natural product, the structurally simpler linear peptide anti-tumour agent spargalin.^[3]

The stereochemistry of the amino acids present in 1-3 was not assigned during the initial isolation; however further work by Süssmuth and co-workers^[4] confirmed the absolute configuration of the amino acids *via* a combination of chiral HPLC/GC and Marfey's analysis. Due to the instability of the α -OH-Gly residue, the configuration could not be confirmed experimentally. Instead, by estimating the distances between protons using NOESY measurements, followed by molecular dynamics simulations, the authors proposed that the α -OH-Gly residue was (S)-configured. These calculations also predicted the presence of a strong intramolecular hydrogen bond network when the α -OH-Gly residue was in the (S)-configuration.




Figure 1. Structure of skyllamycins A-C (1-3).

The skyllamycins also possess interesting biological activity. They were initially discovered in the search for novel agents that inhibit the platelet-derived growth factor (PDGF) signaling pathway. We recently re-isolated them in our search for inhibitors of *Pseudomonas aeruginosa* biofilms.^[1c] Traditional antibiotics target bacteria in the planktonic state, however the formation of biofilms allow bacteria to evade their action.^[5] As such, agents which inhibit biofilm formation or clear pre-attached biofilms, a major contributor to the persistence of nosocomial infections, represent a potentially useful therapeutic strategy. To our knowledge, the skyllamycins are the first known class of cyclic peptide biofilm disruptors. Given their highly unusual structural features, together with their important anti-biofilm activity, we sought to develop a concise total synthesis of skyllamycins A-C (1-3) which is described herein which, to date, have not succumbed to total synthesis.

Retrosynthetically, we envisioned a novel macrocyclization step to generate the α -OH-Gly residue that would necessitate preparation of modified linear peptides 4-6 containing a C-terminal amide and a N-terminal glyoxylamide functionality (Scheme 1A). We reasoned that, due to the inherent instability of the α -OH-Gly in linear peptides,^[6] formation in the final step would reduce the likelihood of decomposition during synthesis. Indeed, the stability of this residue in the natural products is proposed to be due to a hydrogen-bonding network in the mature structure.^[1b] Furthermore, the presence of this intramolecular hydrogen-bonding network during the cyclization reaction could potentially induce α -OH-Gly formation with the desired (S)-stereochemistry. Glyoxylamides 4-6 in turn could be accessed from resin-bound 7-9 by resin cleavage followed by oxidative cleavage of the N-terminal serine residue.^[7] We envisioned that resin-bound peptides 7-9 could be prepared *via* Fmoc-solid-phase peptide synthesis (SPPS) from Sieber amide resin which possesses a linker that can be cleaved under mild acidic conditions (1% TFA in CH₂Cl₂).^[8]


Before the assembly of skyllamycins A-C could begin, synthesis of the unusual amino acids present in the natural products was required (Scheme 1B). The suitably protected β -OH-Leu 10, β -OH-Phe 11, β -OH-O-Me-Tyr 12 were synthesized utilizing a unified strategy, namely organometallic addition to Garner's aldehyde^[9] followed by deprotection, oxidation and protecting group manipulations^[10] (see Supporting Information for synthetic details). The β -OH groups were protected as an oxazolidine, based on the pseudo-proline moiety commonly

[a] Andrew Giltrap and Prof. Richard J. Payne
School of Chemistry
The University of Sydney
Sydney, NSW 2006, AUSTRALIA
E-mail: richard.payne@sydney.edu.au

[b] F. P. Jake Haeckl, Dr Kenji L. Kurita and Prof. Roger L. Linington
Department of Chemistry
Simon Fraser University
Burnaby, British Columbia BC V5A 1S6, Canada


Supporting information for this article is given via a link at the end of the document.

coupled, the latter under microwave conditions, to yield the resin-bound pentapeptide **18**.



To begin the synthesis, Fmoc-D-Trp-OH was loaded onto Sieber amide resin using (7-azabenzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyAOP) and *N*-methylmorpholine (NMM) (Scheme 2). Next, oxazolidine-protected Fmoc- β -OH-O-Me-Tyr-OH (**12**) was coupled for 16 h using 1-[*bis*(dimethylamino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxid hexafluorophosphate (HATU), 1-hydroxy-7-azabenzotriazole (HOAt) and *i*Pr₂NEt to yield resin-bound dipeptide **16**. Due to the hindered nature of the oxazolidine-protected N-terminus, Fmoc-Pro-OH was coupled under optimized conditions, namely HATU, HOAt and *i*Pr₂NEt with microwave irradiation at 50 °C for 1 hour to afford **17**. Protected Fmoc- β -OH-Phe-OH (**11**) and Fmoc-Gly-OH were subsequently

Next, the key on-resin esterification step was carried out utilizing oxazolidine-protected Fmoc- β -OH-Leu-OH **10**, *N,N'*-diisopropylcarbodiimide (DIC) and catalytic *N,N*-dimethylaminopyridine (DMAP) affording resin-bound **22-24**. Importantly, we could not detect any epimerization of the β -OH-Leu moiety under the DMAP-catalyzed esterification conditions. With the key ester bond now in place, Fmoc-D-Leu-OH was effectively coupled to resin-bound peptides **23** and **24** as described previously, using HATU, HOAt and *i*Pr₂NEt at 50 °C for 1 hour under microwave heating. Unfortunately, when these conditions were applied to resin-bound intermediate **22** containing the β -Me-Asp residue, a significant amount of by-product with a mass 18 Da less than the product was observed. This was likely owing to aspartimide formation^[16] promoted by a conformational change in the peptide backbone induced by the β -Me group. Gratifyingly, this side-reaction could be suppressed by excluding an additional base from the coupling reaction, instead utilizing DIC and HOAt at 50 °C with microwave heating. These optimized conditions were subsequently employed, followed by elongation with side chain unprotected Fmoc-Ser-OH to afford resin-bound peptides **7-9**.



Scheme 4. Total Synthesis of skyllamycins A-C (1-3).

between the synthetic and isolated skyllamycins A-C suggested that they were the same enantiomer (see Supporting Information). Finally, the synthetic natural products were analyzed in a *P. aeruginosa* biofilm inhibition assay alongside the authentic natural products. Pleasingly, the synthetic and isolated products **1-3** showed comparable biofilm inhibition activity, further confirming the identity of the synthetic natural products (see Supporting Information).

In conclusion, we have successfully completed the first total synthesis of skyllamycins A-C (**1-3**). A key final macrocyclization step with concomitant formation of the unusual α -OH-Gly moiety was used to construct the natural products, which were identical in all respects to isolated skyllamycins A-C. This work lays the foundation for further studies into the biosynthesis of these natural products, the generation of synthetic skyllamycin analogues, as well as the interrogation of the mechanism of their biofilm inhibitory activity. Studies toward this end will be reported in due course.

Acknowledgements

We thank Dr. Ian Luck and Dr. Nick Proschogo (University of Sydney) for NMR spectroscopy and mass spectrometry technical support, respectively, and Walter Bray (UC, Santa Cruz) for support with the biofilm screening. We thank the Australian Postgraduate Award and John A. Lamberton Scholarship for PhD funding (A.M.G), ARC Future Fellowship to R.J.P (FT130100150), and NSERC Discovery support to R.G.L. (RGPIN-2016-03962).


Keywords: solid-phase synthesis • natural products • biofilms • peptides • cyclic peptides

- [1] a) S. Toki, T. Agatsuma, K. Ochiai, Y. Saitoh, K. Ando, S. Nakanishi, N. A. Lokker, N. A. Giese, Y. Matsuda, *J. Antibiot.* **2001**, *54*, 405-414; b) S. Pohle, C. Appelt, M. Roux, H. P. Fiedler, R. D. Süßmuth, *J. Am. Chem. Soc.* **2011**, *133*, 6194-6205; c) G. Navarro, A. T. Cheng, K. C. Peach, W. M. Bray, V. S. Bernan, F. H. Yildiz, R. G. Linington, *Antimicrob. Agents Chemother.* **2014**, *58*, 1092-1099.
- [2] a) S. Uhlmann, R. D. Süßmuth, M. J. Cryle, *ACS Chem. Biol.* **2013**, *8*, 2586-2596; b) K. Haslinger, C. Brieke, S. Uhlmann, L. Sieverling, R. D. Süßmuth, M. J. Cryle, *Angew. Chem. Int. Ed.* **2014**, *53*, 8518-8522.
- [3] a) T. Takeuchi, H. Iinuma, S. Kunimoto, T. Masuda, M. Ishizuka, M. Takeuchi, M. Hamada, H. Naganawa, S. Kondo, H. Umezawa, *J. Antibiot.* **1981**, *34*, 1619-1621; b) H. Umezawa, S. Kondo, H. Iinuma, S. Kunimoto, Y. Ikeda, H. Iwasawa, D. Ikeda, T. Takeuchi, *J. Antibiot.* **1981**, *34*, 1622-1624.
- [4] V. Schubert, F. Di Meo, P. L. Saadi, S. Bartoschek, H. P. Fiedler, P. Trouillas, R. D. Süßmuth, *Chem. Eur. J.* **2014**, *20*, 4948-4955.
- [5] P. S. Stewart, J. William Costerton, *The Lancet* **2001**, *358*, 135-138.
- [6] L. Lebreton, J. Annat, P. Derrepas, P. Dutartre, P. Renaut, *J. Med. Chem.* **1999**, *42*, 277-290.
- [7] a) K. F. Geoghegan, J. G. Stroh, *Bioconjugate Chem.* **1992**, *3*, 138-146; b) O. El-Mahdi, O. Melnyk, *Bioconjugate Chem.* **2013**, *24*, 735-765.
- [8] P. Sieber, *Tetrahedron Lett.* **1987**, *28*, 2107-2110.
- [9] P. Garner, J. M. Park, *J. Org. Chem.* **1987**, *52*, 2361-2364.
- [10] a) L. Williams, Z. D. Zhang, F. Shao, P. J. Carroll, M. M. Joullie, *Tetrahedron* **1996**, *52*, 11673-11694; b) N. Okamoto, O. Hara, K. Makino, Y. Hamada, *J. Org. Chem.* **2002**, *67*, 9210-9215; c) A. Nishida, H. Sorimachi, M. Iwaida, M. Matsumizu, T. Kawate, M. Nakagawa, *Synlett* **1998**, 389-390.
- [11] T. Wöhr, F. Wahl, A. Nefzi, B. Rohwedder, T. Sato, X. Sun, M. Mutter, *J. Am. Chem. Soc.* **1996**, *118*, 9218-9227.
- [12] S. Schabbert, M. D. Pierschbacher, R. H. Mattern, M. Goodman, *Bioorg. Med. Chem.* **2002**, *10*, 3331-3337.
- [13] D. Sun, P. Lai, W. Xie, J. Deng, Y. Jiang, *Synth. Commun.* **2007**, *37*, 2989-2994.
- [14] R. Krishnamoorthy, L. D. Vazquez-Serrano, J. A. Turk, J. A. Kowalski, A. G. Benson, N. T. Breaux, M. A. Lipton, *J. Am. Chem. Soc.* **2006**, *128*, 15392-15393.
- [15] C. Yue, J. Thierry, P. Potier, *Tetrahedron Lett.* **1993**, *34*, 323-326.
- [16] R. Subirós-Funosas, A. El-Faham, F. Albericio, *Tetrahedron* **2011**, *67*, 8595-8606.

Entry for the Table of Contents

COMMUNICATION

The total synthesis of the complex cyclic peptide natural products skyllamycin A, B and C is described. The successful syntheses hinged on an unprecedented macrocyclization reaction in the last step that furnished the natural products through the generation of the unusual α -OH-Gly moiety.



Andrew M. Giltrap, F. P. Jake Haeckl,
Kenji L. Kurita, Roger G. Linington and
Richard J. Payne*

Page No. – Page No.

Total Synthesis of Skyllamycins A-C