

Reductive Coupling Very Important Paper

International Edition: DOI: 10.1002/anie.201605162 German Edition: DOI: 10.1002/ange.201605162

Nickel-Catalyzed Reductive Amidation of Unactivated Alkyl **Bromides**

Eloisa Serrano and Ruben Martin*

Abstract: A user-friendly, nickel-catalyzed reductive amidation of unactivated primary, secondary, and tertiary alkyl bromides with isocyanates is described. This catalytic strategy offers an efficient synthesis of a wide range of aliphatic amides under mild conditions and with an excellent chemoselectivity profile while avoiding the use of stoichiometric and sensitive organometallic reagents.

Although unactivated alkyl halides are inherently disposed towards destructive β-hydride elimination and homodimerization pathways, these molecules have successfully been employed in a myriad of metal-catalyzed cross-coupling reactions.^[1] At present, the vast majority of these processes are based on stoichiometric, well-defined, and in many instances, air-sensitive organometallic species. Challenged by these drawbacks, recent years have witnessed the development of cross-electrophile coupling processes,^[2] becoming powerful and practical synthetic alternatives to classical crosscoupling reactions, achieving an otherwise similar molecular complexity under milder reaction conditions while avoiding the need for organometallic reagents.

Despite the advances realized, the palette of electrophilic partners in cross-electrophile processes remains rather limited when compared with classical nucleophile/electrophile regimes. It comes as a surprise that isocyanates, privileged synthons in industrial settings,^[3] have been virtually unexplored in cross-electrophile events with organic (pseudo)halides.^[4,5] This is likely due to the strong binding properties of isocyanates to low-valent transition-metal complexes, leading to unproductive dimerization or trimerization pathways.^[6] At present, cross-electrophile coupling reactions with isocvanates as coupling partners remain confined to substrates that rapidly undergo oxidative addition, such as aryl or benzyl halides lacking β-hydrogen atoms, thus preventing undesired pathways (Scheme 1, path a).^[7] Ideally, this field of expertise should include the use of unactivated alkyl halides possessing β -hydrogen atoms,^[1] thus resulting in a new synthetic route for rapidly preparing aliphatic amides, ubiquitous motifs in

[*]	E. Serrano, Prof. R. Martin Institute of Chemical Research of Catalonia (ICIQ) The Barcelona Institute of Science and Technology Av. Països Catalans 16, 43007 Tarragona (Spain) E-mail: rmartinromo@iciq.es
	Prof. R. Martin Catalan Institution for Research and Advanced Studies (ICREA) Passeig Lluïs Companys 23, 08010 Barcelona (Spain)
	Supporting information and the ORCID identification number(s the author/s) of this article can be found under http://dx.doi.org

s) for rg/10. 1002/anie.201605162.



Scheme 1. Amide synthesis through C-C bond formation using isocyanates. $1^\circ = \text{primary}$; $2^\circ = \text{secondary}$; $3^\circ = \text{tertiary}$.

hindered combinations

1°,2° and 3° amides

nucleophile

pharmaceuticals, agrochemicals, peptides, and polymers.^[8] Indeed, a close look into the literature data indicates that there is a paucity of highly chemoselective catalytic C-C bond-forming processes^[9] with improved flexibility, practicality, and generality that would give access to primary, secondary, or even tertiary amides at will, including hindered substrate combinations, while avoiding the handling of carbon monoxide (CO) at high pressures^[10] or well-defined and stoichiometric organometallic species.^[11] among others (Scheme 1, path b).^[12]

As part of our interest in reductive coupling reactions,^[13] we questioned whether a unified catalytic umpolung strategy through the in situ generation of carbogenic nucleophiles (II) from unactivated alkyl halides (I) and their coupling with isocyanates would constitute a generic platform for preparing aliphatic amides (III; Scheme 1, bottom). However, at the outset of our investigations it was unclear whether it would be possible to balance the high reactivity of isocyanates and the commonly observed parasitic β -hydride elimination or homodimerization pathways when using unactivated alkyl halides. Herein, we describe the successful realization of this concept, providing access to primary, secondary, and even tertiary alkyl amides by exploiting a previously unrecognized opportunity through sequential cross-coupling reactions of three different electrophiles.

We began our investigations by studying the reaction of 1a with isocyanate 2a (Scheme 2). The choice of 2a was not arbitrary, as primary amides can be prepared by simple deprotection of the *tert*-butyl group.^[14] After judicious

Angew. Chem. Int. Ed. 2016, 55, 1-6

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Wiley Online Library

Communications



Scheme 2. Optimization of the Ni-catalyzed reductive amidation of **1a** with **2a**. [a] Reaction conditions: **1a** (0.50 mmol), **2a** (0.75 mmol), NiBr₂ (3 mol%), ligand (4.5 mol%), Mn (0.75 mmol), DMF (1 mL) at RT, 16 h. [b] GC yield using *n*-decane as the internal standard. [c] Yield of isolated product. **1a-I** = 1-iodo-4-phenylbutane; **1a-OTs** = 4-phenylbutyl-4-methylbenzenesulfonate.

evaluation of the reaction parameters,^[15] we found that a combination of inexpensive NiBr₂ (3 mol%), L2 (4.5 mol %),^[16] and Mn as the reducing agent in DMF at RT, delivered 3a, which was isolated in 86% yield. Importantly, only traces of β-hydride elimination and homodimerization products were detected in the crude mixtures. Notably, ligand optimization revealed a crucial influence of the substitution pattern on the aromatic ring, with bipyridine ligands lacking ortho substituents (entry 2) or structurally similar phenanthroline ligands L4 and L5 providing inferior results (entries 4 and 5).^[17,18] Strikingly, the utilization of L3 had a deleterious impact on yield when using 2a as substrate (entry 3),^[19] revealing an interesting effect of the substituents located at the ortho position. As shown in entries 6-12, the use of other solvents, precatalysts, reducing agents, or 1a-I/1a-OTs analogues resulted in diminished yields of **3a**,^[20] thus showing the subtleties of our procedure. As expected, control experiments revealed that all of the reaction parameters were critical for success.[15]

Encouraged by these results, we turned our attention to study the preparative scope of our Ni-catalyzed reductive amidation of unactivated alkyl bromides with isocyanates (Scheme 3). As shown for **3a–3 f**, the procedure allowed for the coupling of either aromatic or aliphatic isocyanates with equal ease. Intriguingly, a procedure employing **L3** was particularly efficient when dealing with aromatic isocyanates.^[21,22] At present, we do not have any rational explanation for this behavior. Particularly illustrative was the chemoselectivity profile of the procedure, as substrates containing silyl ethers (**3k**), acetals (**3l**), esters (**3h**, **3o**, **3q**), amides (**3p**), nitriles (**3r**), heterocycles (**3q**), or chlorides (**3j**, **3n**) could all be perfectly accommodated. As shown for **3h** and **3n**, aryl pivalates and chlorides, substrates commonly employed in Ni-catalyzed cross-electrophile couplings, do not



Angewandte

I Edition Chemie

Scheme 3. Scope of alkyl bromides and isocyanates. Reaction conditions: as for Scheme 2, entry 1; Yields of isolated products, average of at least two independent runs. [a] **1a** (4.69 mmol). [b] NiBr₂ (10 mol%), **L2** (15 mol%). [c] [(TMEDA)Ni(*o*-tolyl)Cl] (15 mol%), **L2** (30 mol%). [d] NiBr₂ (10 mol%), **L3** (20 mol%), RNCO (0.5 mmol). [e] NiBr₂ (5 mol%), **L2** (7.5 mol%). TMEDA = tetramethylethylenediamine; cy = cyclohexyl; TBDPS = *tert*-butyldiphenylsilyl.

compete with the efficacy of this method. Boronic esters (3m) were tolerated as well, leaving an additional handle for further functionalization. Although in lower yields, we found that terminal epoxides could also participate in the targeted reaction (3s). Importantly, the reaction could be easily scaled up, obtaining 3a in similar yields. On the basis of these results, we wondered whether our procedure could accommodate unactivated secondary or tertiary alkyl halides. Despite the higher statistical propensity for β -hydride elimination pathways and increased steric hindrance around the C–Br bond, a host of cyclic and acyclic secondary alkyl bromides could be equally accommodated under otherwise identical reaction conditions (3t-3x). A noteworthy observation concerns the preferential formation of 3x from *exo-*1x, reinforcing the idea

www.angewandte.org

2

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

K These are not the final page numbers!

that radical species might come into play.^[23] The successful preparation of **3y** showcases the generality of our approach as unactivated tertiary alkyl halides have rarely been employed in metal-catalyzed cross-electrophile coupling reactions.^[24,25]

A close survey of the literature data reveals that a unified metal-catalyzed strategy for accessing primary, secondary, and tertiary amides remains elusive. In a final venture to unlock the full potential of our procedure, we sought to intercept the insitu generated **IV** upon subsequent addition of a proper electrophile, thus accessing tertiary aliphatic amides in a one-pot fashion (Scheme 4, top left). Although

Primary & tertiary aliphatic amides by reductive amidation procedures



Scheme 4. Sequential C-C bond-forming scenarios.

counterintuitive at first sight, the preparation of 4a-4d demonstrates the feasibility of this approach, constituting a formal cross-coupling reaction of three different electrophiles. On the other hand, primary amides could easily be obtained by deprotection of the *tert*-butyl group with Sc-(OTf)₃ (4e).^[26] Importantly, such a design principle allowed us to rapidly convert 1a into 4f, thus constituting a powerful alternative platform for handling flammable and toxic MeNCO in amidation technologies (Scheme 4, bottom).^[27] Taken together, the results in Scheme 3 and Scheme 4 demonstrate the prospective impact of our catalytic amidation procedure for accessing a wide variety of amides from simple starting materials in a straightforward manner.

Although a comprehensive mechanistic study should await further investigations, deuterium labelling experiments were performed to study the stereochemical course of the reaction (Scheme 5, top). Interestingly, we found a statistical mixture of diastereoisomers in **6** when exposing **5** to our optimized reaction conditions, suggesting the intermediacy of single-electron-transfer (SET) processes.^[23] In line with this notion, a complete racemization was detected when using (*R*)-(3-bromo-butyl)benzene (97 % *ee*) as substrate. We then turned our attention to study the reactivity of the putative Ni⁰(**L3**)₂ (**7**) and NiBr₂(**L3**) (**8**) species (Scheme 5, bottom).^[28,29] These compounds were easily prepared from either Ni(COD)₂ or NiBr₂ and were characterized by X-ray crystallography.^[15] As expected, both **7** and **8** were found to be



Scheme 5. Preliminary mechanistic studies.

catalytically competent as reaction intermediates, delivering **3d** in slightly lower yield to that observed in Scheme 3 (57% yield).^[30] More importantly, we found that **3d** could be obtained regardless of whether Mn was present or not with stoichiometric amounts of **7**. As expected, **8** delivers **3d** with similar yields to those shown for **7**. Although we cannot rigorously rule out other conceivable pathways,^[31] we propose a mechanistic scenario involving alkyl–Ni¹ species generated by comproportionation of in situ generated alkyl–Ni^{II} intermediates and Ni⁰(L)₂, followed by insertion of the isocyanate motif, and a final SET mediated by Mn.^[32,33]

In summary, we have described the first Ni-catalyzed reductive amidation of unactivated alkyl halides with isocyanates for accessing primary, secondary, or even tertiary amides through iterative techniques. The reaction proceeds under mild conditions, thus minimizing unproductive β -hydride elimination or dimerization/trimerization events, while accommodating a wide range of substrates with an excellent chemoselectivity profile. Further investigations on the mechanism and the elaboration of an asymmetric version of the reaction are currently being pursued in our laboratories.

Acknowledgements

We thank ICIQ, the European Research Council (ERC-277883), MINECO (CTQ2015-65496-R and Severo Ochoa Excellence Accreditation 2014–2018, SEV-2013-0319), FEDER, and the Cellex Foundation for support. Johnson Matthey, Umicore, and Nippon Chemical Industrial are acknowledged for a gift of metal and ligand sources. E.S. thanks MINECO for a FPI fellowship. We sincerely thank E. Escudero and E. Martin for all X-ray data.

Keywords: cross-coupling \cdot nickel \cdot reductive coupling \cdot synthetic methods

Angew. Chem. Int. Ed. 2016, 55, 1-6

- For selected reviews on catalytic cross-coupling reactions of unactivated alkyl halides through nucleophile/electrophile regimes, see: a) X. Hu, *Chem. Sci.* 2011, 2, 1867–1886; b) R. Jana, T. P. Pathak, M. S. Sigman, *Chem. Rev.* 2011, 111, 1417– 1492; c) A. C. Frisch, M. Beller, *Angew. Chem. Int. Ed.* 2005, 44, 674–688; *Angew. Chem.* 2005, 117, 680–695; d) M. R. Netherton, G. C. Fu, *Adv. Synth. Catal.* 2004, 346, 1525–1532; e) D. J. Cárdenas, *Angew. Chem. Int. Ed.* 2003, 42, 384–387; *Angew. Chem.* 2003, 115, 398–401.
- [2] For reviews on cross-electrophile couplings of organic halides, see: a) J. Gu, X. Wang, W. Xue, H. Gong, *Org. Chem. Front.* 2015, 2, 1411–1421; b) D. J. Weix, *Acc. Chem. Res.* 2015, 48, 1767–1775; c) T. Moragas, A. Correa, R. Martin, *Chem. Eur. J.* 2014, 20, 8242–8258; d) C. E. I. Knappke, S. Grupe, D. Gärtner, M. Corpet, C. Gosmini, A. Jacobi von Wangelin, *Chem. Eur. J.* 2014, 20, 6828–6842.
- [3] a) S. Ozaki, Chem. Rev. 1972, 72, 457–496; b) H. Ulrich, Chemistry and Technology of Isocyanates, Wiley, New York, 1996.
- [4] For seminal stoichiometric studies of Ni⁰ complexes with isocyanates, see: a) H. Hoberg, J. Organomet. Chem. 1988, 358, 507-517; b) H. Hoberg, K. Summermann, A. Milchereit, J. Organomet. Chem. 1985, 288, 237-248; c) H. Hoberg, E. Hernandez, Angew. Chem. Int. Ed. Engl. 1985, 24, 961-962; Angew. Chem. 1985, 97, 987-988; d) J. F. Villa, H. B. Powell, Inorg. Chim. Acta 1979, 32, 199-204.
- [5] For selected Ni-catalyzed reactions of isocyanates with coupling partners other than organic halides, see: a) T. Miura, M. Morimoto, M. Murakami, J. Am. Chem. Soc. 2010, 132, 15836–15838; b) T. Ozawa, H. Horie, T. Kurahashi, S. Matsubara, Chem. Commun. 2010, 46, 8055–8057; c) B. R. D'Souza, J. Louie, Org. Lett. 2009, 11, 4168–4171; d) K. D. Schleicher, T. F. Jamison, Org. Lett. 2007, 9, 875–878; e) H. A. Duong, M. Cross, J. Louie, J. Am. Chem. Soc. 2004, 126, 11438–11439.
- [6] P. Braunstein, D. Nobel, Chem. Rev. 1989, 89, 1927-1945.
- [7] a) A. Correa, R. Martin, J. Am. Chem. Soc. 2014, 136, 7253– 7725; b) J.-C. Hsieh, C.-H. Cheng, Chem. Commun. 2005, 4554– 4556.
- [8] a) The Amide Linkage: Structural Significance in Chemistry, Biochemistry and Materials Science (Eds.: A. Greenberg, C. M. Breneman, J. F. Liebman), Wiley-Interscience, New York, 2000;
 b) J. S. Carey, D. Laffan, C. Thomson, M. T. Williams, Org. Biomol. Chem. 2006, 4, 2337–2347.
- [9] For selected elegant reviews on amide bond formation, see:
 a) V. R. Pattabiraman, J. W. Bode, *Nature* 2011, 480, 471-479;
 b) C. L. Allen, J. M. Williams, *Chem. Soc. Rev.* 2011, 40, 3405-3415;
 c) R. M. Lanigan, T. D. Sheppard, *Eur. J. Org. Chem.* 2013, 2013, 7453-7465;
 d) E. Valeur, M. Bradley, *Chem. Soc. Rev.* 2009, 38, 606-631.
- [10] For selected examples, see: a) G. Zhang, B. Gao, H. Huang, Angew. Chem. Int. Ed. 2015, 54, 7657-7661; Angew. Chem. 2015, 127, 7767-7771; b) K. Dong, X. Fang, R. Jackstell, G. Laurenczy, Y. Li, M. Beller, J. Am. Chem. Soc. 2015, 137, 6053-6058; c) C. Jiménez-Rodriguez, A. A. Núñez-Magro, T. Seidensticker, G. R. Eastham, M. R. L. Furst, D. J. Cole-Hamilton, Catal. Sci. Technol. 2014, 4, 2332-2339; d) H. Liu, N. Yan, P. J. Dyson, Chem. Commun. 2014, 50, 7848-7851; e) X. Fang, R. Jackstell, M. Beller, Angew. Chem. Int. Ed. 2013, 52, 14089-14093; Angew. Chem. 2013, 125, 14339-14343; f) T. Fukuyama, T. Inouye, I. Ryu, J. Organomet. Chem. 2007, 692, 685-690; g) I. Ryu, K. Nagahara, N. Kambe, N. Sonoda, S. Kreimerman, M. Komatsu, Chem. Commun. 1998, 1953-1954, and references therein.
- [11] Selected examples on using organometallic reagents with βhydrogen atoms. For RMgX, see: a) G. Schäfer, C. Matthey, J. W. Bode, Angew. Chem. Int. Ed. 2012, 51, 9173–9175; Angew.

Chem. **2012**, *124*, 9307–9310. For RLi, see: b) V. Pace, L. Castoldi, W. Holzer, *Chem. Commun.* **2013**, *49*, 8383–8385, and c) I. Coldham, S. Dufour, T. F. N. Haxell, J. J. Patel, G. Sanchez-Jimenez, *J. Am. Chem. Soc.* **2006**, *128*, 10943–10951, and references therein.

- [12] Recently, intermolecular catalytic hydrocarbamoylation reactions of noncyclic alkenes with formamides have been reported for forging aliphatic amide bonds possessing β-hydrogens. In all cases α-branched amides could not be achieved, obtaining exclusive linear selectivity. For examples, see: a) T. Seidensticker, M. R. L. Furst, R. Frauenlob, J. Vondran, E. Paetzold, U. Kragl, A. J. Vorhold, *ChemCatChem* 2015, *7*, 4085–4090; b) Y. Miyazaki, Y. Yamada, Y. Nakao, T. Hiyama, *Chem. Lett.* 2012, *41*, 298–300; c) S. Ko, H. Han, S. Chang, *Org. Lett.* 2003, *5*, 2687–2690.
- [13] For selected recent examples, see: a) Y. Liu, J. Cornella, R. Martin, J. Am. Chem. Soc. 2014, 136, 11212-11215; b) T. Moragas, J. Cornella, R. Martin, J. Am. Chem. Soc. 2014, 136, 17702-17705; c) X. Wang, Y. Liu, R. Martin, J. Am. Chem. Soc. 2015, 137, 6476-6479; d) X. Wang, M. Nakajima, R. Martin, J. Am. Chem. Soc. 2015, 137, 6924-8927.
- [14] P. D. Bailey, T. J. Mills, R. Pettecrew, R. A. Price, in *Compre*hensive Organic Functional Groups Transformation II, Vol. 5 (Eds.: A. R. Katritzky, R. J. K. Taylor), Elsevier, Oxford, 2005, pp. 201–294.
- [15] See the Supporting information for details.
- [16] L2 can be prepared in two steps and on a multigram scale following a slightly modified literature procedure: T. Kauffmann, J. König, A. Woltermann, *Chem. Ber.* 1976, 109, 3864– 3868.
- [17] For the early use of bipyridine and phenanthroline ligands in cross-coupling reactions of unactivated alkyl halides, see: a) J. Zhou, G. C. Fu, J. Am. Chem. Soc. 2004, 126, 1340–1341;
 b) D. A. Powell, T. Maki, G. C. Fu, J. Am. Chem. Soc. 2005, 127, 510–511.
- [18] Whereas L1 and L5 predominantly lead to homodimerization, significant amounts of β-hydride elimination and reduction products were observed when using L4.
- [19] Note, however, that **L3** turned out to be particularly efficient for aryl isocianates (see Scheme 3).
- [20] Full conversion to β -hydride elimination products was observed for **1a-I**. The observed reactivity of **1a-OTs** is in line with the ability of these substrates to couple with other heterocumulenes (see Ref. [13 a]).
- [21] Unlike the utilization of aliphatic isocyanates, equimolar amounts of aromatic isocyanates were critical to prevent the formation of considerable amounts of *N*-acylureas.
- [22] [(TMEDA)Ni(o-tolyl)Cl] turned out to be particularly suited for the coupling of *i*PrNCO, avoiding dimerization or trimerization pathways.
- [23] This hypothesis is reinforced by the significant inhibition observed when reacting 1a with 2a in the presence of radical scavengers such as TEMPO or BHT. The intermediacy of radical-type intermediates gains credence from the observation that the Ni-catalyzed reductive amidation of 6-bromohex-1-ene results in a linear relationship between acyclic and 5-exo-trig cyclization products at different Ni/L2 loadings.
- [24] For remarkable exceptions, see: a) X. Wang, S. Wang, W. Xue, H. Gong, J. Am. Chem. Soc. 2015, 137, 11562-11565; b) C. Zhao, X. Jia, X. Wang, H. Gong, J. Am. Chem. Soc. 2014, 136, 17645-17651.
- [25] Noncaged tertiary alkyl bromides provided traces of products.
- [26] A. K. Mahalingam, X. Wu, M. Alterman, *Tetrahedron Lett.* 2006, 47, 3051–3053.
- [27] E. Broughton, Environ. Health 2005, 4, 6.

www.angewandte.org

© 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!



- [28] The isolation of $Ni^0(L2)_2$ proved to be particularly difficult. Stoichiometric studies were performed with $Ni^0(L3)$ and $NiBr_2$ -(L3), as L3 proved superior for aromatic isocyanates.
- [29] CCDC 1481428 (7) and CCDC 1481429 (8) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- [30] β -hydride elimination, reduction, homodimerization, and the formation of *N*-acylureas account for the mass balance.
- [31] At present, we cannot rule out a significant contribution dealing with Ni¹ intermediates generated via single-electron transfer reduction mediated by Mn, see: a) E. Duñach, A. P. Esteves, M. J. Medeiros, S. Olivero, *New J. Chem.* 2005, *29*, 633–636; b) T. Fujihara, Y. Horimoto, T. Mizoe, F. B. Sayyed, Y. Tani, J. Terao, S. Sakaki, Y. Tsuji, *Org. Lett.* 2014, *16*, 4960–4963; c) M. L. Nadal, J. Bosch, J. M. Vila, G. Klein, S. Ricart, J. M. Moretó, *J. Am. Chem. Soc.* 2005, *127*, 10476–10477.
- [32] For selected comproportionation events en route to Ni¹ species, see: a) J. Cornella, E. Gómez-Bengoa, R. Martin, J. Am. Chem.

Soc. 2013, 135, 1997-2009; b) A. Velian, S. Lin, A. J. M. Miller,
M. W. Day, T. Agapie, J. Am. Chem. Soc. 2010, 132, 6296-6297;
c) V. B. Phapale, E. Buñuel, M. García-Iglesias, D. J. Cárdenas,
Angew. Chem. Int. Ed. 2007, 46, 8790-8795; Angew. Chem.
2007, 119, 8946-8951; d) G. D. Jones, J. L. Martin, C. McFarland, O. R. Allen, R. E. Hall, A. D. Haley, R. J. Brandon, T. Konovalova, P. J. Desrochers, P. Pulay, D. A. Vicic, J. Am. Chem.
Soc. 2006, 128, 13175-13183.

[33] Ni¹ species have been shown to rapidly react with heterocumulenes other than RNCO, see: F. S. Menges, S. M. Craig, N. Tötsch, A. Bloomfield, S. Ghosh, H.-J. Krüger, M. A. Johnson, *Angew. Chem. Int. Ed.* **2016**, *55*, 1282–1285; *Angew. Chem.* **2016**, *128*, 1304–1307.

Received: May 26, 2016 Published online: ■■ ■■, ■■■



Communications



avoids the use of stoichiometric and

sensitive organometallic reagents.

Communications



primary, secondary, and tertiary alkyl bromides with isocyanates gives access

to a wide range of aliphatic amides. The

6 www.angewandte.org

C 2016 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

Angew. Chem. Int. Ed. 2016, 55, 1-6

These are not the final page numbers!