## Accepted Manuscript

Anionic fluoroquinolones as antibacterials against biofilm-producing *Pseudo-monas aeruginosa* 

Timothy E. Long, Lexie C. Keding, Demetria D. Lewis, Michael I. Anstead, T. Ryan Withers, Hongwei D. Yu

PII: DOI: Reference:	S0960-894X(16)30012-9 http://dx.doi.org/10.1016/j.bmcl.2016.01.012 BMCL 23477
To appear in:	Bioorganic & Medicinal Chemistry Letters
Received Date: Revised Date: Accepted Date:	<ul><li>22 December 2015</li><li>5 January 2016</li><li>6 January 2016</li></ul>



Please cite this article as: Long, T.E., Keding, L.C., Lewis, D.D., Anstead, M.I., Ryan Withers, T., Yu, H.D., Anionic fluoroquinolones as antibacterials against biofilm-producing *Pseudomonas aeruginosa*, *Bioorganic & Medicinal Chemistry Letters* (2016), doi: http://dx.doi.org/10.1016/j.bmcl.2016.01.012

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# **ACCEPTED MANUSCRIPT**

### **Graphical Abstract**

To create your abstract, type over the instructions in the template box below. Fonts or abstract dimensions should not be changed or altered.





Bioorganic & Medicinal Chemistry Letters journal homepage: www.elsevier.com

### Anionic fluoroquinolones as antibacterials against biofilm-producing Pseudomonas aeruginosa

Timothy E. Long<sup>a,b\*</sup>, Lexie C. Keding<sup>b</sup>, Demetria D. Lewis<sup>a</sup>, Michael I. Anstead<sup>d</sup>, T. Ryan Withers<sup>e</sup>, and Hongwei D. Yu<sup>b,c,e</sup>

<sup>a</sup> Department of Pharmaceutical Science and Research, School of Pharmacy, Marshall University, Huntington, WV 25755

<sup>b</sup> Departments of Biochemistry and Microbiology, and <sup>c</sup>Pediatrics, Joan C. Edwards School of Medicine, Marshall University, Huntington, WV 25755

<sup>d</sup>Department of Pediatrics, College of Medicine, University of Kentucky, Lexington, KY 40506

<sup>e</sup>Progenesis Technologies, LLC, Huntington WV 25755

#### ARTICLE INFO

Article history: Received Received in revised form Accepted Available online

*Keywords:* fluoroquinolone Pseudomonas mucoid biofilm antibacterial

#### ABSTRACT

*Pseudomonas aeruginosa* is a common biofilm-forming bacterial pathogen implicated in diseases of the lungs. The extracellular polymeric substances (EPS) of respiratory *Pseudomonas* biofilms are largely comprised of anionic molecules such as rhamnolipids and alginate that promote a mucoid phenotype. In this paper, we examine the ability of negatively-charged fluoroquinolones to transverse the EPS and inhibit the growth of mucoid *P. aeruginosa*. Anionic fluoroquinolones were further compared with standard antibiotics via a novel microdiffusion assay to evaluate drug penetration through pseudomonal alginate and respiratory mucus from a patient with cystic fibrosis.

2009 Elsevier Ltd. All rights reserved.

*Pseudomonas aeruginosa* is an opportunistic bacterium that is frequently implicated in burn wound, bladder, and lower respiratory tract infections [1]. The Gram-negative aerobe is also the primary etiology of chronic pulmonary infections in cystic fibrosis (CF) due to its ability to evade host defenses within the accumulated mucus lining the airway of the lungs. When *Pseudomonas* colonizes the lower respiratory tract, neutrophils infiltrate the lungs and release reactive oxygen species (ROS) to suppress the infection. Exposure to ROS is thought to cause DNA-level mutations and activation of gene products that further protects *P. aeruginosa* from the immune system [2]. Among these are the enzymes and proteins involved in the biosynthesis and secretion of rhamnolipids [3] and alginate [4], two components of the extracellular polymeric substance (EPS) found in *Pseudomonas* biofilms of CF airways.

Like the sulfated and carboxylated components of mucus located in the lower respiratory tract, rhamnolipids and alginate possess an overall negative charged (Figure 1) [5]. In CF, the anionic environment of the lungs is amplified by DNA and Factin released from necrotic neutrophils. The physiochemical properties of this dense complex mixture of negatively charged biomolecules consequentially serves as a barrier against cationic and zwitterionic antimicrobials [6]. It was rationalized from here that antibiotics carrying a net -1 or -2 charge would more readily transverse these barriers and reach the bacteria entrenched within due to the lack of ionic interactions with the EPS and mucus. On this premise, we set forth to evaluate anionic fluoroquinolones as growth inhibitors of biofilm-producing *P. aeruginosa* and their ability to penetrate the alginate component of *Pseudomonas* EPS and CF respiratory mucus.



**Figure 1.** Structures of (a) rhamnolipids and (b) alginate in biofilm-producing *Pseudomonas aeruginosa*.

The initial series of anionic fluoroquinolones explored were derived from FDA-approved fluoroquinolones **1a-5a** (Figure 2)

\* Corresponding author. Tel.: +1-304-696-7393; fax: +1-304-696-7309; e-mail: longt@marshall.edu

1

### 2

## CCEPTED MANUSCRIPT

including ciprofloxacin (1a), a first-line treatment for urinary tract infections due to P. aeruginosa. The initial design of the anionic fluoroquinolones was to acetylate the C-7 amino residue of the quinolone ring to preclude zwitterion formation at physiological pH. The reactions were performed in dichloromethane with acetic anhydride (1.5 equiv.), Et<sub>3</sub>N (1.3-2.6 equiv.), and DMAP (0.1 equiv.) to afford the N-acetyl fluoroquinolones 1b-5b. The derivatives and parent compounds were then evaluated for antipseudomonal activity against the reference non-mucoid P. aeruginosa PAO1 and its mucoid derivative, PAO581. The genomes of both bacterial strains have been sequenced [7,8] with the mutation causing the stable production of EPS alginate previously defined [9].



(-)5b R = Ac

Figure 2: N-Protio fluoroquinolones and N-substituted derivatives 1-6. 

Table 1: Zones	of growth	inhibition (m	ım) against	non-mucoid	PAO1	and
mucoid PAO581	for fluoroc	uinolones 1-6	í. <sup>a</sup>			

strain	PAO1	PAO1	PAO581	PAO581
media	LB	PIA	LB	PIA
ciprofloxacin (1a)	30	24	38	26
gatifloxacin (2a)	29	21	31	23
lomefloxacin (3a)	23	12	29	15
moxifloxacin (4a)	24	10	31	14
pazufloxacin (5a)	25	17	30	20
pefloxacin (6)	18	12	19	14
1b	10	0	0	13
1c	18	(20) <sup>c</sup>	21	9
2b	0	0	0	0
3b	0	0	0	0
4b	0	0	0	0
5b	0	0	0	0

 $^a$  Compounds were tested as  $K^{\scriptscriptstyle +}$  salts at 10  $\mu M$  concentration in water.

<sup>b</sup> Pseudomonas isolation agar (PIA) is supplemented with 5% glycerol as a C source for mucoid production. Luria-Bertani (LB) agar lacks glycerol to promote the mucoid phenotype Partial inhibition.

The N-acetyl fluoroquinolones 1b-5b were tested by the conventional disc diffusion (Kirby Bauer) assay on LB and PIA

to compare the ability of each analog to inhibit bacterial viability on regular growth and mucoidy-promoting agar media, respectively (Table 1). The compounds were evaluated as the K<sup>+</sup> salts using 10 mm cellulose discs impregnated with 10 µM of drug. Following overnight air incubation at 37 °C, the plates were examined for zones of growth inhibition and it was discovered that acetylation of the C-7 amino residue negated the inhibitory activity for most of the derivatives against both PAO variants. In contrast, the unmodified fluoroquinolones proved to be active displaying moderate to strong growth inhibitory effects with a zone diameter range of 10-38 mm.

As expected, ciprofloxacin demonstrated the greatest antipseudomonal activity and its N-acetylated form 1b was the lone derivative in the series to inhibit growth. To further explore impact of lipophilicity on the activity, the N-formyl (1c, Figure 2) [10] and 10 additional N-acyl derivatives (1d-1m, Figure 3) of ciprofloxacin [11,12] were prepared for microbiological evaluation. The compounds were once again tested at 10  $\mu$ M as  $K^+$  salts and with the exception of the *N*-formyl analog 1c, each was found to have no inhibitory effects on Pseudomonas growth. It was concluded from the data that the anionic character and lipophilicity (clogP 1.96 - 3.98) furnished to analogs 1b-1m of ciprofloxacin (clogP 1.63) had adversely affected antimicrobial activity. These findings further suggested that an ionizable group at the C-7 position of the quinolone ring may be a pre-requisite for transport and/or gyrase binding in Pseudomonas as noted for the reduced efficacy of pefloxacin (6, Table 1).



Figure 3: N-Acyl fluoroquinolones 1d-1m.

A second series of ciprofloxacin analogs modified with an ionizable carboxylate substituent on the C-7 piperazine ring were subsequently prepared (Scheme 1) and evaluated. Compounds **1n-1p** were synthesized by heating ciprofloxacin (1 equiv.) with their respective cyclic anhydrides (1 equiv.) overnight in DMSO [13]. Activity comparisons were made with the polycationic aminoglycoside tobramycin and the carboxypenicillin ticarcillin, which like the analogs carries a net -2 charge.



Scheme 1: Synthesis of carboxyfluoroquinlones 1n-1p.

In this series, the dianionic *N*-phthalyl derivative **1p** was the only compound found to effectively inhibit the growth of

## ACCEPTED MANUSCRIPT

*Pseudomonas* (Table 2). This was a surprising finding on comparison with the inactive *N*-benzoyl analog **1i**. Moreover, a 53-58% reduction in zone size was observed for bacteria grown on the mucoidy-promoting PIA media in contrast to 20-32% for ciprofloxacin. When compared with tobramycin, the *N*-phthalyl analog exhibited similar activity while ticarcillin was found to be inactive suggesting that the PAO strains used in the study were harboring the genes for one or more penicillin-degrading  $\beta$ -lactamase (e.g., AmpC) [14].

Table 2: Zones of growth inhibitions (mm) for fluoroquinolones 1n-1p.<sup>a</sup>

				<b>_</b>
strain	PAO1	PAO1	PAO581	PAO581
media	LB	PIA	LB	PIA
1n	0	0	0	0
10	0	0	0	0
1p	23	11	31	13
tobramycin	18	13	20	17
ticarcillin	0	0	0	0

<sup>a</sup>Compounds were tested as  $K^+$  salts at 10  $\mu M$  concentrations in water.

With these results, an additional series of carboxy-fluoroquinolones were evaluated. The acetic and butyric acid derivatives 1v and 1w were prepared by acid hydrolysis of their parent methyl esters 1q and 1r, respectively (Scheme 2) [15]. Additional *N*-alkyl derivatives 1s-1u were synthesized from their respective alkyl bromides for comparison with the analogs and the *N*-methyl fluoroquinolone, pefloxacin **6** (Figure 2). The antipseudomonal activity for this series was found to be mostly limited to *N*-alkylated analogs 1r and 1s (Table 3). On comparison with pefloxacin, the methyl ester derivative 1r displayed similar activity against mucoid PAO581.



Scheme 2: Synthesis of N-alkyl fluoroquinolones 1q-1w.

Table 3: Zones of grow	th	inhibitions	(mm)	for	fluoroq	juinolone	s 1q-1	<b>1 w</b> . <sup>a</sup>	
	-								-

strain	PAO1	PAO1	PAO581	PAO581
media	LB	PIA	LB	PIA
1q	0	0	0	0
1r	16	6	17	9
1s	23	0	13	0
1t	0	0	0	0
1u	0	0	13	0
1v	0	0	0	0
1w	0	0	0	0
pefloxacin (6)	18	12	19	14

 $^a\text{Compounds}$  were tested as  $K^{\scriptscriptstyle +}$  salts at 10  $\mu M$  concentrations in water.

Minimum inhibitory and bactericidal concentrations were subsequently determined by broth microdilution for the two most active fluoroquinolones from the study, the *N*-formyl and *N*phthalyl substituted ciprofloxacin analogs **1c** and **1p**, respectively (Table 4). The carboxy-fluoroquinolone **1p** was found to be inferior to both ciprofloxacin and tobramycin in liquid culture. Likewise, the *N*-formyl analog **1c** proved to be two fold less effective at inhibiting the pseudomonal growth compared to ciprofloxacin and tobramycin. Of note, the MIC results (i.e., ciprofloxacin  $\approx$  tobramycin > 1c = 1p) did not correlate to the zone data (i.e., ciprofloxacin > 1p > tobramycin > 1c) performed on solid media. It is believed that the smaller zone sizes exhibited for the charged fluoroquinolones and tobramycin was a consequence of truncated diffusion through the uncharged agar media. These results prompted us to examine the influence of charge on the ability the antibiotics to penetrate through the alginate component of biofilm *Pseudomonas* EPS and CF respiratory mucus.

**Table 4:** Minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) for select compounds.<sup>a</sup>

concentration (inib)	<i>s)</i> for select co.	mpounds.		
strain	PAO1	PAO1	PAO581	PAO581
	MIC	MBC	MIC	MBC
1c	3.125	6.25	3.125	6.25
1p	3.125	6.25	3.125	12.5
ciprofloxacin	1.56	6.25	1.56	6.25
tobramycin	3.125	1.56	0.78	6.25

 $^a$  Compounds were tested as  $K^{\prime}$  salts in Luria-Bertani broth. Values reported in  $\mu M.$ 

To probe the penetrating ability of charged antibiotics through solid media, a novel microdiffusion assay was developed using centrifuge filter columns containing 1% PAO581 alginate [16] or CF sputum in Noble agar. As depicted in Figure 4, the filter columns containing the semi-permeable barriers were inserted into a 1.5 mL centrifuge tube containing a 5 x 10<sup>5</sup> inoculum of *P. aeruginosa* and 50 µL of 25 µM antibiotic was applied to the agar surface. The tubes were sealed and incubated overnight at 37 °C with shaking. The penetrating ability was assessed by calculating the inhibition of antibiotic activity from the OD<sub>600</sub> measurements and Equation 1 with Noble agar, an uncharged medium, representing 100% drug diffusion into the broth culture.



Figure 4: Microdiffusion assay design.

 Table 5: Inhibition of antibiotic activity using PAO581 in the microdiffusion assay.

strain	PAO581 alginate	CF sputum
1c	0.61	0.89
1p	0.80	0.49
ciprofloxacin	0.97	-0.33
tobramycin	0.98	0.0

n = 0 (drug bioavailability not affected)

n = 0 < to 1 (drug bioavailability decreased)

n < 0 (drug bioavailability increased)

The microdiffusion assay revealed that *Pseudomonas* alginate and the components in CF sputum likely modulate antibiotic penetration (Table 5). The data suggests that the negatively charged sugars in alginate (i.e., mannuronate, guluronate) reduces the EPS permeation of both charged and zwitterionic antibiotics through *Pseudomonas* biofilms. Based on the preliminary

#### 4

### **ACCEPTED MANUSCRIPT**

findings from this assay, the anionic fluoroquinolones **1c** and **1p** were less hindered by the alginate barrier compared to ciprofloxacin and tobramycin. Conversely, the constituents found in CF patient sputum did not appear to affect the penetration of these standard antibiotics. Follow up studies using sputum from different CF patient populations will be performed to further corroborate these findings.

In summary, select anionic derivatives of the fluoroquinolone ciprofloxacin were found to possess antipseudomonal activity against biofilm-producing P. aeruginosa. The C-7 position was highly sensitive to chemical modification that may have affected DNA gyrase binding and/or cellular entry. It was further ascertained by the microdiffusion assay that the anionic fluoroquinolones appeared to have less chemical interactions with the alginate constituent from the EPS of PAO581 than ciprofloxacin and tobramycin. Conversely, the data suggests that the standard antibiotics penetrate CF respiratory mucus more effectively than the anionic fluoroquinolones. As a hallmark of the CF pathology, the hypersecretion of respiratory mucus facilitates persistent infection of the lower airways by Pseudomonas and adversely affects drug bioavailability (Figure 5). Treatment efficacy with inhaled and systemic antibiotics is not only predicated by the chemical constituents in the secreted mucus and Pseudomonas EPS but also the physiochemical properties of the drugs. This study established that ciprofloxacin can be chemically modified to modulate permeation in bacterial EPS and respiratory mucus as discerned by the microdiffusion assay. If developed further, this novel in vitro method to evaluate drug penetration against various biological barriers could become useful tool to help guide antimicrobial therapy in patients with chronic pulmonary disease.



Figure 5: Respiratory mucus and EPS of *Pseudomonas* biofilms reduce the bioavailability of inhaled and systemic antibiotics.

#### Acknowledgements

This research was supported in part by the Marshall University School of Pharmacy FRS Grant Program and the Office of Experiential Learning for financial support. Special thanks is also given to NASA West Virginia EPSCoR grant program, the National Center for Research Resources, the National Center for Advancing Translational Sciences, and National Institutes of Health (NIH), through Grant UL1TR000117. HDY is supported by NIH P20GM103434 to the West Virginia IDeA Network for Biomedical Research Excellence, and is the co-founder of Progenesis Technologies, LLC. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

#### **References and notes**

- 1. Driscoll, J. A.; Brody, S. L.; Kollef, M. H. Drugs 2007, 67, 351
- 2. Ciofu, O.; Riis B.; Pressler T.; Poulsen H. E.; Høiby, N. Antimicrob. Agents. Chemother. 2005, 49, 2276
- Vinckx, T.; Wei, Q.; Matthijs, S.; Cornelis, P. *Microbiology* 2010, 156, 678
- Mathee, K.; Ciofu, O.; Sternberg, C.; Lindum, P. W.; Campbell, J. I.; Jensen, P.; Johnsen, A. H.; Givskov, M.; Ohman, D. E.; Molin, S.; Høiby, N.; Kharazmi, A. *Microbiology* 1999, 145, 1349
- 5. Mann, E. E.; Wozniak, D. J. FEMS Microbiol. Rev. 2012, 36, 893
- Hatch, R. A.; Schiller, N. L. Antimicrob. Agents Chemother. 1998, 42, 974
- Stover, C. K.; Pham, X. Q.; Erwin, A. L.; Mizoguchi, S. D.; Warrener, P.; Hickey, M. J.; Brinkman, F. S.; Hufnagle, W. O.; Kowalik, D. J.; Lagrou, M.; Garber, R. L.; Goltry, L.; Tolentino, E.; Westbrock-Wadman, S.; Yuan, Y.; Brody, L. L.; Coulter, S. N.; Folger, K. R.; Kas, A.; Larbig, K.; Lim, R.; Smith, K.; Spencer, D.; Wong, G. K.; Wu, Z.; Paulsen, I. T.; Reizer, J.; Saier, M. H.; Hancock, R. E.; Lory, S.; Olson, M. V. Nature 2000, 406, 959
- Yin, Y.; Withers, T. R.; Govan, J. R.; Johnson, S. L.; Yu ,H. D. Genome Announc. 2013, 1, e00834-13
- Qiu, D.; Eisinger, V. M.; Head, N. E.; Pier, G. B.; Yu, H. D. Microbiology 2008, 154, 2119
- Vavříková, E.; Polanc, S.; Kočevar, M.; Horváti, K.; Bosze, S.; Stolaříková, J.; Vávrová, K.; Vinšová. J. Eur. J. Med. Chem. 2011, 46, 4937
- 11. Ji, C.; Miller, P. A.; Miller, M. J. ACS Med. Chem. Lett. 2015, 6, 707
- Cormier, R.; Burda, W. N.; Harrington, L.; Edlinger, J.; Kodigepalli, K. M.; Thomas, J.; Kapolka, R.; Roma, G.; Anderson, B. E.; Turos, E.; Shaw, L. N. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 6513
- Noël, S.; Gasser, V.; Pesset, B.; Hoegy, F.; Rognan, D.; Schalk, I. J.; Mislin, G. L. Org. Biomol. Chem. 2011, 9, 8288
- Hengzhuang, W.; Ciofu, O.; Yang, L.; Wu, H.; Song, Z.; Oliver, A.; Høiby, N. Antimicrob. Agents Chemother. 2013, 57, 196
- Fardeau, S.; Dassonville-Klimpt, A.; Audic, N.; Sasaki, A.; Pillon, M.; Baudrin, E.; Mullié, C.; Sonnet, P. *Bioorg. Med. Chem.* 2014, 22, 4049
- 16. Damron, F. H.; Yu, H. D. J. Bacteriol. 2011, 193, 286