**JOC** The Journal of Organic Chemistry

## Article

## **Convergent Total Synthesis of Lamellarins and Their Congeners**

Daiki Morikawa, Kazuki Morii, Yuto Yasuda, Atsunori Mori, and Kentaro Okano J. Org. Chem., Just Accepted Manuscript • DOI: 10.1021/acs.joc.0c00998 • Publication Date (Web): 28 May 2020 Downloaded from pubs.acs.org on May 30, 2020

## **Just Accepted**

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

## **Convergent Total Synthesis of Lamellarins and Their Congeners**

Daiki Morikawa,<sup>†</sup> Kazuki Morii,<sup>†</sup> Yuto Yasuda,<sup>†</sup> Atsunori Mori,<sup>†,‡</sup> and Kentaro Okano<sup>\*,†</sup>

<sup>†</sup>Department of Chemical Science and Engineering, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan <sup>‡</sup>Research Center for Membrane and Film Technology, Kobe University, 1-1 Rokkodai, Nada, Kobe 657-8501, Japan

Supporting Information Placeholder



**ABSTRACT:** A convergent total synthesis of lamellarins S and Z is described. The synthesis features a halogen dance of an easily accessible  $\alpha$ , $\beta$ -dibromopyrrole promoted by an ester moiety. The resultant  $\beta$ , $\beta$ '-dibromopyrrole undergoes a ligand-controlled Suzuki-Miyaura coupling to provide a range of diarylated pyrrole derivatives. The established synthetic method was also applicable to the synthesis of ningalin B, and lukianols A and B.

## **INTRODUCTION**

The number of pyrrole alkaloids have continued to grow over recent years.<sup>1</sup> Among them, more than 50 lamellarins have been reported<sup>2</sup> since lamellarins A-D were isolated in 1985 by Faulkner and Clardy from the marine prosobranch mollusks.<sup>3</sup> Ningalins, lukianols, didemnimides, dictyodendrins, polycitrins, and storniamides are also part of this class, as related pyrrole alkaloids, which possess various aryl groups at the two  $\beta$  posi



Figure 1. Lamellarins and the related pyrrole alkaloids.

tions of the pyrrole ring (Figure 1).<sup>2,4</sup> In addition to their structural diversity, their wide range of biological activities attract considerable interest as synthetic targets.<sup>5</sup> The common key feature of the synthesis of these pyrrole alkaloids is construction of the polysubstituted pyrrole core. Most of the reported synthetic methods use a symmetrical pyrrole diester or diarylacetylene as the synthetic intermediates, and most of the syntheses focus on symmetrical natural products.<sup>2,6</sup> To the best of our knowledge, there are only a limited number of synthetic methods that can be applied to a series of these unsymmetrical lamellarins.<sup>7</sup> Herein, we describe an approach toward a divergent synthesis of symmetrical/unsymmetrical lamellarins and their congeners, featuring a halogen dance<sup>8,9</sup> of a dibromopyrrole derivative.

The major challenge toward accomplishing a convergent total synthesis of lamellarins and their congeners is the construction of a multiply arylated pyrrole. We describe an inventive method for introducing the desired functional groups onto the pyrrole in a straightforward and divergent synthesis, superior to the classical Parr-Knorr synthesis and Hantzsch syntheses for this purpose.<sup>10</sup> Recently we have been investigating the synthetic potential of a halogen dance for the regiocontrolled and step-economical synthesis of tri- and tetra-substituted thiophenes and furans (Scheme 1A)." Differential reactivity of the bromo groups enabled a one-pot double Suzuki-Miyaura coupling, and afforded a triarylated thiophene in a regioselective manner. In contrast, attempted regioselective cross-coupling reaction of 2,3,5-tribromothiophene was difficult, and the same compound was afforded in low yield.<sup>12</sup> In addition to the superior regioselectivity, this reaction allows migration of the bromo group which can be used as a handle for further transformation. This enables a highly efficient atom-economical process, and importantly, also avoids late-stage bromination, which often causes side reactions in the presence of highly reactive electron-rich aromatic rings such as pyrrole. A halogen dance with a bromopyrrole derivative has not yet been reported, possibly due to the lower acidity of the pyrrole compared with thiophene and furan. We started the synthetic studies toward the pyrrole alkaloids based on our recent observation that an ester group significantly promoted the halogen dance (Scheme 1B).<sup>13</sup> In preliminary experiments, we investigated the regioselective metalation of tribromopyrrole 1 to obtain dibromopyrrole 2 as a common intermediate for the synthesis of these pyrrole alkaloids through regioselective arylation; however, the use of several organolithiums or Grignard reagents provided 2 as a major product, but with several inseparable byproducts. We anticipated that the readily available dibromopyrrole 3 could be converted into the desired 2 by protonation of organolithium 5, derived from the firstgenerated organolithium 4 through a halogen dance.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60



**Scheme 1.** Strategy for the convergent synthesis of lamellarins and their congeners.

## **RESULTS AND DISCUSSION**

The synthesis began with the regioselective dibromination of the commercially available pyrrole carboxylic acid ethyl ester (6) (Scheme 2). Protection of the pyrrole nitrogen with a 2-(trimethylsilyl)ethoxymethyl (SEM) group provided  $\alpha$ , $\beta$ -dibromopyrrole 7 for the key halogen dance step. Treatment with LDA at -78 ° C resulted in the

smooth migration of the α-bromo group within 10 min, and aqueous workup provided  $\beta$ , $\beta$ '-dibromopyrrole **8** in 82% yield on a multi-gram scale. Based on our preliminary experiments, two bromo groups and an ester moiety are essential for this halogen dance of a bromopyrrole derivative.<sup>14</sup> Both the reaction temperature and the reaction time proved critical for a high-yielding process, due to the instability of either β-lithio-α,β-dibromopyrrole **9** or α-lithio-β,β'-dibromopyrrole **10**.<sup>15</sup>



**Scheme 2.** Halogen dance of  $\alpha$ , $\beta$ -dibromopyrrole 7.

With  $\beta$ , $\beta$ '-dibromopyrrole **8** in hand, we then explored suitable palladium catalysts for the regioselective Suzuki-Miyaura coupling with two equivalents of arylboronate ester **11** (Table 1). First, the reaction with Pd(PPh<sub>3</sub>)<sub>4</sub> furnished monoarylated pyrrole **12**<sup>16</sup> and diarylated pyrrole **13** in 13% and 52% yields, respectively (entry 1). This monoarylation proceeded in a completely regiospecific manner, despite the expected difficulty in controlling the reactivities of two bromo groups at the  $\beta$  positions.<sup>17</sup> A combination of Pd<sub>2</sub>(dba)<sub>3</sub> (5 mol%) and PPh<sub>3</sub> (20 mol%) led to the formation of a 1:1 mixture of **12** and **13** with 36% recovery of the starting dibromopyrrole **8** (entry 2). Prolonged reaction time resulted in significant reduction of the yields of the products, due to hydrolysis of the ester moiety. The use of Pd(PCy<sub>3</sub>)<sub>2</sub> and Pd(*t*-Bu<sub>3</sub>P)<sub>2</sub> provi-

## Table 1. Effects of phosphine ligands<sup>a</sup>



<sup>*a*</sup> Reaction conditions: dibromopyrrole **8** (1.0 equiv, 0.30 mmol), **11** (2.0 equiv, 0.60 mmol), Pd source (0.030 mmol for palladium, 10 mol%), phosphine ligand (0.060 mmol, 20 mol%) Ba(OH)<sub>2</sub>·8H<sub>2</sub>O (6.0 equiv, 1.8 mmol), 1,4-dioxane/H<sub>2</sub>O (4:1), reflux, 15 min. <sup>*b*</sup> The yield

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

60

was determined from the <sup>1</sup>H NMR spectrum of the crude product using 1,1,2,2-tetrachloroethane as an internal standard. <sup>c</sup> Not detected in the crude <sup>1</sup>H NMR spectrum. <sup>d</sup> Arylboronate ester **11** (1.0 equiv, 0.30 mmol).

ded diarylated pyrrole 13 in 63% and 75% yields, respectively (entries 3 and 4). Encouraged by this superior catalytic activity, we performed the same reaction with one equivalent of arylboronate ester 11 in order to obtain monoarylated pyrrole 12; however,  $Pd(t-Bu_3P)_2$  favored the exclusive formation of 13 with 21% recovery of the starting dibromopyrrole 8 (entry 5). These results can be rationalized by "intramolecular catalyst transfer",<sup>18</sup> which has been observed in a Ni-/Pd-catalyzed living chaingrowth polymerization (Scheme 3). The complex 14 formed after transmetalation is converted to palladium (o)- $\pi$  complex 15. Dissociation of the  $\pi$  complex 15 gives the monoarylated pyrrole 12. In contrast, the palladium (o) in 15 is transferred to the distal C-Br bond through oxidative addition to form palladium (II) species 16. Subsequent Suzuki - Miyaura coupling furnishes the diarylated pyrrole 13. We next examined the effects of an additive to suppress the catalyst transfer. Apart from the recent report by Yokozawa's group,<sup>18e</sup> stoichiometric amount of Ph<sub>2</sub>S did not suppress the formation of the diarylated pyrrole,<sup>19</sup> whereas tetrahydrothiophene (THT) reduced the catalytic activity (entries 6 and 7). Additional



**Scheme 3.** Potential catalyst transfer mechanism of the exclusive formation of the diarylated pyrrole.

PhSMe and 1,5-cyclooctadiene<sup>20</sup> did not provide satisfactory results. Among a series of phosphine ligands tested,  $P(4-CF_3-C_6H_4)_3$  led to the formation of monoarylated pyrrole **12** in 36% yield (entry 8); however, more electrondeficient  $P[3,5-(CF_3)_2C_6H_3]_3$  provided neither **12** nor **13** with 85% recovery of the starting dibromopyrrole **8** (entry 9).

In this Suzuki-Miyaura cross coupling the use of Ba(OH)<sub>2</sub> proved effective for the smooth arylation; however, the strongly basic conditions led to a significant reduction of the yield of the products after prolonged reaction time, probably due to hydrolysis of the ethyl ester. This limitation prohibited our further optimization to improve the yield of monoarylated pyrrole 12. In addition, laborious purification process was required to obtain the monoarylated pyrrole 12 in a pure form on a preparative scale. The monoarylated pyrrole 12 was separated from the diarylated pyrrole 13 by column chromatography. Subsequent preparative SEC-HPLC was necessary for separating the monoarylated pyrrole 12 from a minute amount of an unidentified byproduct. During the process, substantial amount of the monoarylated pyrrole was not recovered, which resulted in the low isolated yield (22%) in Scheme 4) compared to the NMR yield (36% in Table 1).

This catalyst screening enabled a stepwise and regiocontrolled introduction of two aryl groups, which led to the synthesis of lukianols  $A^{6c,21}$  and  $B^{21a,21g}$  (Scheme 4). Similar to the report by Banwell,<sup>7a</sup> dibromopyrrole 8 underwent a one-pot double Suzuki-Miyaura coupling with catalytic  $Pd(PPh_3)_4$ , providing the desired product 13 in 78% yield. Deprotection of the SEM group was performed in a stepwise manner,<sup>22</sup> namely, TFA-promoted removal of the (trimethylsilyl)ethyl group and subsequent treatment of the resulting hemiaminal with aqueous sodium carbonate in the presence of hydroxylamine as a formaldehyde scavenger. The unprotected pyrrole underwent Nalkylation with 17 to provide the desired product 18, which was then transformed to the fused pyrrole 19 through hydrolysis of the ester by KOSiMe<sub>3</sub><sup>23</sup> followed by cyclization.<sup>21b</sup> Removal of the benzyl groups with a combination of BCl<sub>3</sub> and C<sub>6</sub>HMe<sub>5</sub><sup>24</sup> provided lukianol A. The established stepwise arylation was effective for the synthesis of lukianol B bearing different aromatic rings. The trimethylsilyl group of the obtained diarylated pyrrole 21 was converted to an iodo group by ICl at  $-78^{\circ}$  C, without affecting the reactive pyrrole ring. Deprotection of the SEM group, lactone formation, and removal of the benzyl groups completed the synthesis of lukianol B. This catalyst-controlled arylation strategy allows introduction of two different aryl groups in a completely regioselective manner. This proved effective for the synthesis of these pyrrole alkaloids, as a late-stage iodination of lukianol A was not regioselective due to the multiple phenol moieties and reactive pyrrole  $\alpha$ -position.<sup>21g</sup>



Scheme 4. Regiocontrolled synthesis of lukianol A and B.

The  $\beta$ ,  $\beta$ '-dibromopyrrole **8** was also the common intermediate for the synthesis of ningalin B<sup>6f,6h,25</sup>, and lamellarins S<sup>26</sup> and Z<sup>27</sup> (Scheme 5). A one-pot double Suzuki-Miyaura coupling with arylboronate ester 23 took place to give the corresponding pyrrole 24 in 59% yield. Hydrolysis of the ester and Pb(OAc)<sub>4</sub>-mediated lactone formation proceeded smoothly. Removal of the SEM group was carried out to provide the compound 25 in 24% yield over three steps from compound 24. N-Alkylation was performed with phenylethyl alcohol 26 under Mitsunobu conditions, and subsequent hydrogenolysis of the six benzyl ethers was carried out to provide ningalin B in 97% yield. The lactone-fused pyrrole 25 was also a synthetic intermediate for the total synthesis of lamellarin S. Thus, *N*-alkylation was carried out with phenylethyl alcohol 27 under the standard Mitsunobu conditions to provide compound 28 in 71% yield. Subsequent treatment with PIFA<sup>28</sup> achieved the oxidative C-C bond formation to construct the lamellarin skeleton. Hydrogenolysis of the benzvl ethers gave lamellarin S in 98% yield. In addition to these natural products, lamellarin Z, which has different aryl groups at the two  $\beta$  positions, was synthesized using the established stepwise arylation from the synthesis of lukianol B. The optimal monoarylation conditions were effective for arylboronate ester 29, and the corresponding monoarylated pyrrole 30 was isolated in 30% yield, with the excellent regioselectivity. The second arylation was performed with arylboronate ester 23 in the presence of catalytic  $Pd(PPh_3)_4$  to provide the desired compound **31** in moderate yield. Similar to the synthetic route of lamellarin S, the ethyl ester was then converted to the lactone. Removal of the SEM group and subsequent N-alkylation took place smoothly with phenylethyl alcohol 27 to provide the desired compound 32. The PIFA-promoted oxidative formation of the lamellarin skeleton and subsequent palladium-catalyzed hydrogenolysis of the benzyl ethers provided lamellarin Z in 80% yield over two steps from compound **32**.

#### CONCLUSIONS

In summary, we have achieved the convergent total synthesis of ningalin B, lukianols A and B, and lamellarins S and Z, by developing an unprecedented ester-promoted halogen dance of the  $\alpha$ , $\beta$ -dibromopyrrole derivative. The two bromo groups at the  $\beta$  positions were converted to the corresponding aryl groups by a regioselective Suzuki-Miyaura coupling. Significant ligand effects were observed for the ratio of monoarylated and diarylated products, which paved the way for the straightforward syntheses of the related pyrrole alkaloids.

## EXPERIMENTAL SECTION

### GENERAL

Analytical thin layer chromatography (TLC) was performed on Merck 60  $F_{254}$  aluminum sheets precoated with a 0.25 mm thickness of silica gel. Melting points (m.p.) were measured on a Yanaco MP-J3 and are uncorrected. Infrared (IR) spectra were recorded on a Bruker Alpha with an ATR attachment (Ge) and are reported in wave numbers (cm<sup>-1</sup>). <sup>1</sup>H NMR (400 MHz) and <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz) spectra were measured on a JEOL ECZ400 spectrometer. Chemical shifts for <sup>1</sup>H NMR are reported in parts per million (ppm) downfield from tetramethylsilane with the solvent resonance as the internal standard (CHCl<sub>3</sub>:  $\delta$  7.26 ppm, THF- $d_7$ :  $\delta$  1.72 ppm, DMSO- $d_5$ :  $\delta$  2.50 ppm, CHD<sub>2</sub>OD:  $\delta$  3.31 ppm, acetone- $d_5$ :  $\delta$  2.05 ppm, tetramethylsilane:  $\delta$  o ppm) and coupling constants are in Hertz

60

1

2

3

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60

#### The Journal of Organic Chemistry



Scheme 5. Convergent synthesis of lamellarins and their congeners.

(Hz). The following abbreviations are used for spin multiplicity: s = singlet, d = doublet, t = triplet, q = quartet, sept = septet, m = multiplet, and br = broad. Chemical shifts for <sup>13</sup>C{<sup>1</sup>H} NMR are reported in ppm from tetramethylsilane with the solvent resonance as the internal standard (CDCl<sub>3</sub>:  $\delta$ 77.16 ppm, THF-*d*s:  $\delta$  67.21 ppm, DMSO-*d*6:  $\delta$  39.52 ppm, CD<sub>3</sub>OD:  $\delta$  49.00 ppm, acetone-*d*6:  $\delta$  29.84 ppm). Highresolution mass spectra (HRMS) were performed on a JEOL JMS-TiooLP AccuTOF LC-Plus (ESI) with a JEOL MS-5414DART attachment.

## MATERIALS

Unless otherwise stated, all reactions were conducted in flame-dried glassware under an inert atmosphere of nitrogen. All work-up and purification procedures were carried out with reagent solvents in air. Unless otherwise noted, materials were obtained from commercial suppliers and used without further purification. Flash column chromatography was performed on Wakogel 60N (45-75 µm, Wako Pure Chemical Industries, Ltd.). Recycling preparative SEC-HPLC was performed with LC-9201 (Japan Analytical Industry Co., Ltd.) equipped with preparative SEC columns (JAI-GEL-1H and JAI-GEL-2H). Anhydrous THF (>99.5%, water content: < 10 ppm) was purchased from Kanto Chemical Co., Inc. and further dried by passing through a solvent purification system (Glass Contour) prior to use. LDA (2.0 M in THF/heptane /ethylbenzene) was purchased from Sigma-Aldrich Co. (Product number: 361798).

## SYNTHESIS OF DIPROMOPYRROLE 8

#### Ethyl 4,5-dibromo-1*H*-pyrrole-2-carboxylate (S1).

A 1-L round-bottomed flask equipped with a Teflon-coated magnetic stirring bar was charged with ethyl 1*H*-pyrrole-2-carboxylate (6) (11.20 g, 80.5 mmol, 1.0 equiv) and CHCl<sub>3</sub> (400 mL). After the solution was cooled to  $0^{\circ}$  C, Br<sub>2</sub> (25.56 g, 160 mmol, 2.0 equiv) in CHCl<sub>3</sub> (8 mL) was added dropwise to the flask over 5 min. The reaction mixture was allowed to warm

to room temperature with stirring over 5 h, at which time the mixture was treated with saturated aqueous sodium thiosulfate (200 mL). After partitioned, the aqueous layer was extracted with CHCl<sub>3</sub> (200 mL). The organic extracts were washed with water (200 mL) and brine (200 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was recrystallized from hexane to provide the title compound S1 as a colorless solid (19.00 g, 64.0 mmol, 80%), whose <sup>1</sup>H NMR spectra was identical to that reported in the literature.<sup>29</sup>  $R_f =$ 0.21 (hexane/diethyl ether = 6:1); M.p. 111-112  $^{\circ}$  C; IR (ATR, cm<sup>-1</sup>): 3230, 2925, 1694, 1412, 1385, 1319, 1235, 1205, 1022, 973, 830, 759, 637, 622; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.45 (br s, 1H), 6.88 (d, 1H, J = 3.2 Hz), 4.33 (q, 2H, J = 7.2 Hz), 1.36 (t, 3H, I = 7.2 Hz; <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  160.5, 124.0, 118.0, 107.6, 100.5, 61.5, 14.5; HRMS (DART/TOF) m/z: [M +  $H^{+}_{1}$  Calcd for  $C_{7}H_{8}^{79}Br^{81}BrNO_{2}$  297.8901; Found 297.8905.

#### Ethyl 4,5-dibromo-1-[[2-(trimethylsilyl)ethoxy]methyl]-1*H*-pyrrole-2-carboxylate (7).

A flame-dried 500-mL Schlenk tube equipped with a Tefloncoated magnetic stirring bar and a rubber septum was charged with S1 (3.715 g, 15.0 mmol, 1.0 equiv) and anhydrous THF (50 mL). After the solution was cooled to 0  $^{\circ}$  C, SEMCl (3.367 g, 20.2 mmol, 1.3 equiv) and NaH (771.9 mg, 19.3 mmol, 1.3 equiv) were added to the Schlenk tube at 0  $^{\circ}$  C. The reaction mixture was allowed to warm to room temperature with stirring over 80 min, at which time the mixture was treated with water (50 mL). The resulting mixture was extracted twice with diethyl ether (70 mL). The combined organic extracts were washed with water (100 mL) and brine (100 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 1:1) to provide the corresponding product 7as a brown oil (4.517 g, 10.6 mmol, 71%).  $R_f = 0.78$  (hexane/diethyl ether = 6:1; IR (ATR, cm<sup>-1</sup>): 1712, 1415, 1315, 1220,

Page 6 of 18

1094, 859, 836, 757, 635; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.06 (s, 1H), 5.83 (s, 2H), 4.29 (q, 2H, *J* = 7.2 Hz), 3.57 (t, 2H, *J* = 8.2 Hz), 1.34 (t, 3H, *J* = 7.2 Hz), 0.90 (t, 2H, *J* = 8.2 Hz), -0.03 (s, 9H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  159.5, 124.5, 120.5, 113.4, 100.5, 75.3, 66.1, 60.6, 17.8, 14.4, -1.4; HRMS (ESI/TOF) *m/z*: [M + Na]<sup>+</sup> Calcd for C<sub>13</sub>H<sub>21</sub><sup>79</sup>Br<sup>81</sup>BrNO<sub>3</sub>SiNa 449.9535; Found 449.9520.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

## Ethyl 3,4-dibromo-1-[(2-(trimethylsilyl)ethoxy)methyl]-1*H*-pyrrole-2-carboxylate (8).

A flame-dried 500-mL Schlenk tube equipped with a Tefloncoated magnetic stirring bar and a rubber septum was charged with 7 (3.66 g, 8.56 mmol, 1.0 equiv) and anhydrous THF (86 mL). After the solution was cooled to ~78  $^\circ~$  C, LDA (2.0 M in THF/heptane/ethylbenzene, 4.7 mL, 9.4 mmol, 1.1 equiv) was added to the Schlenk tube and stirred at -78  $^{\circ}$  C for 10 min. The reaction mixture was treated with water (2 mL) at -78  $^{\circ}$  C and allowed to warm to room temperature with stirring over 1 h, at which time the resulting mixture was treated with saturated aqueous ammonium chloride (90 mL) and extracted twice with diethyl ether (100 mL). The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/ $CH_2Cl_2 = 1:1$ ) to provide the title product 8 as a yellow oil (2.99 g, 7.01 mmol, 82%).  $R_f = 0.32$  (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 1:1); IR (ATR, cm<sup>-1</sup>): 3118, 1692, 1394, 1375, 1312, 1281, 1248, 1168, 1096, 1081, 834; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.06 (s, 1H), 5.63 (s, 2H), 4.36 (q, 2H, J = 7.6 Hz), 3.51 (t, 2H, J = 8.0 Hz), 1.40 (t,  $_{2H, J} = 7.6 \text{ Hz}$ , 0.90 (t,  $_{3H, J} = 8.0 \text{ Hz}$ ), -0.02 (s, 9H);  $_{^{13}}C{^{1}\text{H}}$ NMR (100 MHz, CDCl<sub>3</sub>): δ 159.7, 126.6, 121.0, 108.6, 101.6, 78.6, 66.8, 61.1, 17.9, 14.3, -1.3; HRMS (ESI/TOF) m/z: [M + Na]<sup>+</sup> Calcd for C<sub>13</sub>H<sub>21</sub><sup>79</sup>Br<sup>81</sup>BrNO<sub>3</sub>SiNa 449.9535; Found 449.9534.

## CATALYST SCREENING IN SUZUKI-MIYAURA COUPLING

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 8 (0.30 mmol, 1.0 equiv), 11 (0.60 mmol, 2.0 equiv),  $Ba(OH)_2 \cdot 8H_2O$  (1.8) mmol, 6.0 equiv), palladium catalyst (30 µmol for palladium, 10 mol%), phosphorus ligand (60 µmol, 20 mol%), water (0.6 mL), and 1,4-dioxane (2.4 mL). The flask was placed in a preheated oil bath and heated at 100 ° C for 15 min, at which time the reaction mixture was treated with water (3 mL). The resulting mixture was extracted with diethyl ether (5 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product. The yields of the monoarylated pyrrole 12 and diarylated pyrrole 13 were determined by <sup>1</sup>H NMR analysis using 1,1,2,2-tetrachloroethane as an internal standard by comparing relative values of integration for the peaks observed at 5.71 ppm (2 protons for 13), 5.66 ppm (2 proton for 12), and 5.63 ppm (2 proton for 8) with that of 1,1,2,2tetrachloroethane observed at 5.96 ppm.

## TOTAL SYNTHESIS OF LUKIANOL A

Ethyl 3,4-bis(4-(benzyloxy)phenyl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrrole-2-carboxylate (13). A 100-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 8 (438.5 mg, 1.03 mmol, 1.0 equiv), 11 (1.270 g, 4.09 mmol, 4.0 equiv), Ba(OH)2: 8H2O (1.946 g, 6.17 mmol, 6.0 equiv), Pd(PPh3)4 (115.7 mg, 0.100 mmol, 10 mol%), water (2 mL), and 1,4dioxane (8 mL). The flask was placed in a preheated oil bath and heated at 100  $^{\circ}\,$  C for 15 min, at which time the solution was treated with saturated aqueous ammonium chloride (30 mL). The resulting mixture was extracted twice with ethyl acetate (30 mL). The combined organic extracts were washed with water (50 mL) and brine (50 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/diethyl ether = 8:1 to 3:1, gradient) to provide the corresponding product 13 as a brown oil (511.0 mg, 0.806 mmol, 78%).  $R_f = 0.27$  (hexane/diethyl ether = 3:1); IR (ATR, cm<sup>-1</sup>): 2953, 1694, 1240, 1176, 1104, 1082, 834, 697; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.46-7.31 (m, 10H), 7.12 (d, 2H, J = 8.8 Hz), 7.10 (s, 1H), 7.00 (d, 2H, J = 8.8 Hz), 6.90 (d, 2H, J = 8.8 Hz), 6.81 (d, 2H, J = 8.8 Hz), 5.71 (s, 2H), 5.08 (s, 2H), 5.00 (s, 2H), 4.06 (q, 2H, J = 7.2 Hz), 3.62 (t, 2H, J = 8.0 Hz), 0.99-0.93 (m, 5H), 0.00 (s, 9H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 161.7, 157.6, 157.2, 137.2, 137.1, 131.8, 131.6, 129.3, 128.5, 128.3, 127.9, 127.5, 127.2, 125.4, 124.6, 120.2, 114.5, 114.0, 77.7, 69.9, 66.2, 59.8, 17.9, 13.8, -1.3; HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for  $C_{39}H_{43}NO_5SiNa$ 656.2808; Found 656.2831.

## Ethyl 3,4-bis(4-(benzyloxy)phenyl)-1*H*-pyrrole-2carboxylate (S2).

A 100-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 13 (115.8 mg, 0.183 mmol, 1.0 equiv) and  $CH_2Cl_2$  (4 mL). The solution was cooled to 0 ° C. TFA (100 µL, 1.31 mmol, 7.3 equiv) was added to the flask and the resulting mixture was stirred at 0 ° C for 4 h, at which time the reaction mixture was treated with water (8 mL) and extracted with  $CH_2Cl_2$  (10 mL) three times. The combined organic extracts were washed with saturated aqueous sodium hydrogen carbonate (30 mL), water (30 mL), and brine (30 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude hemiaminal (118.8 mg), which was used for the next step without further purification.

A 20-mL test tube equipped with a Teflon-coated magnetic stirring bar was charged with the crude hemiaminal, sodium carbonate (152.4 mg, 1.44 mmol, 7.9 equiv), NH<sub>2</sub>OH· HCl (13.7 mg, 0.21 mmol, 1.1 equiv), water (0.3 mL), and THF (1.2 mL). The resulting mixture was stirred at room temperature for 4.5 h, at which time the reaction mixture was treated with 1 M aqueous hydrochloric acid (2 mL) and extracted with ethyl acetate (4 mL) three times. The combined organic extracts were washed with water (20 mL) and brine (20 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/methyl acetate = 5:1) to provide the title product S<sub>2</sub> as a colorless solid (75.0 mg, 0.15 mmol, 82%).  $R_f = 0.50$  (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 1:2); M.p. 82-85 ° C; IR (ATR, cm<sup>-1</sup>): 3301, 1681, 1535, 1455, 1379, 1289, 1239, 1176, 1025, 833, 737, 698; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.12 (br s, 1H), 7.47-7.30 (m, 10H), 7.20 (d, 2H, J = 8.8 Hz), 7.05-7.03 (m, 3H), 6.92 (d, 2H, J = 8.8 Hz), 6.83 (d, 2H, J = 8.8 Hz), 5.07 (s, 2H), 5.01 (s, 2H), 4.19 (q, 2H, J

2

3

4

5

6

7

8

9

32

33

57 58 59

60

= 7.2 Hz), 1.17 (t, 3H, J = 7.2 Hz); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 161.3, 157.9, 157.4, 137.3, 137.2, 132.1, 129.6, 129.0, 128.7, 128.07, 128.05, 127.74, 127.65, 127.5, 126.9, 126.5, 120.0, 119.9, 114.7, 114.1, 70.1, 60.3, 14.3; HRMS (ESI/TOF) m/z: [M + Na]<sup>+</sup> Calcd for C<sub>33</sub>H<sub>29</sub>NO<sub>4</sub>Na 526.1994; Found 526.2018.

#### Ethvl 3,4-bis(4-(benzyloxy)phenyl)-1-(2-(4-(benzyloxy)phenyl)-2-oxoethyl)-1H-pyrrole-2-

carboxylate (18). A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with S2 (109.0 mg, 10 0.216 mmol, 1.0 equiv), 17 (93.0 mg, 0.305 mmol, 1.4 equiv), 11 K<sub>2</sub>CO<sub>3</sub> (59.9 mg, 0.433 mmol, 2.0 equiv), and DMF (720 µL). 12 After stirring at room temperature for 3 h, the reaction was 13 quenched with water (4 mL). The resulting mixture was washed twice with diethyl ether (4 mL). The aqueous layer 14 was extracted twice with ethyl acetate (4 mL). The combined 15 organic extracts were washed with brine (8 mL), dried over 16 sodium sulfate, and filtered. The filtrate was concentrated 17 under reduced pressure to provide the title product 18 as a 18 colorless solid (118.0 mg, 0.162 mmol, 75%), which was used 19 for the next step without further purification.  $R_f = 0.53$  (hex-20 ane/methyl acetate = 3:2); M.p. 152-153  $^{\circ}$  C; IR (ATR, cm<sup>-1</sup>): 21 2932, 1688, 1600, 1373, 1233, 1171, 1102, 1015, 834, 739, 697; <sup>1</sup>H 22 NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.02 (d, 2H, J = 8.8 Hz), 7.46-7.30 23 (m, 15H), 7.17 (d, 2H, J = 8.8 Hz), 7.06 (d, 2H, J = 8.8 Hz), 7.03 24 (d, 2H, J = 8.8 Hz), 6.92 (s, 1H), 6.90 (d, 2H, J = 8.8 Hz), 6.8025 (d, 2H, J = 8.8 Hz), 5.74 (s, 2H), 5.16 (s, 2H), 5.08 (s, 2H), 5.00 (s, 2H), 3.93 (q, 2H, J = 7.3 Hz), 0.85 (t, 3H, J = 7.3 Hz); <sup>13</sup>C{<sup>1</sup>H} 26 NMR (100 MHz, CDCl<sub>3</sub>): δ 191.9, 163.3, 162.0, 157.6, 157.2, 27 137.34, 137.25, 136.2, 132.1, 131.1, 130.5, 129.5, 128.9, 128.7, 128.4, 28 128.2, 128.04, 128.00, 127.7, 127.6, 127.5, 127.0, 124.6, 120.3, 115.1, 29 114.6, 114.0, 70.3, 70.1, 59.8, 55.6, 13.8; HRMS (ESI/TOF) m/z: 30  $[M + Na]^+$  Calcd for C<sub>48</sub>H<sub>41</sub>NO<sub>6</sub>Na 750.2832; Found 750.2850. 31

## 3,4-Bis(4-(benzyloxy)phenyl)-1-(2-(4-(benzyloxy)phenyl)-2-oxoethyl)-1H-pyrrole-2-carboxylic acid (S3).

34 A 10-mL screw-top test tube equipped with a Teflon-coated 35 magnetic stirring bar was charged with 18 (61.0 mg, 83.8 36 µmol, 1.0 equiv), potassium trimethylsilanolate (66.3 mg, 37 0.465 mmol, 5.6 equiv), and 1,2-dichloroethane (840 µL). The 38 test tube was placed in a preheated aluminum block and heated at 80  $^{\circ}\,$  C for 15 min. The reaction was quenched with 39 1 M aqueous hydrochloric acid (5 mL). The resulting mixture 40 was extracted with CHCl<sub>3</sub> (4 mL) three times. The combined 41 organic extracts were washed with brine (10 mL), dried over 42 sodium sulfate, and filtered. The filtrate was concentrated 43 under reduced pressure to give a crude product, which was 44 washed with the mixture of diethyl ether (4 mL) and hexane 45 (2 mL) to provide the title compound S<sub>3</sub> as a colorless solid 46 (42.2 mg, 60.3  $\mu$ mol, 72%). R<sub>f</sub> = 0.37 (hexane/methyl acetate = 47 3:2); M.p. 177-180 ° C; IR (ATR, cm<sup>-1</sup>): 2922, 1650, 1600, 1454, 48 1240, 1171, 1026, 835, 735, 697, 639; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.01 (d, 2H, J = 8.8 Hz), 7.46-7.30 (m, 16H), 7.06 (d, 49  $_{2H, J} = 8.8 \text{ Hz}$ , 7.00-6.96 (m, 6H), 6.80 (d, 2H, J = 8.8 Hz), 50 5.76 (s, 2H), 5.15 (s, 2H), 5.07 (s, 2H), 5.00 (s, 2H); <sup>13</sup>C{<sup>1</sup>H} 51 NMR (100 MHz, THF-d<sub>8</sub>): δ 191.8, 163.6, 163.1, 158.4, 157.9, 52 138.6, 138.5, 137.7, 132.8, 130.9, 130.7, 129.8, 129.6, 129.0, 128.9, 53 128.7, 128.5, 128.11, 128.07, 128.0, 127.7, 124.7, 121.0, 115.2, 114.8, 54 114.0, 70.5, 70.2, 56.0; HRMS (ESI/TOF) *m/z*: [M + Na]<sup>+</sup> Calcd 55 for C<sub>46</sub>H<sub>37</sub>NO<sub>6</sub>Na 722.2519; Found 722.2545. 56

### 3,7,8-Tris(4-(benzyloxy)phenyl)-1H-pyrrolo[2,1c][1,4]oxazin-1-one (19).

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with S<sub>3</sub> (65.4 mg, 93.5 µmol, 1.0 equiv), potassium acetate (162.5 mg, 1.66 mmol, 18 equiv), and acetic anhydride (9.4 mL). The flask was placed in a preheated oil bath and heated at 100  $^{\circ}$  C for 30 min. After cooling to room temperature, the acetic anhydride was removed azeotropically with toluene (60 mL) under reduced pressure. The oily residue was dissolved in ethyl acetate (10 mL) and the resulting mixture was washed with saturated aqueous ammonium chloride (5 mL). After partitioned, the aqueous layer was extracted twice with ethyl acetate (5 mL). The combined organic extracts were washed with water (10 mL) and brine (10 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/methyl acetate = 3:1) to provide the corresponding product 19 as a yellow amorphous  $(48.6 \text{ mg}, 71.3 \mu \text{mol}, 76\%)$ . R<sub>f</sub> = 0.59 (hexane/methyl acetate = 3:2); IR (ATR, cm<sup>-1</sup>): 1732, 1608, 1514, 1429, 1415, 1242, 1176, 1026, 834, 751, 697; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.66 (d, 2H, J = 9.2 Hz), 7.46-7.29 (m, 18H), 7.22 (s, 1H), 7.09 (d, 2H, J = 8.8 Hz), 7.04 (d, 2H, J = 8.8 Hz), 6.96 (d, 2H, J = 8.8 Hz), 6.88 (d, 2H, J = 8.8 Hz), 5.12 (s, 2H), 5.08 (s, 2H), 5.04 (s, 2H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 159.8, 158.4, 158.0, 154.5, 142.1, 137.2, 137.0, 136.7, 132.2, 130.0, 129.8, 128.8, 128.73, 128.71, 128.3, 128.2, 128.1, 127.8, 127.6, 126.2, 126.0, 125.1, 123.5, 119.1, 115.3, 114.9, 114.3, 113.1, 102.9, 70.2, 70.1; HRMS (DART/TOF) m/z:  $[M + H]^+$ Calcd for C46H36NO5 682.2594; Found 682.2608.

## Lukianol A.

A flame-dried 20-mL Schlenk tube equipped with a Tefloncoated magnetic stirring bar and a rubber septum was charged with 19 (23.7 mg, 34.8 µmol, 1.0 equiv), pentamethylbenzene (31.0 mg, 209 µmol, 6.0 equiv), and CH<sub>2</sub>Cl<sub>2</sub> (3.5 mL). The solution was cooled to -78  $^{\circ}$  C. BCl<sub>3</sub> (1 M in CH<sub>2</sub>Cl<sub>2</sub>, 209 µL, 0.21 mmol, 6.0 equiv) was added dropwise to the Schlenk tube. After stirring at -78  $^{\circ}$  C for 30 min, the reaction mixture was allowed to warm to 0  $^\circ\,\,$  C and stirred for 20  $\,$ min, at which time the mixture was treated with methanol (4 mL) and concentrated under reduced pressure. The residue was dissolved in methanol (3 mL), washed twice with hexane (2 mL), and concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/acetone = 1:1). The obtained product was washed with a mixture of acetone (2.3 mL) and hexane (0.75 mL) to provide lukianol A as a colorless solid (9.0 mg, 22 µmol, 63%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra (DMSO $d_6$ ) were identical to those reported in the literature.<sup>21a</sup> R<sub>f</sub> = 0.28 (hexane/acetone = 1:1); M.p. 210-214 ° C; IR (ATR, cm<sup>-1</sup>): 3321, 1703, 1612, 1544, 1537, 1519, 1504, 1421, 1259, 1236, 1203, 1174, 1042, 1023, 836, 759, 635, 608; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD): δ 7.84 (s, 1H), 7.61 (d, 2H, J = 8.4 Hz), 7.49 (s, 1H), 7.13 (d, 2H, J = 8.4 Hz), 7.01 (d, 2H, J = 8.4 Hz), 6.87 (d, 2H, J = 8.4 Hz), 6.74 (d, 2H, J = 8.4 Hz), 6.67 (d, 2H, J = 8.4 Hz);<sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>): δ 9.88 (s, 1H), 9.45 (s, 1H), 9.41 (s, 1H), 8.08 (s, 1H), 7.59 (s, 1H), 7.56 (d, 2H, I = 8.4 Hz), 7.05 (d, 2H, I =8.4 Hz), 6.95 (d, 2H, J = 8.4 Hz), 6.87 (d, 2H, J = 8.4 Hz), 6.70 (d, 2H, J = 8.4 Hz), 6.66 (d, 2H, J = 8.4 Hz); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, DMSO-*d*<sub>6</sub>): δ 158.4, 156.6, 156.3, 153.6, 140.8, 131.8, 129.5, 128.8, 127.4, 125.5, 123.9, 123.2, 121.3, 120.0, 115.8, 115.3, 114.6,

111.9, 103.1; HRMS (DART/TOF) m/z:  $[M + H]^+$  Calcd for  $C_{25}H_{18}NO_5$  412.1185; Found 412.1170.

## 1-Bromo-4-(phenylmethoxy)-benzene (S4).

A 500-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 4bromophenol (6.961 g, 40.2 mmol, 1.0 equiv), potassium carbonate (11.15 g, 81.0 mmol, 2.0 equiv), benzyl bromide (5.0 mL, 42.2 mmol, 1.05 equiv), and acetonitrile (40 mL). The flask was placed in a preheated oil bath and heated at reflux for 1 h. After cooling to room temperature, the resulting mixture was filtered through a pad of Celite. The filtrate was concentrated under reduced pressure to provide the corresponding product **S4** as a colorless solid (9.79 g, 37.2 mmol, 93%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra were identical to those reported in the literature.<sup>30</sup>

## 4,4,5,5-Tetramethyl-2-[4-(phenylmethoxy)phenyl]-1,3,2dioxaborolane (11).

A flame-dried 50-mL Schlenk tube equipped with a Tefloncoated magnetic stirring bar and a rubber septum was charged with S4 (1.00 g, 3.80 mmol, 1.0 equiv) and anhydrous THF (19 mL). After the solution was cooled to -78 ° C, n-BuLi (1.59 M in *n*-hexane, 2.62 mL, 4.18 mmol, 1.1 equiv) was added to the Schlenk tube. The reaction mixture was stirred at -78 ° C for 10 min. To the white suspension was added *i*-PrOBpin (850.0 mg, 4.57 mmol, 1.2 equiv) at  $-78^{\circ}$  C. After stirring at -78 ° C for 10 min, the resulting mixture was allowed to warm to room temperature with stirring over 30 min, at which time the reaction mixture was treated with water (20 mL). The mixture was extracted twice with diethyl ether (20 mL). The combined organic extracts were washed with brine (40 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/ $CH_2Cl_2 = 10:1$  to hexane/methyl acetate = 3:1, gradient) to provide the corresponding product 11 as a colorless solid (922.5 mg, 2.97 mmol, 78%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra were identical to those reported in the literature.31

## 2-Bromo-1-[4-(phenylmethoxy)phenyl]-ethanone (17).

A 100-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 1-[4-(benzyloxy)phenyl]-ethanone (452.6 mg, 2.00 mmol, 1.0 equiv), CuBr<sub>2</sub> (893.8 mg, 4.00 mmol, 2.0 equiv), CHCl<sub>3</sub> (4 mL), and ethyl acetate (4 mL). The flask was placed in a preheated oil bath and heated at 80 ° C for 3 h. After cooling to room temperature, the resulting mixture was filtered through a pad of Celite. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 3:1 to hexane/diethyl ether = 10:1, gradient) to provide the corresponding product 17 as a colorless solid (350.6 mg, 1.15 mmol, 57%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra were identical to those reported in the literature.<sup>32</sup>

## TOTAL SYNTHESIS OF LUKIANOL B

Ethyl 3-(4-(benzyloxy)phenyl)-4-bromo-1-((2-(trimethylsilyl)ethoxy)methyl)-1*H*-pyrrole-2-carboxylate (12). A 100-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 8 (427.0 mg, 1.00 mmol, 1.0 equiv), 11 (465.8 mg, 1.50 mmol, 1.5 equiv), Ba(OH)2· 8H2O (1.893 g, 6.00 mmol, 6.0 equiv), Pd2(dba)3 (45.9 mg, 50.1 µmol, mol%), tris[4-5 (trifluoromethyl)phenyl]phosphine (92.0 mg, 0.197 mmol, 20 mol%), water (2 mL), and 1,4-dioxane (8 mL). The flask was placed in a preheated oil bath and heated at 100  $^{\circ}\,$  C for 15 min, at which time the reaction was guenched with water (10 mL) and saturated aqueous ammonium chloride (10 mL). The resulting mixture was extracted with ethyl acetate (15 mL) three times. The combined organic extracts were washed with water (40 mL) and brine (40 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/diethyl ether = 10:1 to 5:1, gradient) followed by preparative SEC-HPLC to provide the title compound 12 as a colorless oil (118 mg, 0.222 mmol, 22%). Rf = 0.31 (hexane/diethyl ether = 3:1); IR (ATR, cm<sup>-1</sup>): 1699, 1288, 1246, 1179, 1100, 1084, 1018, 858, 833, 736, 697; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.46 (d, 2H, J = 7.6 Hz), 7.40 (dd, 2H, J = 8.0, 7.6 Hz), 7.33 (t, 1H, J = 8.0 Hz), 7.22 (d, 2H, J = 8.2 Hz), 7.08 (s, 1H), 7.00 (d, 2H, J = 8.2 Hz), 5.66 (s, 2H), 5.11 (s, 2H), 4.06 (q, 2H, J = 7.2 Hz), 3.58 (t, 2H, J = 8.2 Hz), 0.99-0.91 (m, 5H), 0.00 (s, 9H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 160.8, 158.2, 137.1, 132.9, 131.6, 128.6, 128.0, 127.6, 126.7, 119.8, 113.9, 99.4, 77.8, 70.0, 66.5, 60.2, 17.9, 13.8, -1.3; HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for  $C_{26}H_{32}^{81}$ BrNO<sub>4</sub>SiNa 554.1161; Found 554.1165.

# Ethyl 4-(4-(benzyloxy)-3-(trimethylsilyl)phenyl)-3-(4-(benzyloxy)phenyl)-1-((2-

# (trimethylsilyl)ethoxy)methyl)-1*H*-pyrrole-2-carboxylate (21).

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 12 (288.1 mg, 0.543 mmol, 1.0 equiv), 20 (621.5 mg, 1.63 mmol, 3.0 equiv), Ba(OH)2: 8H2O (1.026 g, 3.25 mmol, 6.0 equiv), Pd(PPh3)4 (62.2 mg, 53.8 µmol, 10 mol%), water (1.1 mL), and 1,4dioxane (4.3 mL). The flask was placed in a preheated oil bath and heated at 100  $^{\circ}\,$  C for 15 min, at which time the reaction was quenched with saturated aqueous ammonium chloride (10 mL). The resulting mixture was extracted twice with ethyl acetate (10 mL). The combined organic extracts were washed with brine (20 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/diethyl ether = 10:1) followed by preparative SEC-HPLC to provide the title compound 21 as a colorless oil (167.8 mg, 0.238 mmol, 44%).  $R_f =$ 0.43 (hexane/diethyl ether = 3:1); IR (ATR, cm<sup>-1</sup>): 1693, 1236, 1175, 1104, 1080, 836, 736, 696; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.46-7.31 (m, 10H), 7.14-7.05 (m, 5H), 6.91 (d, 2H, J = 8.0 Hz), 6.73 (d, 1H, J = 8.0 Hz), 5.72 (s, 2H), 5.07 (s, 2H), 5.01 (s, 2H), 4.06 (q, 2H, J = 7.2 Hz), 3.62 (t, 2H, J = 8.2 Hz), 0.99-0.93 (m, 5H), 0.09 (s, 9H), 0.00 (s, 9H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 161.8, 161.7, 157.7, 137.3, 137.2, 135.2, 131.8, 131.7, 130.1, 128.5, 128.4, 127.9, 127.7, 127.4, 126.6, 125.2, 124.8, 120.2, 114.0, 110.2, 77.6, 70.0, 69.8, 66.2, 59.8, 17.9, 13.8, -0.9, -1.3; HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for  $C_{42}H_{51}NO_5Si_2Na$ 728.3203; Found 728.3184.

50

51

52

53

54

55

56

3

4

5

7

9

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

4-(4-(benzyloxy)-3-iodophenyl)-3-(4-Ethvl (benzyloxy)phenyl)-1-((2-2 (trimethylsilyl)ethoxy)methyl)-1H-pyrrole-2-carboxylate (S5). A 20-mL test tube equipped with a Teflon-coated magnetic stirring bar was charged with 21 (46.7 mg, 66.1 µmol, 1.0 equiv) and CH<sub>2</sub>Cl<sub>2</sub> (1.7 mL). After the solution was cooled to 6 -78 ° C, ICl (0.1 M in CH<sub>2</sub>Cl<sub>2</sub>, 730 μL, 73 μmol, 1.1 equiv) was added dropwise to the test tube. After stirring at  $-78^{\circ}$  C for 8

20 min, the mixture was allowed to warm to  $-40^{\circ}$  C for 20 min, at which time the mixture was treated with saturated 10 aqueous sodium thiosulfate (3 mL). After partitioned, the 11 aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 mL) three time. 12 The combined organic extracts were washed with brine (6 13 mL), dried over sodium sulfate, and filtered. The filtrate was 14 concentrated under reduced pressure to provide the title 15 compound S5 as a brown oil (48.4 mg, 63.7  $\mu$ mol, 96%). Rf = 0.47 (hexane/diethyl ether = 2:1); M.p.  $187-188^{\circ}$  C; IR (ATR, 16 17 cm<sup>-1</sup>): 1695, 1281, 1240, 1176, 1104, 1082, 1043, 1020, 858, 834, 805, 735, 697; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.62 (d, 1H, J = 18 2.2 Hz), 7.48-7.31 (m, 10H), 7.12-7.09 (m, 3H), 6.92 (d, 2H, J = 19 8.7 Hz, 6.87 (dd, 1H, J = 8.6, 2.2 Hz), 6.63 (d, 1H, J = 8.6 Hz), 20 5.71 (s, 2H), 5.09 (s, 2H), 5.08 (s, 2H), 4.06 (q, 2H, I = 7.2 Hz), 21 3.61 (t, 2H, J = 8.2 Hz), 0.99-0.94 (m, 5H), 0.01 (s, 9H);  ${}^{13}C{}^{1}H$ 22 NMR (100 MHz, CDCl<sub>3</sub>): δ 161.7, 157.8, 155.6, 139.0, 137.2, 136.7, 23 131.8, 131.7, 129.4, 129.3, 128.7, 128.6, 128.0, 127.9, 127.6, 127.1, 24 125.4, 123.2, 120.4, 114.2, 112.3, 86.5, 77.8, 70.9, 70.1, 66.4, 60.0, 25 17.9, 13.9, -1.3; HRMS (DART/TOF) m/z: [M + H]<sup>+</sup> Calcd for 26 C<sub>39</sub>H<sub>43</sub>INO<sub>5</sub>Si 760.1955; Found 760.1927. 27

> Ethvl 4-(4-(benzyloxy)-3-iodophenyl)-3-(4-(benzyloxy)phenyl)-1H-pyrrole-2-carboxylate (S6).

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with S5 (220 mg, 0.290 mmol, 1.0 equiv) and CH<sub>2</sub>Cl<sub>2</sub> (18 mL). After the solution was cooled to 0  $^{\circ}$  C, TFA (222 µL, 2.90 mmol, 10 equiv) was added dropwise to the flask. After stirring at 0 ° C for 4 h, to the flask was added TFA (111 µL, 1.45 mmol, 5.0 equiv). The reaction mixture was stirred at o  $^{\circ}$  C for 4 h, at which time the reaction was guenched with water. After partitioned, the aqueous layer was extracted twice with CH<sub>2</sub>Cl<sub>2</sub> (6 mL). The combined organic extracts were washed with saturated aqueous sodium hydrogen carbonate (20 mL) and brine (20 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude hemiaminal (186 mg) as a yellow amorphous, which was used for the next step without further purification.

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with the crude hemiaminal, Na<sub>2</sub>CO<sub>3</sub> (246 mg, 2.32 mmol, 8.0 equiv), NH<sub>2</sub>OH · HCl (20.4 mg, 0.316 mmol, 1.1 equiv), THF (1.9 mL), and water (0.44 mL). After stirring at room temperature for 15 h, the reaction was quenched with water (5 mL) and 1 M aqueous hydrochloric acid (2 mL). The resulting mixture was extracted with ethyl acetate (5 mL) three times. The combined organic extracts were washed with brine (20 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/ $CH_2Cl_2$  = 1:2 to hexane/methyl acetate = 3:1, gradient). The obtained product was washed with hexane to provide the title compound S6 as a yellow amorphous (123 mg, 0.196

mmol, 68%).  $R_f = 0.43$  (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 1:2); IR (ATR, cm<sup>-1</sup>): 1669, 1530, 1454, 1378, 1282, 1240, 1175, 1040, 1024, 834, 736, 697; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.20 (br s, 1H), 7.66 (d, 1H, J = 2.3 Hz), 7.49-7.31 (m, 10H), 7.18 (d, 2H, J = 8.7 Hz), 7.03 (d, 1H, J = 3.2 Hz), 6.94-6.89 (m, 3H), 6.65 (d, 1H, J = 8.7 Hz), 5.09 (s, 2H×2), 4.20 (q, 2H, *J* = 7.2 Hz), 1.17 (t, 3H, *J* = 7.2 Hz); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 161.4, 157.9, 155.6, 138.9, 137.1, 136.6, 132.0, 129.6, 129.4, 128.8, 128.6, 128.5, 127.93, 127.86, 127.6, 127.0, 126.7, 124.7, 120.3, 119.9, 114.1, 112.3, 86.5, 70.8, 70.0, 60.3, 14.2; HRMS (DART/TOF) m/z:  $[M + H]^+$  Calcd for C<sub>33</sub>H<sub>29</sub>INO<sub>4</sub> 630.1141; Found 630.1136.

#### 4-(4-(benzyloxy)-3-iodophenyl)-3-(4-Ethvl (benzyloxy)phenyl)-1-(2-(4-(benzyloxy)phenyl)-2oxoethyl)-1H-pyrrole-2-carboxylate (S7).

A 10-mL screw-top test tube equipped with a Teflon-coated magnetic stirring bar was charged with S6 (116 mg, 0.185 mmol, 1.0 equiv), 17 (84.5 mg, 0.277 mmol, 1.5 equiv), K<sub>2</sub>CO<sub>3</sub> (50.9 mg, 0.368 mmol, 2.0 equiv), and DMF (620 µL). After stirring at room temperature for 4 h, the reaction was quenched with water (4 mL). The resulting mixture was extracted twice with diethyl ether (4 mL). The combined organic extracts were washed with water (5 mL) and brine (5 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography  $(hexane/CH_2Cl_2 = 1:1 to hexane/methyl acetate = 3:1, gradi$ ent) to provide the title compound S7 as a yellow amorphous (141 mg, 0.165 mmol, 89%). Rf = 0.43 (hexane/diethyl ether = 2:1); IR (ATR, cm<sup>-1</sup>): 1688, 1681, 1600, 1280, 1233, 1171, 1100, 833, 736, 697, 632; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  8.02 (d, 2H, J = 8.8 Hz), 7.66 (d, 1H, J = 2.3 Hz), 7.48-7.31 (m, 15H), 7.16 (d, 2H, J = 8.8 Hz), 7.07 (d, 2H, J = 8.8 Hz), 6.93-6.89 (m, 4H), 6.63 (d, 1H, J = 8.8 Hz), 5.74 (s, 2H), 5.16 (s, 2H), 5.10 (s, 2H), 5.07 (s, 2H), 3.94 (q, 2H, J = 7.2 Hz), 0.85 (t, 3H, J = 7.2 Hz); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 191.7, 163.1, 161.7, 157.6, 155.4, 138.9, 137.2, 136.6, 136.0, 131.9, 130.9, 130.3, 130.0, 129.5, 129.2, 128.7, 128.52, 128.50, 128.3, 128.2, 128.0, 127.84, 127.80, 127.5, 127.0, 122.8, 120.3, 114.9, 114.0, 112.2, 86.4, 70.7, 70.1, 69.9, 59.7, 55.5, 13.7; HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for C<sub>48</sub>H<sub>40</sub>INO<sub>6</sub>Na 876.1798; Found 876.1786.

## 4-(4-(Benzyloxy)-3-iodophenyl)-3-(4-(benzyloxy)phenyl)-1-(2-(4-(benzyloxy)phenyl)-2oxoethyl)-1H-pyrrole-2-carboxylic acid (S8).

A 10-mL screw-top test tube equipped with a Teflon-coated magnetic stirring bar was charged with S7 (38.3 mg, 44.9 µmol, 1.0 equiv), potassium trimethylsilanolate (38.7 mg, 0.271 mmol, 6.0 equiv), and 1,2-dichloroethane (0.45 mL). The test tube was placed in a preheated aluminum block and heated at 80  $^{\circ}$  C for 15 min, at which time the reaction mixture was treated with 1 M aqueous hydrochloric acid (3 mL). After partitioned, the aqueous layer was extracted with  $CHCl_3$  (3 mL) three times. The combined organic extracts were washed with brine (3 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was washed with the mixture of diethyl ether (2 mL) and hexane (1 mL) to provide the title compound S8 as a colorless solid (23.8 mg, 28.8  $\mu$ mol, 64%). R<sub>f</sub> = 0.18 (hexane/methyl acetate = 2:1); M.p. 187-188° C; IR (ATR, cm<sup>-1</sup>): 1686, 1650, 1600, 1451, 1287, 1239, 1172, 1011, 736, 696; <sup>1</sup>H NMR (400 MHz, THF- $d_8$ ):  $\delta$  8.02 (d, 2H, J =

55

56

57 58 59

60

8.8 Hz), 7.64 (d, 1H, *J* = 2.0 Hz), 7.48-7.24 (m, 15H), 7.14 (d, 2H, *J* = 8.8 Hz), 7.11-7.09 (m, 3H), 6.91-6.87 (m, 3H), 6.74 (d, 1H, *J* = 8.8 Hz), 5.84 (s, 2H), 5.18 (s, 2H), 5.064 (s, 2H), 5.059 (s, 2H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, THF-*d*<sub>8</sub>):  $\delta$  191.6, 163.6, 162.9, 158.6, 156.2, 139.4, 138.6, 138.0, 137.7, 132.7, 130.94, 130.87, 130.7, 129.8, 129.5, 129.3, 129.0, 128.9, 128.5, 128.1, 128.0, 127.9, 127.6, 123.1, 121.2, 115.2, 114.2, 112.6, 86.4, 71.1, 70.6, 70.2, 56.0; HRMS (ESI/TOF) *m*/*z*: [M - H]<sup>-</sup> Calcd C<sub>46</sub>H<sub>35</sub>INO<sub>6</sub> 824.1509; Found 824.1533.

## Lukianol B.

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with S8 (44.4 mg, 53.8 µmol, 1.0 equiv), potassium acetate (95.7 mg, 0.975 mmol, 18 equiv), and acetic anhydride (5.4 mL). The flask was placed in a preheated oil bath and heated at 100  $^{\circ}$  C for 30 min. After cooling to room temperature, the acetic anhydride was removed azeotropically with toluene (50 mL) under reduced pressure. The residue was dissolved in ethyl acetate (6 mL) and the resulting mixture was washed with saturated aqueous ammonium chloride (6 mL). After partitioned, the aqueous layer was extracted twice with ethyl acetate (6 mL). The combined organic extracts were washed with brine (10 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/methyl acetate = 3:1) to provide the corresponding product 22 (27.3 mg) as a yellow amorphous, which was used for the next reaction without further purification.  $R_f = 0.29$  (hexane/methyl acetate = 3:1).

A flame-dried 20-mL Schlenk tube equipped with a Tefloncoated magnetic stirring bar and a rubber septum was charged with 22 (27.3 mg, 33.8 µmol, 1.0 equiv), pentamethylbenzene (29.3 mg, 198 µmol, 6.0 equiv), and CH<sub>2</sub>Cl<sub>2</sub> (3.4 mL). The solution was cooled to -78 ° C. BCl<sub>3</sub> (1 M in CH<sub>2</sub>Cl<sub>2</sub>, 200 µL, 0.20 mmol, 6.0 equiv) was added dropwise to the Schlenk tube. After stirring at  $-78^{\circ}$  C for 30 min, the reaction mixture was allowed to warm to 0  $^{\circ}$  C and stirred for 20 min, at which time the mixture was treated with methanol (4 mL) and concentrated under reduced pressure. The residue was dissolved in methanol (3 mL) and the resulting mixture was washed with hexane (3 mL) three times. The methanol layer was concentrated under reduced pressure to give a crude product, which was washed with acetone (2 mL) to provide lukianol B as a colorless solid (13.1 mg, 24.4 µmol, 45% over 2 steps), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra (acetone $d_6$ ) were identical to those reported in the literature.<sup>21a</sup> R<sub>f</sub> =  $0.26 (CH_2Cl_2/methanol = 20:1); M.p. > 275^{\circ} C; IR (ATR, cm^{-1})$ <sup>1</sup>): 2924, 1697, 1680, 1658, 1612, 1536, 1518, 1503, 1433, 1422, 1415, 1408, 1274, 1239, 1208, 1181, 836; <sup>1</sup>H NMR (400 MHz, acetone*d*<sub>6</sub>): δ 9.15 (s, 1H), 8.81 (s, 1H), 8.44 (s, 1H), 7.93 (s, 1H), 7.65 (d, 2H, J = 8.8 Hz), 7.63 (d, 1H, J = 2.0 Hz), 7.61 (s, 1H), 7.18 (d, 2H, *J* = 8.8 Hz), 7.00 (dd, 1H, *J* = 8.8, 2.0 Hz), 6.96 (d, 2H, *J* = 8.8 Hz), 6.84-6.81 (m, 3H);  ${}^{13}C{}^{1}H$  NMR (100 MHz, acetone- $d_6$ ):  $\delta$ 159.5, 157.9, 156.5, 154.4, 142.9, 140.1, 133.0, 130.9, 130.5, 128.5, 51 127.3, 126.8, 124.9, 123.4, 120.8, 116.8, 115.7, 115.6, 113.8, 103.8, 52 84.4; HRMS (DART/TOF) m/z:  $[M + H]^+$  Calcd for 53 C<sub>25</sub>H<sub>17</sub>INO<sub>5</sub> 538.0151; Found 538.0177.

#### 2,4-Diiodophenol (S9).

A 500-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 2-iodo-phenol

(3.820 g, 17.4 mmol, 1.0 equiv), NaOH (1.114 g, 28.9 mmol 1.7 equiv), and methanol (43 mL). After the solution was cooled to o  $^{\circ}$  C, I<sub>2</sub> (6.166 g, 24.3 mmol, 1.4 equiv) in methanol (43 mL) was added dropwise to the flask at 0  $^{\circ}$  C. After stirring for 85 min at 0  $^{\circ}\,$  C, I<sub>2</sub> (668.0 mg, 2.63 mmol, 0.15 equiv) in methanol (5 mL) was added dropwise to the flask at o  $^{\circ}$  C and the resulting mixture was stirred for 40 min, at which time the reaction mixture was treated with saturated aqueous sodium thiosulfate (100 mL) and concentrated under reduced pressure. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) and the resulting mixture was washed with saturated aqueous ammonium chloride (50 mL). After partitioned, the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (50 mL). The combined organic extracts were washed with brine (100 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/ $CH_2Cl_2 = 2:1$ ) to provide the corresponding product So as a colorless solid (3.966 g, 11.5 mmol, 66%), whose  ${}^{1}$ H and <sup>13</sup>C NMR spectra were identical to those reported in the literature.33

#### 2,4-Diiodo-1-(phenylmethoxy)-benzene (S10).

A 200-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with **S9** (3.732 g, 10.8 mmol, 1.0 equiv), potassium carbonate (2.986 g, 21.6 mmol, 2.0 equiv), benzyl bromide (1.849 g, 10.8 mmol, 1.0 equiv), and acetonitrile (22 mL). The flask was placed in a preheated oil bath and heated at 80  $^{\circ}$  C for 30 min. After cooling to room temperature, the resulting mixture was filtered through a pad of Celite. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane) to provide the corresponding product **S10** as a colorless solid (4.389 g, 10.1 mmol, 93%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra were identical to those reported in the literature.<sup>34</sup>

# (2-(Benzyloxy)-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)trimethylsilane (20).

A flame-dried 100-mL Schlenk tube equipped with a Tefloncoated magnetic stirring bar and a rubber septum was charged with S10 (1.005 g, 2.30 mmol, 1.0 equiv), TMSCl (1.5 mL, 12 mmol, 5 equiv), and anhydrous THF (11.5 mL). After the solution was cooled to -78  $^{\circ}$  C, *n*-BuLi (1.55 M in *n*hexane, 1.48 mL, 2.3 mmol, 1.0 equiv) was added to the Schlenk tube. The reaction mixture was stirred at  $-78^{\circ}$  C for 30 min, the resulting mixture was allowed to warm to room temperature with stirring over 1 h, at which time the reaction mixture was treated with water (20 mL) and saturated aqueous ammonium chloride (5 mL). The resulting mixture was extracted twice with diethyl ether (15 mL). The combined organic extracts were washed with brine (30 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane) to provide a mixture of the corresponding product S11 and the disilylated byproduct (673.3 mg).

A flame-dried 50-mL Schlenk tube equipped with a Tefloncoated magnetic stirring bar and a rubber septum was charged with **S11** (673.3 mg, 1.51 mmol, 1.0 equiv) and anhydrous THF (7.6 mL). After the solution was cooled to  $-78^{\circ}$  C, *n*-BuLi (1.55 M in *n*-hexane, 1.1 mL, 1.7 mmol, 1.1 equiv) was

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

54

55

56

57 58 59

60

added to the Schlenk tube. The reaction mixture was stirred at -78 ° C for 5 min, to the solution was added *i*-PrOBpin (337.4 mg, 1.81 mmol, 1.2 equiv) at  $-78^{\circ}$  C. After stirring at - $78^{\circ}$  C for 10 min, the resulting mixture was allowed to warm to room temperature with stirring over 1 h, at which time the reaction mixture was treated with water (10 mL). The resulting mixture was extracted twice with diethyl ether (10 mL). The combined organic extracts were washed with brine (20 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane to hexane/ $CH_2Cl_2 = 2:1$ , gradient) to provide the corresponding product 20 as a colorless solid (388.7 mg, 1.02 mmol, 44% over 2 steps).  $R_f = 0.26$  (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 2:1); M.p. 99-100 ° C; IR (ATR, cm<sup>-1</sup>): 1588, 1371, 1352, 1279, 1231, 1145, 1102, 854, 838, 695, 669; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.83 (s, 1H), 7.78 (d, 1H, J = 8.2 Hz), 7.41-7.30 (m, 5H), 6.85 (d, 1H, J = 8.2 Hz, 5.09 (s, 2H), 1.31 (s, 12H), 0.24 (s, 9H);  $^{13}C{^{1}H}$ NMR (100 MHz, CDCl<sub>3</sub>): δ 166.0, 142.1, 138.3, 137.0, 128.6, 127.9, 127.6, 127.4, 120.7, 109.9, 83.6, 69.8, 25.0, -0.7; HRMS  $(ESI/TOF) m/z: [M + Na]^+$  Calcd for C<sub>22</sub>H<sub>32</sub>BO<sub>3</sub>SiNa 405.2148; Found 405.2145.

## TOTAL SYNTHESIS OF NINGALIN B

Ethyl 3,4-bis[3,4-bis(benzyloxy)phenyl]-1-[(2-(trimethylsilyl)ethoxy)methyl]-1H-pyrrole-2-carboxylate (24).

25 A 50-mL round-bottomed flask equipped with a Teflon-26 coated magnetic stirring bar was charged with 8 (424.6 mg, 27 0.994 mmol, 1.0 equiv), Ba(OH)2.8H2O (1.88 g, 5.97 mmol, 28 6.0 equiv), 23 (1.66 g, 3.99 mmol, 4.0 equiv), Pd(PPh<sub>3</sub>)<sub>4</sub> (113.9 29 mg, 98.6 µmol, 10 mol%), water (2 mL), and 1,4-dioxane (8 30 mL). The flask was placed in a preheated oil bath and heated 31 at 100  $^{\circ}$  C for 15 min, at which time the solution was treated 32 with water (5 mL). The reaction mixture was extracted twice with diethyl ether (10 mL). The combined organic extracts 33 were washed with water and brine, dried over sodium sulfate, 34 and filtered. The filtrate was concentrated under reduced 35 pressure to give a crude product, which was purified by silica 36 gel column chromatography (hexane/diethyl ether = 5:1 to 37 hexane/diethyl ether = 3:1, gradient) and followed by prepar-38 ative SEC-HPLC to provide the title product 24 as a pale yel-39 low oil (499 mg, 0.590 mmol, 59%). Rf = 0.21 (hexane/diethyl 40 ether = 3:1); IR (ATR, cm<sup>-1</sup>): 2949, 1695, 1530, 1498, 1455, 1426, 41 1373, 1355, 1247, 1212, 1136, 1081, 1023, 836; <sup>1</sup>H NMR (400 MHz, 42 CDCl<sub>3</sub>): δ 7.44-7.21 (m, 20H), 7.07 (s, 1H), 6.89 (d, 1H, J = 8.2 43 Hz), 6.84 (d, 1H, J = 1.6 Hz), 6.76 (d, 1H, J = 8.2 Hz), 6.73 (dd, 44 1H, J = 8.0 Hz, 2.3 Hz), 6.65 (dd, 1H, J = 8.2, 2.3 Hz), 6.61 (d, 45 1H, J = 2.3 Hz, 5.69 (s, 2H), 5.15 (s, 2H), 5.10 (s, 2H), 5.02 (s, 2H), 4.75 (s, 2H), 3.98 (q, 2H, J = 7.2 Hz), 3.60 (t, 2H, J = 8.1 46 Hz), 0.95 (t, 2H, I = 8.1 Hz), 0.89 (t, 3H, I = 7.2 Hz), -0.01 (s, 47 9H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 161.7, 148.6, 148.3, 48 147.8, 147.2, 137.6, 137.5, 137.4, 131.5, 129.5, 128.6, 128.5, 127.9, 49 127.8, 127.7, 127.4, 127.3, 127.2, 125.2, 124.5, 123.9, 120.6, 120.3, 50 117.7, 115.0, 114.8, 114.4, 77.7, 71.4, 71.3, 70.9, 66.4, 59.9, 17.9, 51 13.9, -1.3; HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for 52 C<sub>53</sub>H<sub>55</sub>NO<sub>7</sub>SiNa 868.3646; Found 868.3667. 53

## 7,8-Bis(benzyloxy)-1-[3,4-

bis(benzyloxy)phenyl]chromeno[3,4-b]pyrrol-4(3H)-one (25).

A 10-mL screw-top test tube equipped with a Teflon-coated magnetic stirring bar was charged with compound 24 (245.1 mg, 0.290 mmol, 1.0 equiv), 18-crown-6 (153.3 mg, 0.580 mmol, 2.0 equiv), KOH (162.7 mg, 10.0 mmol, 10.0 equiv), and EtOH (1.5 mL). The test tube was placed in a preheated aluminum block and heated at reflux for 14.5 h, at which time the resulting mixture was added 18-crown-6 (76.2 mg, 0.288 mmol, 1.0 equiv) and KOH (80.3 mg, 1.43 mmol, 4.9 equiv). After stirring at reflux for 20.5 h, the reaction mixture was treated with 1 M aqueous hydrochloric acid, at which time yellow solid was precipitated and filtered to give a crude **S12** as a yellow solid (281.0 mg), which was used for the next step without further purification.

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with the crude S12 (281.0 mg) and anhydrous ethyl acetate (29 mL). To the solution was added Pb(OAc)<sub>4</sub> (258.4 mg, 0.58 mmol), and the flask was placed in a preheated oil bath and heated at reflux for 30 min.  $Pb(OAc)_4$  (130.2 mg, 0.29 mmol) was added to the solution, and the reaction mixture was heated at reflux for 20 min. Pinacol (113.2 mg, 0.956 mmol) was added to the flask, and the reaction mixture was treated with water (30 mL). The resulting mixture was extracted with ethyl acetate (30 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/ $CH_2Cl_2$  = 1:1 to  $CH_2Cl_2$ , gradient) to provide the corresponding product S13 as a yellow solid (89.6 mg, 0.110 mmol).

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with compound S13 (89.6 mg, 0.110 mmol, 1.0 equiv) and CH<sub>2</sub>Cl<sub>2</sub> (2.8 mL). TFA (0.60 mL, 18 mmol, xs) was added to the flask. The resulting mixture was stirred at room temperature for 10 min. After the reaction mixture was concentrated under reduced pressure, the flask was charged with anhydrous THF (0.8 mL), water (0.2 mL), NH<sub>2</sub>OH·HCl (6.9 mg, 99 µmol, 0.90 equiv), and Na<sub>2</sub>CO<sub>3</sub> (35.1 mg, 0.331 mmol, 3.0 equiv). After stirring at room temperature for 25 min, the reaction mixture was treated with water (1 mL). The resulting mixture was extracted with ethyl acetate (2 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was washed with diethyl ether (10 mL) to provide the title compound **25** as a pale yellow solid (48.3 mg, 70.4 µmol, 24% over 3 steps).  $R_f = 0.30$  (CH<sub>2</sub>Cl<sub>2</sub>/methanol = 100:1); M.p. 178-179

C; IR (ATR, cm<sup>-1</sup>): 3242, 3063, 3033, 1696, 1549, 1516, 1499, 1272, 1207, 1023, 735, 695; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  9.60 (br s, 1H), 7.51<sup>-7.29</sup> (m, 15H), 7.22<sup>-7.19</sup> (m, 3H), 7.17 (s, 1H), 7.14 (d, 1H, *J* = 3.2 Hz), 7.12<sup>-7.08</sup> (m, 2H), 7.07 (d, 1H, *J* = 1.8 Hz), 7.03 (d, 1H, *J* = 8.2 Hz), 6.965 (dd, 1H, *J* = 7.8, 1.8 Hz), 6.959 (s, 1H), 5.26 (s, 2H), 5.21 (s, 2H), 5.18 (s, 2H), 4.84 (s, 2H); <sup>13</sup>Cl<sup>4</sup>H] NMR (100 MHz, CDCl<sub>3</sub>):  $\delta$  156.7, 149.1, 148.8, 148.7, 146.1, 145.2, 137.3, 137.1, 136.6, 136.5, 128.73, 128.70, 128.66, 128.5, 128.3, 128.2, 128.1, 128.0, 127.8, 127.6, 127.5, 127.4, 126.6, 123.0, 121.3, 116.7, 116.4, 115.0, 111.1, 108.1, 103.6, 71.5, 71.1, 70.9; HRMS (ESI/TOF) *m/z*: [M + Na]<sup>+</sup> Calcd for C<sub>45</sub>H<sub>35</sub>NO<sub>6</sub>Na 708.2362; Found 708.2367.

## 7,8-Bis(benzyloxy)-3-[3,4-bis(benzyloxy)phenethyl]-1-[3,4-bis(benzyloxy)phenyl]chromeno[3,4-b]pyrrol-4(3*H*)one (S14).

A 10-mL screw-top test tube equipped with a Teflon-coated magnetic stirring bar was charged with compound 25 (42.9 62.4 μmol, 1.0 equiv), 2-[3,4-bis(benzyloxy)mg. phenyl]ethanol (26) (31.3 mg, 93.6 µmol, 1.5 equiv), and PPh<sub>3</sub> (48.9 mg, 0.187 mmol, 3.0 equiv). After the flask was evacuated and backfilled with N2, anhydrous THF (0.3 mL) and DIAD (1.9 M, 99 µL, 0.19 mmol, 3.0 equiv) were added to the flask. The solution was stirred at room temperature for 15 min, at which time the reaction mixture was treated with water (1 mL). The resulting mixture was extracted with diethyl ether (1 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/ $CH_2Cl_2 = 4:1$  to CH<sub>2</sub>Cl<sub>2</sub>, gradient). The amorphous was washed with hexane (2 mL) to provide the titled product S14 as a white amorphous (43.9 mg, 43.7  $\mu$ mol, 70%). Rf = 0.21 (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 4:1); IR (ATR, cm<sup>-1</sup>): 1714, 1510, 1499, 1454, 1376, 1262, 1229, 1159, 1024, 745, 695; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.48-7.28 (m, 21H), 7.24-7.17 (m, 6H), 7.11 (s, 1H), 7.10-7.08 (m, 2H), 6.96 (d,  $_{2H, J} = 7.8 \text{ Hz}$ , 6.93 (d, 1H, J = 1.8 Hz), 6.83-6.80 (m, 2H), 6.61 (dd, 1H, I = 8.2, 2.3 Hz), 6.56 (d, 1H, I = 1.8 Hz), 6.51 (s, 1H), 5.21 (s, 2H×2), 5.10 (s, 2H), 5.08 (s, 2H), 4.97 (s, 2H), 4.79 (s, 2H), 4.53 (t, 2H, J = 6.6 Hz), 3.00 (t, 2H, J = 6.6 Hz); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 155.5, 149.1, 149.0, 148.7, 148.6, 147.8, 146.3, 144.9, 137.4, 137.3, 137.2, 137.0, 136.7, 136.6, 132.2, 131.5, 128.8, 128.7, 128.58, 128.55, 128.51, 128.4, 128.1, 128.0, 127.9, 127.8, 127.59, 127.56, 127.4, 127.35, 127.32, 127.1, 122.9, 121.8, 119.0, 116.5, 115.7, 115.5, 115.0, 114.9, 111.0, 108.1, 103.5, 71.6, 71.5, 71.3, 71.2, 71.1, 70.9, 51.0, 37.6; HRMS (ESI/TOF) m/z: [M  $+ Na^{+}$  Calcd for C<sub>67</sub>H<sub>55</sub>NO<sub>8</sub>Na 1024.3825; Found 1024.3788.

#### Ningalin B.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

A 25-mL pear-shaped flask equipped with a Teflon-coated magnetic stirring bar was charged with compound S14 (31.9 mg, 31.8 µmol, 1.0 equiv), Pd/C (31.9 mg, 100 wt%), EtOH (0.5 mL), and ethyl acetate (0.5 mL). After the flask was evacuated and backfilled with H<sub>2</sub> three times, the resulting mixture was stirred at room temperature for 30 min. The resulting mixture was filtered through a pad of Celite, and the filter cake was washed with hot methanol (10 mL). The filtrate was concentrated under reduced pressure to provide ningalin B as a pale gray solid (14.3 mg, 30.8  $\mu$ mol, 97%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra (DMSO-*d*<sub>6</sub>/CD<sub>3</sub>OD (4:1)) were not in good agreement with those reported in the literature.<sup>25a</sup> The <sup>1</sup>H NMR spectrum (CD<sub>3</sub>OD) was identical to that reported in Fenical' s supporting information (page 10). <sup>1</sup>H and <sup>13</sup>C NMR spectra (DMSO- $d_6$ /CD<sub>3</sub>OD (4:1)) were in good agreement with those reported.<sup>6h,25b</sup>  $R_f = 0.52$  (CH<sub>2</sub>Cl<sub>2</sub>/methanol = 3:1); M.p. >250 ° C; IR (ATR, cm<sup>-1</sup>): 3269, 2927, 1682, 1416, 1366, 1255, 1230, 1168, 816; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 7.14 (s, 1H), 6.89 (s, 1H), 6.84 (d, 1H, J = 7.8 Hz), 6.82 (d, 1H, J = 1.8 Hz), 6.79 (s, 1H), 6.71-6.65 (m, 2H), 6.54 (d, 1H, I = 1.8 Hz), 6.46(dd, 1H, J =8.2, 1.8 Hz), 4.56 (t, 2H, J = 7.0 Hz), 2.96 (t, 2H, J = 7.0 Hz); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>/CD<sub>3</sub>OD (4:1)) δ 7.19 (s, 1H), 7.06 (s, 1H), 6.81 (d, 1H, J = 8.4 Hz), 6.77 (d, 1H, J = 2.0Hz), 6.75 (s, 1H), 6.63 (dd, 1H, J = 7.6, 2.4 Hz), 6.62 (d, 1H, J = 7.8 Hz), 6.59 (d, 1H, J = 2.0 Hz), 6.43 (dd, 1H, J = 8.0, 2.0 Hz),

4.49 (t, 2H, *J* = 7.4 Hz), 2.87 (t, 2H, *J* = 7.4 Hz); <sup>13</sup>C[<sup>1</sup>H] NMR (100 MHz, DMSO-*d*<sub>6</sub>/CD<sub>3</sub>OD (4:1)):  $\delta$  155.2, 146.3, 145.5, 145.4, 145.3, 145.0, 144.1, 142.4, 133.0, 129.5, 126.8, 125.7, 121.1, 120.0, 119.5, 117.3, 116.5, 116.1, 115.8, 114.3, 110.0, 108.9, 103.7, 50.3, 37.5; HRMS (ESI/TOF) *m*/*z*: [M + Na]<sup>+</sup> Calcd for C<sub>25</sub>H<sub>19</sub>NO<sub>8</sub>Na 484.1008; Found 484.1031.

## 2-[3,4-Di(benzyloxy)phenyl]-4,4,5,5-tetramethyl-1,3,2dioxaborolane (23).

A flame-dried 100-mL two-necked test tube equipped with a Teflon-coated magnetic stirring bar and a rubber septum was charged with 3,4-dibenzyloxy-1-iodobenzene35 (1.04 g, 2.53 mmol, 1.0 equiv) and anhydrous THF (13 mL). After the solution was cooled to  $-78^{\circ}$  C, the resulting mixture was treated with n-BuLi (1.59 M in n-hexane, 1.65 mL, 2.63 mmol, 1.0 equiv) and stirred at -78 ° C for 5 min. To the solution was added *i*-PrOBpin (522.2 mg, 2.81 mmol, 1.1 equiv) at -78 ° C. After stirring at  $-78^{\circ}$  C for 10 min, the resulting mixture was allowed to warm to room temperature with stirring over 40 min, at which time the reaction mixture was treated with saturated aqueous ammonium chloride (10 mL). After partitioned, the aqueous layer was extracted with diethyl ether (10 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/diethyl ether = 10:1) to provide the title product 23 as a colorless solid (795.6 mg, 1.91 mmol, 76%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra were identical to those reported in the literature.<sup>36</sup>

## 2-[3,4-Di(benzyloxy)phenyl]ethanol (26).

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 2-(3,4dihydroxyphenyl)ethyl alcohol (347.4 mg, 2.25 mmol, 1.0 equiv), K<sub>2</sub>CO<sub>3</sub> (937.3 mg, 6.75 mmol, 3.0 equiv), benzyl bromide (643  $\mu$ L, 5.41 mmol, 2.4 equiv), and acetonitrile (2.25 mL). The flask was placed in a preheated oil bath and heated at reflux for 15 h. After cooling to room temperature, the resulting mixture was filtered through a pad of Celite. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/diethyl ether = 3:1) to provide the title product **26** as a colorless solid (631 mg, 1.88 mmol, 84%), whose <sup>1</sup>H NMR spectrum was identical to that reported in the literature.<sup>37</sup>

## TOTAL SYNTHESIS OF LAMELLARIN S

7,8-Bis(benzyloxy)-3-[3-(benzyloxy)-4-

methoxyphenethyl]-1-[3,4-

# bis(benzyloxy)phenyl]chromeno[3,4-*b*]pyrrol-4(3*H*)-one (28).

A 10-mL screw-top test tube equipped with a Teflon-coated magnetic stirring bar was charged with compound **25** (55.0 mg, 80.0  $\mu$ mol, 1.0 equiv), 3-benzyloxy-4-methoxybenzeneethanol (**27**) (31.9 mg, 0.123 mmol, 1.5 equiv), PPh<sub>3</sub> (61.8 mg, 0.236 mmol, 3.0 equiv). After the flask was evacuated and backfilled with N<sub>2</sub>, anhydrous THF (0.4 mL), and DIAD (1.9 M, 126  $\mu$ L, 0.240 mmol, 3.0 equiv) were added to the flask. The solution was stirred at room temperature for 20 min, at which time the reaction mixture was treated with water (1 mL). The resulting mixture was extracted with di-

23

24

59

60

ethyl ether (1 mL) three times. The combined organic ex-1 tracts were washed with water and brine, dried over sodium 2 sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified 3 4 by silica gel column chromatography (hexane/ $CH_2Cl_2 = 1:3$  to hexane/ $CH_2Cl_2 = 1:4$ , gradient) to provide the title product 28 5 as a yellow amorphous (52.4 mg, 56.6  $\mu$ mol, 71%). Rf = 0.34 6 (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 1:4); IR (ATR, cm<sup>-1</sup>): 1709, 1588, 1513, 1419, 7 1260, 1137, 1025, 736, 696; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.49-8 7.28 (m, 15H), 7.25-7.16 (m, 7H), 7.13 (s, 1H), 7.11-7.07 (m, 2H), 9 6.98 (d, 1H, J = 8.2 Hz), 6.95 (s, 1H), 6.91 (d, 1H, J = 1.8 Hz), 10 6.83 (dd, 1H, J = 8.2, 2.1 Hz), 6.78 (d, 1H, J = 8.2 Hz), 6.65 (dd, 11 1H, J = 8.0, 2.1 Hz), 6.48 (d, 1H, J = 2.3 Hz), 6.47 (s, 1H), 5.222 12 (s, 1H), 5.219 (s, 2H), 5.11 (s, 2H), 4.94 (s, 2H), 4.80 (s, 2H), 13 4.52 (t, 2H, J = 6.9 Hz), 3.81 (s, 3H), 3.00 (t, 2H, J = 6.9 Hz); 14 <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 155.4, 149.1, 148.7, 148.5, 15 148.1, 146.3, 144.9, 137.3, 137.1, 137.0, 136.64, 136.58, 132.2, 130.5, 128.73, 128.66, 128.6, 128.5, 128.4, 128.1, 128.01, 127.99, 127.83, 16 17 127.76, 127.60, 127.56, 127.3, 127.1, 122.9, 121.6, 119.0, 116.5, 115.0, 114.8, 112.0, 111.0, 108.1, 103.5, 71.5, 71.3, 71.1, 70.87, 70.86, 56.1, 18 51.1, 37.5 (five aromatic carbon signals are missing due to 19 overlapping); HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for 20 C<sub>61</sub>H<sub>51</sub>NO<sub>8</sub>Na 948.3512; Found 948.3548. 21

## 2,3,11-Tris(benzyloxy)-14-(3,4-bis(benzyloxy)phenyl)-12methoxy-8,9-dihydro-6*H*-chromeno[4',3':4,5]pyrrolo[2,1*a*]isoquinolin-6-one (S15).

25 A 50-mL round-bottomed flask equipped with a Teflon-26 coated magnetic stirring bar was charged with compound 28 27 (33.1 mg, 35.7  $\mu$ mol, 1.0 equiv) and anhydrous CH<sub>2</sub>Cl<sub>2</sub> (1 mL). After the resulting solution was cooled to  $-40^{\circ}$  C, a solution 28 of PIFA (18.1 mg, 42.1 µmol, 1.2 equiv) and BF3 · OEt2 (11 µL, 87 29 µmol, 2.4 equiv) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (2.6 mL) was added 30 dropwise to the flask. The reaction mixture was stirred at -31 40  $^{\circ}$  C for 1 h, at which time the reaction was guenched with 32  $_{2}$  M aqueous NH<sub>3</sub> (3 mL). The reaction mixture was extracted 33 twice with CH<sub>2</sub>Cl<sub>2</sub> (3 mL). The combined organic extracts 34 were washed with water and brine, dried over sodium sulfate, 35 and filtered. The filtrate was concentrated under reduced 36 pressure to give a crude product, which was purified by silica 37 gel column chromatography (hexane/diethyl ether = 1:1 to 38 CH<sub>2</sub>Cl<sub>2</sub>, gradient). The solid was washed twice with diethyl 39 ether (2 mL) to provide the titled product S15 as a colorless 40 solid (20.1 mg, 21.8  $\mu$ mol, 61%). R<sub>f</sub> = 0.14 (hexane/diethyl ether = 1:1); M.p. 167-168 ° C; IR (ATR, cm<sup>-1</sup>): 1704, 1543, 1512, 41 1484, 1454, 1420, 1270, 1210, 1165, 1037, 1026, 735, 696; <sup>1</sup>H NMR 42 (400 MHz, CDCl<sub>3</sub>): δ 7.48-7.28 (m, 19H), 7.24-7.21 (m, 4H), 43 7.17-7.15 (m, 2H), 7.10 (d, 1H, J = 8.2 Hz), 7.05 (d, 1H, J = 1.844 Hz), 6.98 (dd, 1H, J = 8.2, 1.8 Hz), 6.92 (s, 1H), 6.73 (s, 1H), 45 6.72 (s, 1H), 6.53 (s, 1H), 5.29 (d, 1H, J = 12.4 Hz), 5.24 (d, 1H, J 46 = 12.4 Hz), 5.19 (s, 2H), 5.148 (s, 2H), 5.145 (d, 1H, J = 12.4 Hz), 47 5.10 (d, 1H, J = 12.4 Hz), 4.85-4.78 (m, 1H), 4.73 (d, 1H, J = 12.8 48 Hz), 4.68 (d, 1H, J = 12.4 Hz), 4.64-4.57 (m, 1H), 3.18 (s 3H), 49 3.06-2.92 (m, 2H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 155.6, 50 149.6, 148.7, 148.5, 148.2, 148.1, 146.2, 144.9, 137.2, 136.83, 136.77, 51 136.7, 136.0, 128.8, 128.7, 128.60, 128.56, 128.5, 128.14, 128.08, 52 128.0, 127.8, 127.4, 127.32, 127.27, 127.20, 127.18, 126.4, 124.0, 53 120.5, 117.3, 115.4, 114.9, 113.8, 113.3, 110.9, 109.1, 107.8, 103.3, 71.3, 54 71.02, 70.97, 70.5, 55.2, 42.5, 28.7 (four aromatic carbon sig-55 nals and a benzylic position carbon are missing due to overlapping); HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for 56 C61H49NO8Na 946.3356; Found 946.3365. 57 58

## Lamellarin S.

A 10 mL pear-shaped flask equipped with a Teflon-coated magnetic stirring bar was charged with compound S17 (17.1 mg, 18.5 µmol, 1.0 equiv), Pd/C (17.1 mg, 100 wt%), EtOH (0.3 mL), and EtOAc (0.3 mL). After the flask was evacuated and backfilled with H<sub>2</sub> three times, the resulting mixture was stirred at room temperature for 14 h. The resulting mixture was filtered through a pad of Celite, and the filter cake was washed with hot methanol (8 mL). The filtrate was concentrated under reduced pressure to provide lamellarin S as a pale gray solid (8.6 mg, 18 µmol, 98%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra (CD<sub>3</sub>OD) were identical to those reported in the literature.<sup>26a,26b</sup>  $R_f = 0.69$  (chloroform/ethanol = 3:1); M.p. >250 ° C; IR (ATR, cm<sup>-1</sup>): 3251, 2929, 2854, 1680, 1597, 1420, 1279, 1252, 1201, 1159, 1040; <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 6.99 (d, 1H, J = 8.0 Hz), 6.88 (d, 1H, J = 1.4 Hz), 6.83 (s, 1H), 6.77(dd, 1H, J = 8.0, 1.4 Hz), 6.76 (s, 1H), 6.71 (s, 1H), 6.62 (s, 1H), $4.73^{-}4.53$  (m, 2H), 3.38 (s, 3H), 3.01 (t, 2H, J = 6.2 Hz);  ${}^{13}C{}^{1}H$ NMR (100 MHz, CD<sub>3</sub>OD): δ 157.6, 148.1, 147.4, 147.3, 146.7, 146.6, 143.4, 138.6, 129.9, 128.7, 128.1, 123.7, 120.2, 119.1, 117.5, 116.4, 115.8, 113.8, 111.3, 110.4, 109.8, 104.2, 55.6, 43.5, 29.3; HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for  $C_{26}H_{19}NO_8Na$ 496.1008; Found 496.1028.

#### 4-Methoxy-3-(phenylmethoxy)-benzeneethanol (27).

A 500-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring was charged with (methoxymethyl)triphenylphosphonium chloride (14.18 g, 41.4 mmol, 1.5 equiv) and THF (140 mL). After the solution was cooled to o<sup>°</sup> C, potassium *tert*-butoxide (6.184 g, 55.1 mmol, 2.0 equiv) was added potionwise over 3 min to the flask and the resulting mixture was stirred at 0  $^\circ\,$  C for 30 min. To the solution added 4-methoxy-3-(phenylmethoxy)-benzaldehyde was (6.701 g, 27.7 mmol, 1.0 equiv) at 0  $^{\circ}$  C. The resulting mixture was allowed to warm to room temperature with stirring over 2 h, at which time the reaction mixture was treated with saturated aqueous ammonium chloride (70 mL). The resulting mixture was extracted twice with diethyl ether (50 mL). The combined organic extracts were washed with water (70 mL) and brine (70 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was dissolved in diethyl ether (40 mL) and CH<sub>2</sub>Cl<sub>2</sub> (40 mL). To the resulting solution was added hexane (300 mL), the resulting precipitate (triphenylphosphine oxide) was removed by filtration. The filtrate was concentrated under reduced pressure to give a colorless solid (13.96 g). This procedure was repeated once again to provide a crude enol ether **S16** as a colorless solid (8.94 g), which was used for the next step without further purification. A 1-L round-bottomed flask equipped with a Teflon-coated magnetic stirring bar was charged with the enol ether S16, THF (50 mL), and 6 M aqueous hydrochloric acid (50 mL). The resulting mixture was stirred at room temperature for 45 min, at which time the mixture was extracted with diethyl ether (100 mL) three times. The combined organic extracts were washed with water (100 mL) three times and brine (20 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude aldehyde S17, which was used for the next step without further purification.

A 500-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with the crude aldehyde S17 and methanol (50 mL). After cooling to  $\circ^{\circ}$  C, NaBH<sub>4</sub> (3.021 g, 79.9 mmol) was added potionwise to the flask at o<sup>°</sup> C. The resulting mixture was allowed to warm to room temperature with stirring over 2 h. Additional NaBH4 (1.496 g, 39.5 mmol) was added potionwise to the reaction mixture at o  $^{\circ}$  C. The resulting mixture was stirred for 17 h, at which time the reaction mixture was treated with water (80 mL) and concentrated under reduced pressure. The resulting mixture was extracted with ethyl acetate (80 mL) three times. The combined organic extracts were washed with brine (80 mL), dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/methyl acetate = 4:1 to 2:1, gradient) followed by SEC-HPLC to provide the title compound 27 as a colorless solid (2.346 g, 9.08 mmol, 33% over 3 steps), whose 'H NMR spectrum was identical to that reported in the literature.38

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16 17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

## TOTAL SYNTHESIS OF LAMELLARIN Z

## Ethyl 3-[3-(benzyloxy)-4-methoxyphenyl]-4-bromo-1-[(2-(trimethylsilyl)ethoxy)methyl]-1*H*-pyrrole-2-carboxylate (30).

A 100-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 8 (512.5 mg, 1.20 mmol, 1.0 equiv), Ba(OH)2.8H2O (2.27 g, 7.18 mmol, 6.0 equiv), 29 (557.0 mg, 1.64 mmol, 1.4 equiv), Pd<sub>2</sub>(dba)<sub>3</sub>·CHCl<sub>3</sub> µmol. (55.2 mg, 60.3 5 mol%). tris[4-(trifluoromethyl)phenyl]-phosphine (111.7 mg, 0.24 mmol, 20 mol%), water (2.4 mL), and 1,4-dioxane (9.6 mL). The flask was placed in a preheated oil bath and heated at 100  $^{\circ}$  C for 15 min, at which time the solution was treated with water (10 mL). The reaction mixture was extracted with diethyl ether (10 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/diethyl ether = 10:1 to hexane/diethyl ether = 5:1, gradient) and followed by preparative SEC-HPLC to provide the title product 30 as a pale yellow oil (202.0 mg, 0.36 mmol, 30%). R<sub>f</sub> = 0.23 (hexane/diethyl ether = 5:1); IR (ATR, cm<sup>-1</sup>): 1700, 1541, 1508, 1428, 1377, 1253, 1223, 1106, 1181, 1024, 856, 745, 702; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.45 (d, 2H, J = 7.3 Hz), 7.39-7.27 (m, 3H), 7.04 (s, 1H), 6.91 (d, 1H, J = 8.2 Hz, 6.88 (d, 1H, J = 1.8 Hz), 6.86 (dd, 1H, J = 8.2, 1.8 Hz), 5.64 (s, 2H), 5.15 (s, 2H), 4.00 (q, 2H, J = 7.1 Hz), 3.92 (s, 3H), 3.57 (t, 2H, J = 8.2 Hz), 0.96 (t, 3H, J = 7.1 Hz), 0.93 (q, 2H, J = 8.2 Hz), 0.00 (s, 9H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 160.7, 149.1, 147.3, 137.3, 132.8, 128.5, 127.8, 127.4, 126.7, 126.5, 123.5, 119.9, 116.6, 110.9, 99.4, 77.8, 71.1, 66.5, 60.1, 56.0, 17.8, 13.8, -1.4; HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for C<sub>27</sub>H<sub>34</sub>NO<sub>5</sub>SiNa 584.1267; Found 584.1291.

## Ethyl [3-(benzyloxy)-4-methoxyphenyl]-4-[3,4bis(benzyloxy)phenyl]-1-[(2-

# (trimethylsilyl)ethoxy)methyl]-1*H*-pyrrole-2-carboxylate (31).

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with **30** (110.0 mg, 0.196 mmol, 1.0 equiv),  $Ba(OH)_2 \cdot 8H_2O$  (370.1 mg, 1.17 mmol, 6.0 equiv), **23** (163.7 mg, 0.393 mmol, 2.0 equiv),  $Pd(PPh_3)_4$ (22.6 mg, 19.6 µmol, 10 mol%), water (0.4 mL), and 1,4-

dioxane (1.6 mL). The flask was placed in a preheated oil bath and heated at 100  $^{\circ}$  C for 15 min, at which time the solution was treated with water (2 mL). The reaction mixture was extracted with diethyl ether (3 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/diethyl ether = 5:1 to hexane/diethyl ether = 3:1) followed by preparative SEC-HPLC to provide the title product **31** as a pale yellow oil (109.5 mg, 0.142 mmol, 72%).  $R_f = 0.20$ (hexane/diethyl ether = 3:1); IR (ATR, cm<sup>-1</sup>): 2954, 1695, 1530, 1506, 1455, 1355, 1249, 1217, 1102, 1024, 859, 737, 697; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.43-7.23 (m, 15H), 7.06 (s, 1H), 6.84 (d, 1H, J = 8.2 Hz), 6.81 (d, 1H, J = 1.8 Hz), 6.78-7.75 (m, 2H), 6.64 (dd, 1H, J = 8.2, 2.3 Hz), 6.61 (d, 1H, J = 1.8 Hz), 5.69 (s, 2H), 5.10 (s, 2H), 5.01 (s, 2H), 4.77 (s, 2H), 4.01 (q, 2H, *J* = 7.2 Hz), 3.88 (s, 3H), 3.61 (t, 2H, J = 8.2 Hz), 0.96 (t, 3H, J = 7.2 Hz), 0.95 (t, 2H, J = 8.2 Hz), 0.00 (s, 9H);  ${}^{13}C{}^{1}H$  NMR (100 MHz, CDCl<sub>3</sub>): δ 161.6, 148.6, 148.5, 147.4, 147.2, 137.5, 137.3, 137.1, 131.5, 128.5, 128.44, 128.39, 127.9, 127.8, 127.72, 127.65, 127.30, 127.25, 127.1, 125.1, 124.5, 123.6, 120.6, 120.3, 116.8, 114.91, 114.88, 111.1, 77.7, 71.3, 70.92, 70.86, 66.3, 59.8, 56.0, 17.9, 13.9, -1.4; HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for C<sub>47</sub>H<sub>51</sub>NO<sub>7</sub>SiNa 792.3333; Found 792.3345.

## 8-(Benzyloxy)-1-(3,4-bis(benzyloxy)phenyl)-7methoxychromeno[3,4-b]pyrrol-4(3H)-one (S20).

A 10-mL screw-top test tube equipped with a Teflon-coated magnetic stirring bar was charged with compound **31** (108.5 mg, 0.141 mmol, 1.0 equiv), 18-crown-6 (74.3 mg, 0.281 mmol, 2.0 equiv), KOH (79.0 mg, 1.41 mmol, 10 equiv), and EtOH (705  $\mu$ L). The test tube was placed in a preheated aluminum block and heated at reflux for 19 h, at which time the solution was treated with 1 M aqueous hydrochloric acid. The desired carboxylic acid **S18** was collected by filtration as a yellow solid (108.4 mg), which was used for the next step without further purification.

A 50-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with the crude S18 (108.4 mg) in anhydrous ethyl acetate (14 mL). To the solution was added Pb(OAc)<sub>4</sub> (93.8 mg, 0.212 mmol). The flask was placed in a preheated oil bath and heated at reflux for 20 min. After  $Pb(OAc)_4$  (31.5 mg, 0.0710 mmol) was added to the solution, and the reaction mixture was heated at reflux for 15 min. Pinacol (43.2 mg, 0.366 mmol) was added to the flask, and resulting mixture was treated with water (14 mL). The reaction mixture was extracted with diethyl ether (10 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/diethyl ether = 3:1) to provide the corresponding product S19 as a yellow solid (54.1 mg, 73.1 µmol).

A 10-mL screw-top test tube equipped with a Teflon-coated magnetic stirring bar was charged with compound **S19** (54.1 mg, 73.1  $\mu$ mol, 1.0 equiv) and CH<sub>2</sub>Cl<sub>2</sub> (1.8 mL). TFA (400  $\mu$ L, 5.22 mmol) was added to the flask. The resulting mixture was stirred at room temperature for 15 min. After the reaction mixture was concentrated under reduced pressure, the flask was charged with anhydrous THF (585  $\mu$ L), water (146  $\mu$ L),

2

3

4

5

6

7

8

9

25

26

27

NH<sub>2</sub>OH·HCl (4.7 mg, 68 µmol, 0.92 equiv), and Na<sub>2</sub>CO<sub>3</sub> (23.8 mg, 0.225 mmol, 3.1 equiv). The reaction mixture was stirred at room temperature for 3 h, at which time NH2OH·HCl (4.9 mg, 71 µmol, 1.0 equiv) and Na2CO3 (22.3 mg, 0.210 mmol, 3.0 equiv) were added to the flask. After stirring at room temperature for 4 h, the reaction mixture was treated with water (0.6 mL). The resulting mixture was extracted with ethyl acetate (1 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, 10 which was washed with diethyl ether (5 mL) to provide the 11 title compound **S20** as a pale yellow solid (36.1 mg, 59.2 µmol, 12 42% over 3 steps).  $R_f = 0.31$  (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 100:1); IR (ATR, 13 cm<sup>-1</sup>): 3237, 1696, 1549, 1494, 1402, 1272, 1192, 1026, 736, 696; 14 <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 9.56 (br s, 1H), 7.51-7.45 (m, 15 4H), 7.38-7.28 (m, 6H), 7.23-7.17 (m, 3H), 7.15 (d, 1H, J = 2.3 16 Hz), 7.14 (s, 1H), 7.09-7.05 (m, 3H), 7.02 (d, 1H, J = 8.2 Hz), 17 6.97 (dd, 1H, J = 8.2, 1.8 Hz), 6.95 (s, 1H), 5.25 (s, 2H), 5.18 (s, 2H), 4.82 (s, 2H), 3.92 (s, 3H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, 18 CDCl<sub>3</sub>): δ 156.5, 149.8, 149.2, 148.7, 146.3, 144.6, 137.3, 137.1, 19 136.4, 128.71, 128.68, 128.5, 128.1, 128.0, 127.9, 127.6, 127.5, 127.40, 20 127.35, 126.7, 123.0, 121.4, 116.7, 116.5, 115.0, 110.5, 107.4, 101.2, 21 71.50, 71.46, 70.7, 56.3 (one aromatic carbon signal is missing 22 due to overlapping); HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd 23 for C<sub>39</sub>H<sub>31</sub>NO<sub>6</sub>Na 632.2049; Found 632.2035. 24

## 8-(Benzyloxy)-3-[3-(benzyloxy)-4-methoxyphenethyl]-1-[3,4-bis(benzyloxy)phenyl]-7-methoxychromeno[3,4*b*]pyrrol-4(3*H*)-one (32).

28 A 10-mL screw-top test tube equipped with a Teflon-coated 29 magnetic stirring bar was charged with compound S20 (38.3 3-benzyloxy-4equiv), mg, 62.8 μmol, 1.0 30 methoxybenzeneethanol (27) (24.3 mg, 94.2 µmol, 1.5 equiv), 31 and PPh<sub>3</sub> (49.3 mg, 0.188 mmol, 3.0 equiv). After the flask was 32 evacuated and backfilled with N<sub>2</sub>, anhydrous THF (320 µL) 33 and DIAD (1.9 M, 100 µL, 0.19 mmol, 3.0 equiv) were added to 34 the flask. The solution was stirred at room temperature for 15 35 min, the resulting mixture was concentrated under reduced 36 pressure to give a crude product, which was purified by silica 37 gel column chromatography (hexane/ $CH_2Cl_2 = 1:3$  to  $CH_2Cl_2$ , 38 gradient). The amorphous was washed twice with methanol 39 (1 mL) to provide the title compound 32 as a white amorphous (46.0 mg, 54.1  $\mu$ mol, 86%). Rf = 0.21 (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 40 1:3); IR (ATR, cm<sup>-1</sup>): 1707, 1513, 1455, 1419, 1260, 1235, 1137, 1025, 41 739, 697; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.47-7.43 (m, 4H), 42 7.37-7.28 (m 6H), 7.25-7.16 (m, 7H), 7.10 (s, 1H), 7.08-7.04 (m, 43 2H), 6.97 (d, 1H, J = 8.2 Hz), 6.94 (s, 1H), 6.91 (d, 1H, J = 2.344 Hz), 6.83 (dd, 1H, J = 8.2, 1.8 Hz), 6.79 (d, 1H, J = 8.2 Hz), 6.67 45 (dd, 1H, J = 8.2, 1.8 Hz), 6.51 (d, 1H, J = 1.8 Hz), 6.49 (s, 1H),46 5.22 (s, 2H), 5.10 (s, 2H), 4.95 (s, 2H), 4.78 (s, 2H), 4.54 (t, 2H, 47 *J* = 6.8 Hz), 3.92 (s, 3H), 3.81 (s, 3H), 3.02 (t, 2H, *J* = 6.8 Hz); 48 <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 155.5, 149.7, 149.1, 148.54, 49 148.51, 148.1, 146.5, 144.3, 137.3, 137.1, 137.0, 136.4, 132.2, 130.5, 50 128.7, 128.6, 128.54, 128.46, 128.02, 127.99, 127.9, 127.8, 127.6, 51 127.35, 127.31, 127.1, 122.9, 121.6, 118.9, 116.5, 115.0, 114.8, 112.0, 52 110.4, 107.3, 101.0, 71.5, 71.3, 70.9, 70.6, 56.3, 56.1, 51.1, 37.6 (four 53 aromatic carbon signals are missing due to overlapping); HRMS (ESI/TOF) m/z:  $[M + Na]^+$  Calcd for  $C_{55}H_{47}NO_8Na$ 54 872.3229; Found 872.3199. 55

## Lamellarin Z.

56

57 58 59

60

A 25-mL two-necked pear-shaped flask equipped with a Teflon-coated magnetic stirring bar was charged with compound 32 (45.0 mg, 52.9 µmol, 1.0 equiv) and anhydrous CH<sub>2</sub>Cl<sub>2</sub> (4.4 mL). After the resulting solution was cooled to -40  $^{\circ}$  C, a solution of PIFA (26.5 mg, 61.6  $\mu$ mol, 1.2 equiv) and BF3·OEt2 (17 µL, 0.14 mmol, 2.6 equiv) in anhydrous CH2Cl2 (0.9 mL) was added dropwise to the flask. The reaction mixture was stirred at  $-40^{\circ}$  C for 1.5 h, at which time the resulting mixture was treated with 2 M aqueous NH<sub>3</sub> (5 mL). The mixture was extracted with CH2Cl2 (5 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was washed with diethyl ether (6 mL) to provide the corresponding product S21 as a colorless solid (36.9 mg).

A 25-mL two-necked pear-shaped flask equipped with a Teflon-coated magnetic stirring bar was charged with compound S21 (36.1 mg, 42.6 µmol, 1.0 equiv), Pd/C (35.8 mg, 100 wt%), EtOH (0.7 mL), and ethyl acetate (0.7 mL). After the flask was evacuated and backfilled with H<sub>2</sub> three times, the resulting mixture was stirred at room temperature for 17 h, at which time the resulting mixture was filtered through a pad of Celite with hot methanol (15 mL) followed by DMSO (3 mL). The filtrate was concentrated under reduced pressure to provide lamellarin Z as a gray solid (20.7 mg, 42.5 µmol, 80% over 2 steps), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra (DMSO-*d*<sub>6</sub>) were identical to those reported in the literature.<sup>27</sup>  $R_f = 0.43$  $(CH_2Cl_2/methanol = 10:1); M.p. > 250^{\circ} C; IR (ATR, cm^{-1}): 3410,$ 1684, 1587, 1488, 1425, 1276, 1245, 1206, 1158, 1041, 1024, 870, 761; <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 9.41 (s, 1H), 9.19 (s, 1H), 9.14 (s, 1H), 8.98 (s, 1H), 7.00 (s, 1H), 6.94 (d, 1H, I = 7.8 Hz), 6.76 (s, 1H), 6.73 (s, 1H), 6.71-6.67 (m, 2H), 6.54 (s, 1H), 4.72-4.63 (m, 1H), 4.55-4.46 (m, 1H), 3.81 (s, 3H), 3.28 (s, 3H),  $3.04^{-}2.93$  (m, 2H);  ${}^{13}C{}^{1}H$  NMR (100 MHz, DMSO- $d_6$ ):  $\delta$  154.5, 147.9, 147.0, 146.2, 146.0, 145.4, 144.7, 142.8, 136.1, 127.2, 127.0, 125.5, 121.6, 118.2, 117.8, 116.7, 115.3, 114.8, 112.5, 110.3, 109.3, 108.4, 100.7, 55.9, 54.7, 42.0, 27.6; HRMS (ESI/TOF) m/z:  $[M + Na]^+$ Calcd for C<sub>27</sub>H<sub>21</sub>NO<sub>8</sub>Na 510.1165; Found 510.1165.

## 2-Benzyloxy-4-bromo-1-methoxybenzene (S22).

A 100-mL round-bottomed flask equipped with a Tefloncoated magnetic stirring bar was charged with 5-bromo-2methoxy-phenol (3.05 g, 15.0 mmol, 1.0 equiv), K<sub>2</sub>CO<sub>3</sub> (4.15 g, 30.0 mmol, 2.0 equiv), benzyl bromide (2.57 g, 15.0 mmol, 1.0 equiv), and acetonitrile (15 mL). The flask was placed in a preheated oil bath and heated at reflux for 4 h. After cooling to room temperature, the resulting mixture was filtered through a pad of Celite. The filtrate was concentrated under reduced pressure to give the title product S22 as a colorless solid (4.35 g, 14.8 mmol, 99%), whose <sup>1</sup>H and <sup>13</sup>C NMR spectra were identical to those reported in the literature.<sup>39</sup>

## 2-[3-(Benzyloxy)-4-methoxyphenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (29).

A flame-dried 100-mL two-necked round-bottomed flask equipped with a Teflon-coated magnetic stirring bar and a rubber septum was charged with 2-benzyloxy-4-bromo-1methoxybenzene (S22) (880.2 mg, 3.00 mmol, 1.0 equiv) and anhydrous THF (15 mL). After the solution was cooled to -78  $^{\circ}$  C, the reaction mixture was treated with *n*-BuLi (1.59 M in n-hexane, 1.9 mL, 3.0 mmol, 1.0 equiv) and stirred at - $78^{\circ}$  C for 5 min. To the solution was added *i*-PrOBpin (658.3) mg, 3.54 mmol, 1.2 equiv) at  $-78^{\circ}$  C. After stirring at  $-78^{\circ}$  C for 10 min, the reaction mixture was allowed to warm to room temperature with stirring over 1 h, at which time the reaction mixture was treated with saturated aqueous ammonium chloride (15 mL). The reaction mixture was extracted with diethyl ether (15 mL) three times. The combined organic extracts were washed with water and brine, dried over sodium sulfate, and filtered. The filtrate was concentrated under reduced pressure to give a crude product, which was purified by silica gel column chromatography (hexane/ $CH_2Cl_2 = 1:1$  to hexane/diethyl ether = 10:1, gradient) to provide the title product 29 as a colorless solid (795.6 mg, 2.34 mmol, 78%). Rf = 0.21 (hexane/CH<sub>2</sub>Cl<sub>2</sub> = 1:1); M.p. 112-113  $^{\circ}$  C; IR (ATR, cm<sup>-1</sup>): 2978, 1600, 1416, 1372, 1352, 1295, 1257, 1219, 1137, 1026, 966, 855, 737, 698, 685; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 7.51-7.28 (m, 7H), 6.91 (d, 1H, J = 8.2 Hz), 5.15 (s, 2H), 3.89 (s, 3H), 1.33 (s, 12H); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ 152.4, 147.8, 137.2, 129.1, 128.5, 127.8, 127.6, 120.8, 119.3, 111.1, 83.6, 70.9, 55.8, 24.9; HRMS (DART/TOF) m/z:  $[M + H]^+$  Calcd for  $C_{20}H_{26}BO_4$ 341.1924; Found 341.1909.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.xxxxxx. Tabular comparisons of NMR data with the natural products and copies of <sup>1</sup>H and <sup>13</sup>C NMR spectra for all new compounds (PDF).

## AUTHOR INFORMATION

#### Corresponding Author

Kentaro Okano\*

\*E-mail: okano@harbor.kobe-u.ac.jp

Notes

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

The authors declare no competing financial interest.

## ACKNOWLEDGMENT

This work was supported by JSPS KAKENHI Grant Numbers (JP16K05774, JP16H01153, JP18H04413, JP19H02717) and the Sasakawa Scientific Research Grant. We thank Mr. Yoshiki Yamane for performing some preliminary experiments. We also thank Dr. Alison McGonagle from Edanz Group (https://en-author-services.edanzgroup.com/) for editing a draft of this manuscript.

## REFERENCES

(1) Zhang, H. W.; Dong, M. L.; Chen, J. W.; Wang, H.; Tenney, K.; Crews, P. Bioactive secondary metabolites from the marine sponge genus *Agelas. Mar. Drugs* **2017**, *15*, 351. (b) Wood, J. M.; Furkert, D. P.; Brimble, M. A. 2-Formylpyrrole natural products: origin, structural diversity, bioactivity and synthesis. *Nat. Prod. Rep.* **2019**, *36*, 289-306.

(2) (a) Fan, H.; Peng, J.; Hamann, M. T.; Hu, J.-F. Lamellarins and related pyrrole-derived alkaloids from marine organisms. *Chem. Rev.* **2008**, *108*, 264-287. (b) Iwao, M.; Fukuda, T.; Ishibashi, F. Synthesis and biological activity of lamellarin alkaloids: an overview. *Heterocycles* **2011**, *83*, 491-529.

(3) Andersen, R. J.; Faulkner, D. J.; He, C. H.; Van Duyne, G. D.; Clardy, J. Metabolites of the marine prosobranch mollusk *Lamellaria* sp. *J. Am. Chem. Soc.* **1985**, *107*, 5492-5495.

(4) Warabi, K.; Matsunaga, S.; van Soest, R. W. M.; Fusetani, N. Dictyodendrins A-E, the frst telomerase-inhibitory marine natural products from the sponge *Dictyodendrilla verongiformis. J. Org. Chem.* **2003**, *68*, 2765-2770.

(5) (a) Blunt, J. W.; Copp, B. R.; Keyzers, R. A.; Munro, M. H. G.; Prinsep, M. R. Marine natural products. *Nat. Prod. Rep.* **2013**, 30, 237-323. (b) Khadka, D. B.; Cho, W. J. Topoisomerase inhibitors as anticancer agents: a patent update. *Expert Opin. Ther. Pat.* **2013**, 23, 1033-1056. (c) Bailly, C. Anticancer properties of lamellarins. *Mar. Drugs* **2015**, *13*, 1105-1123.

(6) For a recent review: (a) Imbri, D.; Tauber, J.; Opatz, T. Synthetic approaches to the lamellarins-a comprehensive review. Mar. Drugs 2014, 12, 6142-6177. (b) Heim, A.; Terpin, A.; Steglich, W. Biomimetic synthesis of lamellarin G trimethyl ether. Angew. Chem, Int. Ed. 1997, 36, 155-156. (c) Boger, D. L.; Boyce, C. W.; Labroli, M. A.; Sehon, C. A.; Jin, Q. Total syntheses of ningalin A, lamellarin O, lukianol A, and permethyl storniamide A utilizing heterocyclic azadiene Diels-Alder reactions. J. Am. Chem. Soc. 1999, 121, 54-62. (d) Liu, J.-H.; Yang, Q.-C.; Mak, T. C. W.; Wong, H. N. C. Highly regioselective synthesis of 2,3,4-trisubstituted 1H-pyrroles: a formal total synthesis of lukianol A. J. Org. Chem. 2000, 65, 3587-3595. (e) Gupton, J. T.; Clough, S. C.; Miller, R. B.; Lukens, J. R.; Henry, C. A.; Kanters, R. P. F.; Sikorski, J. A. The application of vinylogous iminium salt derivatives to the synthesis of ningalin B hexamethyl ether. Tetrahedron 2003, 59, 207-215. (f) Peschko, C.; Winklhofer, C.; Terpin, A.; Steglich, W. Biomimetic syntheses of lamellarin and lukianol-type alkaloids. Synthesis 2006, 3048-3057. (g) Ohta, T.; Fukuda, T.; Ishibashi, F.; Iwao, M. Design and synthesis of lamellarin D analogues targeting topoisomerase I. J. Org. Chem. 2009, 74, 8143-8153. (h) Li, Q.; Jiang, J.; Fan, A.; Cui, Y.; Jia, Y. Total synthesis of lamellarins D, H, and R and ningalin B. Org. Lett. 2011, 13, 312-315. (i) Lade, D. M.; Pawar, A. B.; Mainkar, P. S.; Chandrasekhar, S. Total synthesis of lamellarin D trimethyl ether, lamellarin D, and lamellarin H. J. Org. Chem. 2017, 82, 4998-5004.

(7) (a) Banwell, M. G.; Flynn, B. L.; Hamel, E.; Hockless, D. C. R. Convergent syntheses of the pyrrolic marine natural products lamellarin-O, lamellarin-Q, lukianol-A and some more highly oxygenated congeners. Chem. Commun. 1997, 207-208. (b) Ishibashi, F.; Miyazaki, Y.; Iwao, M. Total syntheses of lamellarin D and H. The first synthesis of lamellarin-class marine alkaloids. Tetrahedron 1997, 53, 5951-5962. (c) Ridley, C. P.; Reddy, M. V. R.; Rocha, G.; Bushman, F. D.; Faulkner, D. J. Total synthesis and evaluation of lamellarin  $\alpha$  20-sulfate analogues. *Biorg. Med.* Chem. 2002, 10, 3285-3290. (d) Pla, D.; Marchal, A.; Olsen, C. A.; Albericio, F.; Álvarez, M. Modular total synthesis of lamellarin D. J. Org. Chem. 2005, 70, 8231-8234. (e) Imbri, D.; Tauber, J.; Opatz, T. A high-yielding modular access to the lamellarins: synthesis of lamellarin G trimethyl ether, lamellarin n and dihydrolamellarin n. Chem. Eur. J. 2013, 19, 15080-15083. (f) Ueda, K.; Amaike, K.; Maceiczyk, R. M.; Itami, K.; Yamaguchi, J. β-Selective C-H arylation of pyrroles leading to concise syntheses of lamellarins C and I. J. Am. Chem. Soc. 2014, 136, 13226-13232. (g) Zheng, K.-L.; You, M.-Q.; Shu, W.-M.; Wu, Y.-D.; Wu, A.-X. Acid-mediated intermolecular [3 + 2] cycloaddition toward pyrrolo[2,1*a*]isoquinolines: total synthesis of the lamellarin core and lamellarin G trimethyl ether. Org. Lett. 2017, 19, 2262-2265.

(8) Representative reviews on halogen dance: (a) Schlosser, M. The organometallic approach to molecular diversity - halogens as helpers. *Eur. J. Org. Chem.* **2001**, 3975-3984. (b) Schnürch, M.; Spina, M.; Khan, A. F.; Mihovilovic, M. D.; Stanetty, P. Halogen dance reactions-a review. *Chem. Soc. Rev.* **2007**, *36*, 1046-1057. (c) Erb, W.; Mongin, F. Halogen 'dance': a way to extend the boundaries of arene deprotolithiation. *Tetrahedron* **2016**, *72*, 4973-4988, and references cited therein.

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

(9) Representative total syntheses using halogen dance: (a) Comins, D. L.; Saha, J. K. Concise synthesis of mappicine ketone and (±)-mappicine. J. Org. Chem. 1996, 61, 9623-9624. (b) Arzel, E.; Rocca, P.; Grellier, P.; Labaeïd, M.; Frappier, F.; Guéritte, F.; Gaspard, C.; Marsais, F.; Godard, A.; Quéguiner, G. New synthesis of benzo-δ-carbolines, cryptolepines, and their salts: In vitro cytotoxic, antiplasmodial, and antitrypanosomal activities of  $\delta$ caribolines, benzo-δ-carbolines, and cryptolepines. J. Med. Chem. 2001, 44, 949-960. (c) Bouillon, A.; Voisin, A. S.; Robic, A.; Lancelot, J.-C.; Collot, V.; Rault, S. An efficient two-step total synthesis of the quaterpyridine nemertelline. J. Org. Chem. 2003, 68, 10178-10180. (d) Stangeland, E. L.; Sammakia, T. Use of thiazoles in the halogen dance reaction: application to the total synthesis of WS75624 B. J. Org. Chem. 2004, 69, 2381-2385. (e) Keaton, K. A.; Phillips, A. J. Titanium(II)-mediated cyclizations of (silyloxy)enynes: a total synthesis of (-)-7-demethylpiericidin A1. J. Am. Chem. Soc. 2006, 128, 408-409. (f) Ohtawa, M.; Ogihara, S.; Sugiyama, K.; Shiomi, K.; Harigaya, Y.; Nagamitsu, T.; Omura, S. Enantioselective total synthesis of atpenin A5. J. Antibiot. 2009, 62, 289-294. (g) Schäckermann, J.-N.; Lindel, T. Synthesis and photooxidation of the trisubstituted oxazole fragment of the marine natural product salarin C. Org. Lett. 2017, 19, 2306-2309. (h) Marcus Vinicius Nora de, S. Halogen dance reaction and its application in organic synthesis. Curr. Org. Chem. 2007, 11, 637-646, and references cited therein.

(10) (a) Knorr, L. Einwirkung des Diacetbernsteinsäureesters auf Ammoniak und primäre Aminbasen. *Ber. Deut. Chem. Ges.* 1885, *18*, 299-311. (b) Paal, C. Synthese von Thiophen- und Pyrrolderivaten. *Ber. Deut. Chem. Ges.* 1885, *18*, 367-371. (c) Minetto, G.; Raveglia, L. F.; Sega, A.; Taddei, M. Microwave-assisted Paal-Knorr reaction - Three-step regiocontrolled synthesis of polysubstituted furans, pyrroles and thiophenes. *Eur. J. Org. Chem.* 2005, 5277-5288. (d) Khaghaninejad, S.; Heravi, M. M. *Advances in Heterocyclic Chemistry, Vol 111*, Katritzky, A. R., Ed. Elsevier Academic Press Inc: San Diego, 2014; Vol. 111, pp 95-146.

(11) (a) Okano, K.; Sunahara, K.; Yamane, Y.; Hayashi, Y.; Mori,
A. One-pot halogen dance/Negishi coupling of dibromothiophenes for regiocontrolled synthesis of multiply arylated thiophenes. *Chem. Eur. J.* 2016, 22, 16450-16454. (b) Miyagawa, N.; Murase, Y.; Okano, K.; Mori, A. Benzoxazole-directed halogen dance of bromofuran. *Synlett* 2017, 28, 1106-1110. (c) Hayashi, Y.; Okano, K.; Mori, A. Synthesis of thieno[3,2-b]indoles via halogen dance and ligand-controlled one-pot sequential coupling reaction. *Org. Lett.* 2018, 20, 958-961. (d) Okano, K.; Murase, Y.; Mori, A. Synthesis of furan-fused silole and phosphole by onepot halogen dance/homocoupling of bromofurfural derivative. *Heterocycles* 2019, 99, 1444-1451.

(12) Bey, E.; Marchais-Oberwinkler, S.; Negri, M.; Kruchten, P.;
Oster, A.; Klein, T.; Spadaro, A.; Werth, R.; Frotscher, M.; Birk, B.; Hartmann, R. W. New Insights into the SAR and binding modes of bis(hydroxyphenyl)thiophenes and -benzenes: influence of additional substituents on 17β-hydroxysteroid dehydrogenase type 1 (17β-HSD1) inhibitory activity and selectivity. *J. Med. Chem.* 2009, 52, 6724<sup>-6</sup>743.

(13) (a) Yamane, Y.; Sunahara, K.; Okano, K.; Mori, A. Magnesium bisamide-mediated halogen dance of bromothiophenes. *Org. Lett.* 2018, 20, 1688-1691. (b) Mari, D.; Miyagawa, N.; Okano, K.; Mori, A. Regiocontrolled halogen dance of bromothiophenes and bromofurans. *J. Org. Chem.* 2018, *83*, 14126-14137.

(14) Instead of the ester group, an oxazoline was also an effective substituent on the pyrrole for the halogen dance, to provide the corresponding  $\beta$ , $\beta$ '-dibromopyrrole. However, introduction of the aryl groups did not proceed smoothly. (15) The desired compound **8** was not obtained at  $-40^{\circ}$  C or o  $^{\circ}$  C. Prolonged reaction time (1 h) resulted in significant reduction in the yield of compound **8** (<10% yield).

(16) The regioselectivity was determined by further transformation of the ethyl ester to its lactone. The chemical shift of the methylene protons of the ethyl ester was shifted to 4.0 ppm from 4.3 ppm (the methylene protons of the ethyl ester of compound **8**), indicating that the installed aryl group shielded the methylene protons.

(17) Handy reported Suzuki-Miyaura cross coupling of a quite similar  $\beta$ , $\beta$ '-dibromopyrrole; however, they obtained almost equimolar amounts of the monoarylated pyrrole and the diarylated pyrrole, even with substoichiometric amount (0.6 equiv to the  $\beta$ , $\beta$ '-dibromopyrrole) of an arylboronic acid, see: (a) Handy, S. T.; Zhang, Y. Regioselective couplings of dibromopyrrole esters. Synthesis 2006, 3883-3887. α-Selective Suzuki-Miyaura coupling of  $\alpha$ , $\beta$ -dibromopyrroles, see: (b) Handy, S. T.; Sabatini, J. J. Regioselective dicouplings: application to differentially substituted pyrroles. Org. Lett. 2006, 8, 1537-1539. (c) Zhang, Y.; Handy, S. T. A solvent-induced reversal of regioselectivity in the Suzuki coupling of pyrrole esters. Open Org. Chem. J. 2008, 2, 58-64. For recent reviews, see: (d) Rossi, R.; Bellina, F.; Lessi, M. Selective palladium-catalyzed Suzuki-Miyaura reactions of polyhalogenated heteroarenes. Adv. Synth. Catal. 2012, 354, 1181-1255. (e) Fairlamb, I. J. S. Regioselective (site-selective) functionalisation of unsaturated halogenated nitrogen, oxygen and sulfur heterocycles by Pd-catalysed cross-couplings and direct arylation processes. Chem. Soc. Rev. 2007, 36, 1036-1045. (f) Banwell, M. G.; Goodwin, T. E.; Ng, S.; Smith, J. A.; Wong, D. J. Palladium-catalysed cross-coupling and related reactions involving pyrroles. Eur. J. Org. Chem. 2006, 3043-3060.

(18) (a) Sheina, E. E.; Liu, J.; Iovu, M. C.; Laird, D. W.; McCullough, R. D. Chain growth mechanism for regioregular nickel-initiated cross-coupling polymerizations. Macromolecules 2004, 37, 3526-3528. (b) Yoshikai, N.; Matsuda, H.; Nakamura, E. Ligand exchange as the first irreversible step in the nickelcatalyzed cross-coupling reaction of Grignard reagents. J. Am. Chem. Soc. 2008, 130, 15258-15259. (c) Miyakoshi, R.; Yokoyama, A.; Yokozawa, T. Catalyst-transfer polycondensation. Mechanism of Ni-catalyzed chain-growth polymerization leading to welldefined poly(3-hexylthiophene). J. Am. Chem. Soc. 2005, 127, 17542-17547. (d) Yokozawa, T.; Ohta, Y. Transformation of stepgrowth polymerization into living chain-growth polymerization. Chem. Rev. 2016, 116, 1950-1968. (e) Yokozawa, T.; Harada, N.; Sugita, H.; Ohta, Y. Intramolecular catalyst transfer on a variety of functional groups between benzene rings in a Suzuki-Miyaura coupling reaction. Chem. Eur. J. 2019, 25, 10059-10062.

(19) Apart from Yokozawa's report (Ref. 18e) using 1,4dibromobenzene derivatives, two bromo groups on the adjacent positions may facilitate the catalyst transfer to provide the diarylated pyrrole **13**.

(20) Seo, T.; Ishiyama, T.; Kubota, K.; Ito, H. Solid-state Suzuki - Miyaura cross-coupling reactions: olefin-accelerated C - C coupling using mechanochemistry. *Chem. Sci.* **2019**, *10*, 8202-8210.

(21) Isolation of lukianols A and B: (a) Yoshida, W. Y.; Lee, K. K.; Carroll, A. R.; Scheuer, P. J. A Complex pyrrolo-oxazinone and its iodo derivative isolated from a tunicate. *Helv. Chim. Acta* **1992**, *75*, 1721-1725. Total syntheses of lukianol A: (b) Fürstner, A.; Weintritt, H.; Hupperts, A. A new, titanium-mediated approach to pyrroles: first synthesis of lukianol A and lamellarin O dimethyl ether. *J. Org. Chem.* **1995**, *60*, 6637-6641. (c) Liu, J.-H.; Yang, Q.-C.; Mak, T. C. W.; Wong, H. N. C. Highly regioselective synthesis of 2,3,4-trisubstituted 1*H*-pyrroles: a formal total synthesis of lukianol A. *J. Org. Chem.* **2000**, *65*, 3587-3595. (d) Kim,

S.; Son, S.; Kang, H. Efficient syntheses of 2-carbomethoxy-3,4-disubstituted pyrroles by the condensation of vinylogous amides with aminomalonate. *Bull. Korean Chem. Soc.* 2001, 22, 1403-1406.
(e) Lu, Y.; Arndtsen, B. A. A direct phosphine-mediated synthesis of pyrroles from acid chlorides and α,β-unsaturated imines. *Org. Lett.* 2009, *II*, 1369-1372. Total syntheses of lukianols A and B: (f) Takamura, K.; Matsuo, H.; Tanaka, A.; Tanaka, J.; Fukuda, T.; Ishibashi, F.; Iwao, M. Total synthesis of the marine natural products lukianols A and B. *Tetrahedron* 2013, *69*, 2782-2788. Formal syntheses of lukianol A: (g) Smith, J. A.; Ng, S.; White, J. The regioselective synthesis of aryl pyrroles. *Org. Biomol. Chem.* 2006, *4*, 2477-2482. (h) Hinze, C.; Kreipl, A.; Terpin, A.; Steglich, W. Synthesis of simple 3,4-diarylpyrrole-2,5-dicarboxylic acids and lukianol A by oxidative condensation of 3-arylpyruvic acids with ammonia. *Synthesis* 2007, 608-612.

(22) The use of TBAF led to the significant reduction in yield of the desired product, due to hydrolysis of the ethyl ester. TFApromoted removal of the trimethylsilylethyl group: (a) Yamaguchi, J.; Seiple, I. B.; Young, I. S.; O'Malley, D. P.; Maue, M.; Baran, P. S. Synthesis of 1,9-dideoxy-pre-axinellamine. *Angew. Chem., Int. Ed.* **2008**, *47*, 3578-3580. (b) Subsequent treatment of the resulting hydroxymethylated pyrrole with aqueous sodium carbonate was effective for the high-yielding process.

(23) Laganis, E. D.; Chenard, B. L. Metal silanoates: organic soluble equivalents for O<sup>-2</sup>. *Tetrahedron Lett.* **1984**, *25*, 5831-5833.

(24) Okano, K.; Okuyama, K.; Fukuyama, T.; Tokuyama, H. Mild debenzylation of aryl benzyl ether with BCl<sub>3</sub> in the presence of pentamethylbenzene as a non-Lewis-basic cation scavenger. *Synlett* **2008**, 1977-1980.

(25) Isolation of ningalin B: (a) Kang, H. J.; Fenical, W. Ningalins A-D: novel aromatic alkaloids from a Western Australian ascidian of the genus Didemnum. J. Org. Chem. 1997, 62, 3254-3262. Total syntheses of ningalin B: (b) Boger, D. L.; Soenen, D. R.; Boyce, C. W.; Hedrick, M. P.; Jin, Q. Total synthesis of ningalin B utilizing a heterocyclic azadiene Diels-Alder reaction and discovery of a new class of potent multidrug resistant (MDR) reversal agents. J. Org. Chem. 2000, 65, 2479-2483. (c) Bullington, J. L.; Wolff, R. R.; Jackson, P. F. Regioselective preparation of 2-substituted 3,4-diaryl pyrroles: a concise total synthesis of ningalin B. J. Org. Chem. 2002, 67, 9439-9442. (d) Hasse, K.; Willis, A. C.; Banwell, M. G. A Total synthesis of the marine alkaloid ningalin B from (S)-proline. Aust. J. Chem. 2009, 62, 683-691. (e) Wu, C.-K.; Weng, Z.; Yang, D.-Y. One-pot construction of 1phenylchromeno[3,4-b]pyrrol-4(3H)-one: application to total synthesis of ningalin B and a pyrrolocoumarin-based electrochromic switch. Org. Lett. 2019, 21, 5225-5228. Formal syntheses of ningalin B: (f) Iwao, M.; Takeuchi, T.; Fujikawa, N.; Fukuda, T.; Ishibashi, F. Short and flexible route to 3,4-diarylpyrrole marine alkaloids: syntheses of permethyl storniamide A, ningalin B, and lamellarin G trimethyl ether. Tetrahedron Lett. 2003, 44, 4443-4446. (g) Gupton, J. T.; Giglio, B. C.; Eaton, J. E.; Rieck, E. A.; Smith, K. L.; Keough, M. J.; Barelli, P. J.; Firich, L. T.; Hempel, J. E.; Smith, T. M.; Kanters, R. P. F. The application of vinylogous iminium salt derivatives to efficient formal syntheses of the marine alkaloids lamellarin G trimethyl ether and ningalin B. Tetrahedron 2009, 65, 4283-4292. (h) Kumar, V.; Awasthi, A.; Metya, A.; Khan, T. A metal-free domino process for regioselective synthesis of 1,2,4-trisubstituted pyrroles: application toward the formal synthesis of ningalin B. J. Org. Chem. 2019, 84, 11581-11595. (i) Watanabe, T.; Mutoh, Y.; Saito, S. Synthesis of lactone-fused

pyrroles by ruthenium-catalyzed 1,2-carbon migrationcycloisomerization. *Org. Biomol. Chem.* **2020**, *18*, 81-85.

(26) Isolation of lamellarin S: (a) Urban, S.; Capon, R. J. Lamellarin-S: A new aromatic metabolite from an Australian tunicate, *Didemnum* sp. *Aust. J. Chem.* **1996**, *49*, 711-713. Total synthesis of lamellarin S: (b) Hasse, K.; Willis, A. C.; Banwell, M. G. Modular total syntheses of lamellarin G trimethyl ether and lamellarin S. *Eur. J. Org. Chem.* **2011**, 88-99.

(27) Isolation of lamellarin Z: Davis, R. A.; Carroll, A. R.; Pierens, G. K.; Quinn, R. J. New lamellarin alkaloids from the Australian ascidian, *Didemnum chartaceum*. *J. Nat. Prod.* **1999**, *62*, 419-424.

(28) (a) Kita, Y.; Tohma, H.; Hatanaka, K.; Takada, T.; Fujita, S.; Mitoh, S.; Sakurai, H.; Oka, S. Hypervalent iodine-induced nucleophilic substitution of *para*-substituted phenol ethers. Generation of cation radicals as reactive intermediates. *J. Am. Chem. Soc.* **1994**, *116*, 3684-3691. (b) Moreno, I.; Tellitu, I.; San Martín, R.; Domínguez, E. A new entrance to the preparation of phenanthrene and phenanthrenoid heterocycles. *Synlett* **2001**, 1161-1163.

(29) Lubriks, D.; Sokolovs, I.; Suna, E. Iodonium salts are key intermediates in Pd-catalyzed acetoxylation of pyrroles. *Org. Lett.* **2011**, *1*3, 4324<sup>-</sup>4327.

(30) Chakraborti, A. K.; Chankeshwara, S. V. Counterattack mode differential acetylative deprotection of phenylmethyl ethers: applications to solid phase organic reactions. *J. Org. Chem.* **2009**, *74*, 1367-1370.

(31) Ebisawa, M.; Ueno, M.; Oshima, Y.; Kondo, Y. Synthesis of dictyomedins using phosphazene base catalyzed diaryl ether formation. *Tetrahedron Lett.* **2007**, *48*, 8918-8921.

(32) Roman, G.; Vlahakis, J. Z.; Vukomanovic, D.; Nakatsu, K.; Szarek, W. A. Heme oxygenase inhibition by 1-aryl-2-(1*H*-imidazol-1-yl/1*H*-1,2,4-triazol-1-yl)ethanones and their derivatives. *ChemMedChem* **2010**, 5, 1541-1555.

(33) Petrone, D. A.; Lischka, M.; Lautens, M. Harnessing reversible oxidative addition: application of diiodinated aromatic compounds in the carboiodination process. *Angew. Chem. Int. Ed.* **2013**, 52, 10635-10638.

(34) Hamamoto, H.; Hattori, S.; Takemaru, K.; Miki, Y. Hypervalent iodine(III)-LiX combination in fluoroalcohol solvent for aromatic halogenation of electron-rich arenecarboxylic acids. *Synlett* **2011**, 1563-1566.

(35) Shen, S.-D.; Zhang, G.-P.; Lei, M.; Hu, L.-H. First total synthesis of salvianolic acid C, tournefolic acid A, and tournefolal. *Arkivoc* **2012**, 204-213.

(36) Desage-El Murr, M.; Nowaczyk, S.; Le Gall, T.; Mioskowski, C. Synthesis of pulvinic acid and norbadione A analogues by Suzuki-Miyaura cross-coupling of benzylated intermediates. *Eur. J. Org. Chem.* **2006**, 1489-1498.

(37) Hamada, A.; Yaden, E. L.; Horng, J. S.; Ruffolo, R. R., Jr.; Patil, P. N.; Miller, D. D. N-Substituted imidazolines and ethylenediamines and their action on α- and β-adrenergic receptors. *J. Med. Chem.* **1985**, *28*, 1269–1273.

(38) Shu, P.; Xiao, X.; Zhao, Y.; Xu, Y.; Yao, W.; Tao, J.; Wang, H.; Yao, G.; Lu, Z.; Zeng, J.; Wan, Q. Interrupted Pummerer reaction in latent-active glycosylation: glycosyl donors with a recyclable and regenerative leaving group. *Angew. Chem. Int. Ed.* **2015**, *54*, 14432-14436.

(39) Cardinal, S.; Paquet-Côté, P.-A.; Azelmat, J.; Bouchard, C.; Grenier, D.; Voyer, N. Synthesis and anti-inflammatory activity of isoquebecol. *Biorg. Med. Chem.* **2017**, *25*, 2043–2056.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53