# Communications



X. Wang,\* G. D. Cuny,\* T. Noël\* \_\_\_\_\_

A Mild, One-Pot Stadler–Ziegler Synthesis of Arylsulfides Facilitated by Photoredox Catalysis in Batch and Continuous-Flow **Visible advance**: A mild, one-pot Stadler– Ziegler process for C–S bond formation has been developed. The method employs the photoredox catalyst [Ru-(bpy)<sub>3</sub>Cl<sub>2</sub>]·6 H<sub>2</sub>O irradiated with visible light. A variety of aryl–alkyl and diaryl

R-SH + Ar-NH<sub>2</sub>

sulfides were prepared from readily available arylamines and aryl/alkylthiols in good yields. The use of a photo microreactor led to a significant improvement with respect to safety and efficiency.

► R-S-Ar

photoredox catalyst, visible light

*t*BuONO, acid (cat.), MeCN, RT – Minimum formation of diazosulfide

- In continuous-flow: 15 s to completion

#### Photoredox Catalysis

# A Mild, One-Pot Stadler–Ziegler Synthesis of Arylsulfides Facilitated by Photoredox Catalysis in Batch and Continuous-Flow\*\*

Xiao Wang,\* Gregory D. Cuny,\* and Timothy Noël\*

Many aromatic molecules containing an aryl carbon–sulfur bond display properties that afford them utility in a broad array of fields, including pharmacology.<sup>[1-3]</sup> Transition metalcatalyzed cross-coupling reactions are now among the most important reaction classes for constructing aromatic C–N<sup>[4]</sup> and C–O<sup>[5]</sup> bonds. However, the discovery of efficient aromatic C–S bond forming processes remains a challenge.<sup>[6]</sup> Several Pd-,<sup>[7a-c]</sup> Cu-,<sup>[7d,e]</sup> and Ni-catalyzed<sup>[7f]</sup> coupling reactions of aryl halides (or boronic acids) and thiols have been reported for generating arylsulfides (Scheme 1). Nevertheless, high reaction temperatures, the necessity of strong base, as well as limited substrate scope and functional group compatibility remain impediments of these approaches. In addition, many of these methods also suffer from catalyst poisoning by the sulfur-containing substrates and products.

A more traditional process to construct C–S bonds is the Stadler–Ziegler reaction,<sup>[8–10]</sup> in which arylamines are converted to the corresponding diazonium salts and then allowed to react with thiolates to yield arylsulfides. The broad scope and efficiency of this reaction makes it attractive. In addition, arylamines are usually more readily available and less expensive than the corresponding arylhalides. Since the pioneering work by Stadler<sup>[8]</sup> and later by Ziegler,<sup>[9]</sup> this process has been applied extensively in industry for manufacturing arylsulfides.<sup>[11]</sup> Various modifications have also been made to achieve improved yields and milder reaction conditions.<sup>[12]</sup> However, these methods require multiple steps including both preparation and isolation of the diazo-

```
[*] Dr. X. Wang, Prof. Dr. G. D. Cuny
    Harvard NeuroDiscovery Center, Harvard Medical School and
    Brigham & Women's Hospital
    65 Landsdowne Street, Cambridge, MA 02139 (USA)
    E-mail: xwang21@partners.org
    Prof. Dr. G. D. Cuny
    Department of Pharmacological and Pharmaceutical Sciences,
    College of Pharmacy, University of Houston
    549 A Science and Research Bldg 2, Houston, TX 77204 (USA)
    E-mail: gdcuny@central.uh.edu
    Dr. T. Noël
    Department of Chemical Engineering and Chemistry, Micro Flow
    Chemistry & Process Technology, Eindhoven University of Tech-
    nology, Den Dolech 2, 5612 AZ Eindhoven (The Netherlands)
    E-mail: t.noel@tue.nl
[**] X.W. and G.D.C. thank the Harvard NeuroDiscovery Center for
    financial support. T.N. would like to acknowledge financial support
    from the Dutch Science Foundation for a VENI grant. The kind
    donation of [Ru(bpy)<sub>3</sub>Cl<sub>2</sub>]·6 H<sub>2</sub>O by Furuya Metal Co., Ltd. is
    appreciated by T.N. We are grateful to Prof. Dennis P. Curran for his
    valuable suggestions.
```

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201303483.

Transition-metal-catalyzed cross-coupling reactions to form C-S bond:



Traditional Stadler-Ziegler reaction to form C-S bond:



 $\ensuremath{\textit{Scheme 1.}}$  Comparison of various methods for constructing aromatic C–S bonds.

nium salts and preparation of the sodium thiolates prior to the coupling step. More importantly, both the diazonium salts and the subsequent intermediate diazosulfides are potentially explosive, especially when heated.<sup>[13]</sup> Herein we report a single-step, one-pot Stadler–Ziegler process facilitated by photoredox catalysis for preparing a variety of aryl-alkyl and diaryl sulfides at room temperature with minimum formation of diazosulfides.

To avoid the isolation of diazonium salts and to enhance overall reaction efficiency, we envisioned combining the diazotization and coupling steps in a one-pot protocol (Scheme 1). Alkyl nitrites, which require only a catalytic amount of organic acid to generate the reactive [NO]<sup>+</sup> intermediate, were considered to replace sodium nitrite for diazonium formation.<sup>[12d,14]</sup> For the sulfurous coupling fragment, thiols would be used directly instead of thiolates.

We commenced our investigations by exploring the direct Stadler–Ziegler reaction between 4-methoxythiophenol (1) and aniline (2). Preliminary results revealed that in the presence of amyl nitrite and a catalytic amount of *p*toluenesulfonic acid monohydrate, 2 could be converted to the phenyldiazonium salt in situ. This compound readily reacted with 1 in MeCN at room temperature either in the presence or absence of light to afford disulfide 4 and diazosulfide 5. Interestingly, only trace amounts of the desired diarylsulfide 3 were detected (Table 1, entries 1 and 2). However, in the presence of the photoredox catalyst [Ru-(bpy)<sub>3</sub>Cl<sub>2</sub>]·6H<sub>2</sub>O and visible light,<sup>[15,16]</sup> 3 was the predominant

www.angewandte.org

© 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

• These are not the final page numbers!

**Table 1:** The effects of light, catalyst and nitrite source on product distribution in the photocatalytic Stadler-Ziegler reaction.<sup>[17]</sup>



Entry <sup>[a]</sup>	Light <sup>[b]</sup>	SET <sup>[c]</sup> catalyst	Loading (mol%)	Nitrite	<b>3</b> [%]	<b>4</b> [%]	5 [%]
1	$\checkmark$	-	_	amylONO	1	26	68
2	×	-	-	amylONO	2	28	62
3	$\checkmark$	[Ru(bpy)₃Cl₂] ·6 H₂O	1.0	amylONO	59	30	9
4	×	[Ru(bpy)₃Cl₂] ·6 H₂O	1.0	amylONO	3	30	68
5	$\checkmark$	[Ru(bpy)₃Cl₂] ·6 H₂O	1.0	<i>t</i> BuONO	86	12	1
6	$\checkmark$	$[Ru(phen)_3Cl_2]$ ·6 H <sub>2</sub> O	1.0	<i>t</i> BuONO	77	17	2
7	$\checkmark$	Cu <sub>2</sub> O	0.5	<i>t</i> BuONO	62	35	<1

[a] LC yields were recorded, with Ph<sub>2</sub>O as an internal standard. [b] 20 W fluorescent light bulb was used. [c] SET = single electron transfer, bpy = 2,2'-bipyridine, phen = phenanthroline.

product. For example, when the phenyldiazonium salt was generated in situ in the presence of a household 20 W fluorescent light source and 1 mol%  $[Ru(bpy)_3Cl_2] \cdot 6H_2O$ , **3** was produced in 59% yield after only 1 h (Table 1, entry 3). Furthermore, when this reaction was conducted in the absence of light irradiation, **3** was only produced in negligible amounts (Table 1, entry 4). Further optimization showed that *tert*-butylnitrite (*t*BuONO) was a better nitrite source, leading to an increased yield of **3** to 85% (entry 5). [Ru-(phen)\_3Cl\_2] \cdot 6H\_2O also proved effective, affording **3** in 77% yield (entry 6). However, copper(I) oxide, a commonly used single electron transfer (SET) catalyst in the Sandmeyer and related reactions, gave a lower yield of **3** (entry 7).

Once the optimal conditions were established, the substrate scope of this transformation was explored (Scheme 2). Reactions of coupling components with *meta-* and *para*substituents provided good to excellent yields of the arylsulfides. In these cases, the electronic property of the substituents had little impact on the product yields. *Ortho*-substituted reactants generally led to slightly diminished yields (**3d**, **g**, **h**), depending on the size of the substituent. Alkylthiols exhibited similar reactivities as compared to arylthiols (**3m**, **n**). Nheterocyclic reactants resulted in somewhat lower product yields (**3j**, **k**, **l**). Notably, the compatibility of aryl halides in this transformation makes it a suitable complement to the existing Ullmann-type cross-coupling reactions (**3e**, **f**, **g**, **h**). In the case where both OH and SH groups were presented, the S-thiolation product (**3b**) predominated.

To further address the safety concerns associated with the diazonium intermediates, a continuous-flow protocol was developed.<sup>[18]</sup> Due to its small dimensions, microreactor



**Scheme 2.** Substrate scope of the photocatalytic Stadler–Ziegler process.

technology provides reduced safety hazards and high surface-to-volume ratios<sup>[19]</sup> The latter is particularly advantageous for photochemical syntheses since it allows for a homogeneous irradiation of the reaction medium.<sup>[20,21]</sup> An operationally simple microfluidic setup was assembled as shown in Figure 1 and the reagents were introduced through a single syringe pump into a visible light transparent capillary microreactor (464  $\mu$ L, PFA tubing, 500  $\mu$ m inner diameter), which was irradiated with blue LED's. By separating *tert*-butylnitrite from other reagents, the diazonium intermediates are solely generated in the microreactor and are immediately consumed during the course of the reaction.

The photocatalytic Stadler-Ziegler reaction between 4methoxythiophenol (1) and aniline (2) was conducted in continuous flow under otherwise similar experimental conditions as described for the batch experiments. The formation of segmented gas-liquid flow served as an indication for the loss of nitrogen from the diazonium intermediates and, therefore, progress of the reaction was qualitatively monitored (Figure 1c). The short length scale in the microreactor provides increased photon flux. This results in a significant acceleration of the direct Stadler-Ziegler reaction; full conversion could be obtained within 15s residence time (Scheme 3). In total 13.2 mmol of 3a per hour can be synthesized in continuous flow, which represents a 78-fold improvement when compared to the corresponding batch experiment. As can be seen from Scheme 3, all three reactions could be completed within 15 s.

www.angewandte.org





**Figure 1.** a) Schematic representation of the microfluidic setup used for the continuous-flow synthesis of arylsulfides through a photocatalytic Stadler–Ziegler reaction. b) Picture of the capillary microreactor irradiated with blue LED light. More details about the photomicroreactor assembly can be found in the Supporting Information. c) Segmented gas–liquid flow at the outlet of the photo microreactor.



**Scheme 3.** Formation of biarylsulfides through a photocatalytic Stadler–Ziegler reaction in continuous-flow.

The traditional Stadler–Ziegler reaction is well known for its complicated set of possible mechanisms.<sup>[12]</sup> Normally, it has been proposed to proceed through a  $S_{RN}1$  chain mechanism,<sup>[11a,12b,22]</sup> or in some cases, a  $S_{RN}2$  mechanism.<sup>[12c]</sup> In those mechanisms, with loss of nitrogen, the diazosulfide can generate the sulfide product using the sulfide radical anion as a single electron transfer reagent ( $S_{RN}1$  and  $S_{RN}2$ ), and/or a disulfide by radical dimerization ( $S_{RN}2$ ). In the absence of photoredox catalyst (Table 1, entries 1 and 2), disulfide **4** was observed along with the unconsumed diazosulfide **5**, which might indicate an inefficient  $S_{RN}2$  mechanism. The significantly decreased amount of **4** and **5** observed under the photoredox conditions (entry 3) may reflect a more efficient SET route over the  $S_{RN}1/S_{RN}2$  processes. Initially, we surmised that the  $[Ru(bpy)_3Cl_2] \cdot 6H_2O$  catalyst facilitates the overall transformation in favor of the desired sulfide product, through the oxidative quenching of  $*Ru(bpy)_3^{2+}$  by either the diazonium salt (Scheme 4, pathway A),<sup>[16b,23]</sup> the diazo-sulfide (Scheme 4, pathway B), or the disulfide (Scheme 4, pathway C).<sup>[24,25]</sup>



**Scheme 4.** Possible mechanisms for the photocatalytic Stadler–Ziegler reaction.

To explore the mechanistic details, experiments in Table 1, entries 3 and 4 were performed with quantification of different species at multiple time points (Figure 2). Initially, the reaction was protected from light, during which the amounts of diazosulfide **5** and disulfide **4** accumulated and stabilized at a maximum level after 1 h, with **5** as the predominant species. After 1 h, the reaction was irradiated with light.<sup>[26]</sup> The sulfide product **3** started to form rapidly, while **5** was consumed at approximately the same rate. Monitoring the reaction revealed that the maximum yield of **3** was approached at



Figure 2. HPLC yields of sulfide 3, disulfide 4, and diazosulfide 5 over time (0–5 h).

www.angewandte.org

© 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

These are not the final page numbers!

4 h when diazosulfide **5** was undetectable.<sup>[17]</sup> These results suggest that diazosulfide **5** is a reactive, photo-labile intermediate in the catalytic cycle (Scheme 4, pathway B). The amount of disulfide **4** remained largely constant, thus it is less likely to be a major reactive intermediate. From these results, we can conclude that the reaction mainly proceeds through pathway B. Other minor and competing mechanisms, including  $S_{RN}1$  and  $S_{RN}2$ , are less pronounced under our photocatalytic conditions. Further mechanistic studies of this reaction are in progress.

In conclusion, a novel one-pot Stadler–Ziegler process to form C–S bonds has been developed that operates under mild reaction conditions. By employing the photoredox catalyst  $[Ru(bpy)_3Cl_2]\cdot 6H_2O$  irradiated with visible light, arylsulfides can be prepared from readily available arylamines and aryl/ alkylthiols in good yields. Notably, the process eliminates the need for diazonium salt isolation and minimizes formation of diazosulfides, both of which are potential explosion hazards. To further address the safety concerns associated with diazonium species, a scalable continuous-flow protocol was developed. The use of microreactors led to an improved irradiation of the reaction medium, which resulted in significant accelerations (full conversion within 15 s residence time) when compared to its batch counterpart.

#### **Experimental Section**

General batch reaction procedure:<sup>[18]</sup> In a 40 mL clear glass vial with PTFE septum (from I-Chem) punched with a disposable syringe needle (for venting N<sub>2</sub>), photoredox catalyst [Ru(bpy)<sub>3</sub>Cl<sub>2</sub>]·6H<sub>2</sub>O (0.010 mmol) was added to a mixture of thiol (1.0 mmol), amine (1.3 mmol), and TsOH·H<sub>2</sub>O (0.015 mmol) in MeCN (7.0 mL) at room temperature. The catalyst normally dissolves completely in 5 min upon stirring. Under visible light generated by a 20 W fluorescent bulb, tert-butylnitrite (2.0 mmol) was slowly added to the mixture. The reaction was monitored by TLC and HPLC, and was stopped after 5 to 16 h. For safety concern, water (0.10-0.20 mL) was added to the reaction mixture to moisturize any unconsumed diazonium salt. Silica gel (0.50-1.0 g) was next added and then the mixture was concentrated. The wet silica gel was loaded onto a flash column eluting with cyclohexane/EtOAc (100:0 to 4:1) to give the desired arylsulfide. The silica gel was also moistened with a half column volume of water before disposal.

General continuous-flow procedure: An oven-dried volumetric flask (10.0 mL) was charged with TsOH·H<sub>2</sub>O (8.8 mg, 0.046 mmol) and [Ru(bpy)<sub>3</sub>Cl<sub>2</sub>]·6H<sub>2</sub>O (22.4 mg, 0.030 mmol). The vessel was fitted with a septum and purged with argon. Next, thiol (3.0 mmol) and aniline (3.9 mmol) were added through a syringe and acetonitrile was added to make the solution volume 10 mL. A second oven-dried volumetric flask (10.0 mL) was fitted with a septum and purged with argon. tert-Butylnitrite (720 µL, 6 mmol) was added through a syringe and acetonitrile was added to make the solution volume 10 mL. These two solutions were loaded in 10 mL BD Discardit II plastic syringes and fitted to a single syringe pump. The different solutions were introduced into the photo microreactor device, as described in Figure 1a, with the appropriate flow rates to provide the different residence times. When exiting the photo microreactor, the reaction was collected in a vial, which was covered in aluminum foil. Typically, each experiment is preceded by a flush of four reactor volumes in order to ensure steady-state data collection. Next, a sample was collected in order to obtain exactly 1.5 mmol of product. The organic phase was concentrated in vacuo and purified by column chromatography eluting with petroleum ether or hexanes and  $0\!\!-\!\!10\,\%$  ethyl acetate.

Received: April 24, 2013 Published online:

Keywords: C–S bond formation  $\cdot$  microflow chemistry  $\cdot$  one-pot processes  $\cdot$  visible-light photoredox catalysis

- G. Liu, J. T. Link, Z. Pei, E. B. Reilly, S. Leitza, B. Nguyen, K. C. Marsh, G. F. Okasinski, T. W. von Geldern, M. Ormes, K. Fowler, M. Gallatin, *J. Med. Chem.* 2000, 43, 4025.
- [2] Y. Wang, S. Chackalamannil, Z. Hu, J. W. Clader, W. Greenlee, W. Billard, H. Binch, G. Crosby, V. Ruperto, R. A. Duffy, R. McQuade, J. E. Lachowicz, *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2247.
- [3] L. Llauger, H.-Z. He, J.-N. Kim, J. Aguirre, N. Rosen, U. Peters, P. Davies, G. Chiosis, J. Med. Chem. 2005, 48, 2892.
- [4] a) D. S. Surry, S. L. Buchwald, Angew. Chem. 2008, 120, 6438;
   Angew. Chem. Int. Ed. 2008, 47, 6338; b) B. Schlummer, U. Scholz, Adv. Synth. Catal. 2004, 346, 1599.
- [5] a) P. J. Fagan, E. Hauptman, R. Shapiro, A. Casalnuovo, J. Am. Chem. Soc. 2000, 122, 5043; b) A. Aranyos, D. W. Old, A. Kiyomori, J. P. Wolfe, J. P. Sadighi, S. L. Buchwald, J. Am. Chem. Soc. 1999, 121, 4369; c) G. Mann, C. Incarvito, A. L. Rheingold, J. F. Hartwig, J. Am. Chem. Soc. 1999, 121, 3224.
- [6] a) T. Kondo, T. Mitsudo, *Chem. Rev.* 2000, 100, 3205; b) C. C. Eichman, J. P. Stambuli, *Molecules* 2011, 16, 590.
- [7] a) T. Migita, T. Shimizu, Y. Asami, J.-I. Shiobara, Y. Kato, M. Kosugi, *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1385; b) M. Murata, S. L. Buchwald, *Tetrahedron* **2004**, *60*, 7397; c) M. A. Fernández-Rodríguez, Q.-L. Shen, J. F. Hartwig, *J. Am. Chem. Soc.* **2006**, *128*, 2180; d) G. Y. Li, G. Zheng, A. F. Noonan, *J. Org. Chem.* **2001**, *66*, 8677; e) L. Wang, W.-Y. Zhou, S.-C. Chen, M.-Y. He, Q. Chen, *Synlett* **2011**, 3041; f) P. Guan, C.-S. Cao, Y. Liu, Y.-F. Li, P. He, Q. Chen, G. Liu, Y.-H. Shi, *Tetrahedron Lett.* **2012**, *53*, 5987.
- [8] O. Stadler, Ber. Dtsch. Chem. Ges. 1884, 17, 2075.
- [9] J. H. Ziegler, Ber. Dtsch. Chem. Ges. 1890, 23, 2469.
- [10] G. E. Hilbert, T. B. Johnson, J. Am. Chem. Soc. 1929, 51, 1526.
- [11] a) A. N. Abeywickrema, A. L. J. Beckwith, J. Am. Chem. Soc.
  1986, 108, 8227; b) S. Perumal, R. Chandrasekaran, V. Vijayabaskar, D. A. Wilson, Magn. Reson. Chem. 1995, 33, 779; c) G. Smith, T. Ruhland, G. Mikkelsen, K. Andersen, C. T. Christoffersen, L. H. Alifrangis, A. Mørk, S. P. Wren, N. Harris, B. M. Wyman, G. Brandt, Bioorg. Med. Chem. Lett. 2004, 14, 4027; d) G. Smith, G. Mikkelsen, J. Eskildsen, C. Bundgaard, Bioorg. Med. Chem. Lett. 2006, 16, 3981; e) Y. Saito, N. Wada, S. Kusano, T. Miyazawa, S. Takahashi, Y. Toyokawa, I. Kajiwara, Patent US 4932999, 1990.
- [12] a) G. Petrillo, M. Novi, G. Garbarino, C. Dell'Erba, *Tetrahedron Lett.* 1985, 26, 6365; b) G. Petrillo, M. Novi, G. Garbarino, C. Dell'Erba, *Tetrahedron* 1986, 42, 4007; c) M. Barbero, I. Degani, N. Diulgheroff, S. Dughera, R. Fochi, M. Migliaccio, J. Org. Chem. 2000, 65, 5600; d) M. Barbero, M. Crisma, I. Degani, R. Fochi, P. Perracino, Synthesis 1998, 1171; e) G. K. S. Prakash, D. Hoole, D. S. Ha, J. Wilkinson, G. A. Olah, Arkivoc 2002, 13, 50.
- [13] For a reported explosion, see: J. Laquidara, Chem. Eng. News 2001, 79, 6.
- [14] a) L. Friedman, J. F. Chelbowski, J. Org. Chem. 1968, 33, 1636;
   b) D. J. Adams, A. Goddard, J. H. Clark, D. J. Macquarrie, Chem. Commun. 2000, 987.
- [15] For reviews of photoredox catalysts, see: a) A. Juris, V. Balzani, F. Barigelletti, S. Campagna, P. Belser, A. von Zelewsky, *Coord. Chem. Rev.* **1988**, *84*, 85; b) J. M. R. Narayanam, C. R. J. Stephenson, *Chem. Soc. Rev.* **2011**, *40*, 102; c) J. W. Tucker, C. R. J. Stephenson, *J. Org. Chem.* **2012**, *77*, 1617; d) J. Xuan, W.-

Angew. Chem. Int. Ed. 2013, 52, 1-6

© 2013 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim

### www.angewandte.org

These are not the final page numbers!

### Angewandte Communications

J. Xiao, Angew. Chem. 2012, 124, 6934; Angew. Chem. Int. Ed. 2012, 51, 6828.

- [16] a) D. A. Nagib, D. W. C. MacMillan, *Nature* 2011, 480, 224; b) D.
  Kalyani, K. B. McMurtrey, S. R. Neufeldt, M. S. Sanford, J. Am. Chem. Soc. 2011, 133, 18566; c) E. L. Tyson, M. S. Ament, T. P.
  Yoon, J. Org. Chem. 2013, 78, 2046.
- [17] No other side products, besides compounds **4** and **5**, were observed within detectable range in these experiments.
- [18] Although the amount of diazosulfide is minimized in this transformation, appropriate safety measurements are recommended for the batch experiments.
- [19] For reviews pertaining to flow chemistry, see: a) T. Noël, S. L. Buchwald, *Chem. Soc. Rev.* 2011, 40, 5010; b) M. Oelgemöller, O. Shvydkiv, *Molecules* 2011, 16, 7522; c) R. L. Hartman, J. P. McMullen, K. F. Jensen, *Angew. Chem.* 2011, 123, 7642; *Angew. Chem. Int. Ed.* 2011, 50, 7502; d) T. Noël, V. Hessel, *ChemSusChem* 2013, 6, 405.
- [20] For selected papers concerning the combination of photochemistry and continuous microflow, see: a) J. W. Tucker, Y. Zhang, T. F. Jamison, C. R. J. Stephenson, Angew. Chem. 2012, 124, 4220; Angew. Chem. Int. Ed. 2012, 51, 4144; b) F. Lévesque, P. H. Seeberger, Angew. Chem. 2012, 124, 1738; Angew. Chem. Int. Ed. 2012, 51, 1706; c) R. S. Andrews, J. J. Becker, M. R. Gagne, Angew. Chem. 2012, 124, 4216; Angew. Chem. Int. Ed. 2012, 51, 4140.
- [21] For a review on the combination of photoredox catalysis and continuous-flow microreactors: T. Noël, X. Wang, V. Hessel, *Chim Oggi* 2013, 31(3), 10.

- [22] a) J. F. Bunnett, Acc. Chem. Res. 1978, 11, 413; b) C. Dell'Erba,
   A. Houmam, M. Novi, G. Petrillo, J. Pinson, J. Org. Chem. 1993, 58, 2670.
- [23] For the use of diazonium salts in photoredox catalysis: a) D. P. Hari, T. Hering, B. König, Org. Lett. 2012, 14, 5334; b) D. P. Hari, P. Schroll, B. König, J. Am. Chem. Soc. 2012, 134, 2958; c) T. Hering, D. P. Hari, B. König, J. Org. Chem. 2012, 77, 10347; d) P. Schroll, D. P. Hari, B. König, ChemistryOpen 2012, 1, 130; e) H. Cano-Yelo, A. Deronzier, J. Chem. Soc. Perkin Trans. 2 1984, 1093; f) H. Cano-Yelo, A. Deronzier, Tetrahedron Lett. 1984, 25, 5517; g) For a recent review about this topic: D. P. Hari, B. König, Angew. Chem. 2013, 125, 4832; Angew. Chem. Int. Ed. 2013, 52, 4734.
- [24] For mechanistic studies of the cleavage of disulfide radical anions, see: S. Antonello, K. Daasbjerg, H. Jensen, F. Taddei, F. Maran, J. Am. Chem. Soc. 2003, 125, 14905.
- [25] For the use of thiocompounds in photoredox catalysis: a) Y. Cheng, J. Yang, Y. Qu, P. Li, Org. Lett. 2012, 14, 98; b) P. Guillo, O. Hamelin, P. Batat, G. Jonusauskas, N. D. McClenaghan, S. Menage, Inorg. Chem. 2012, 51, 2222; c) J.-M. Zen, S.-L. Liou, A. S. Kumar, M.-S. Hsia, Angew. Chem. 2003, 115, 597; Angew. Chem. Int. Ed. 2003, 42, 577; d) T. Miyashita, M. Matsuda, Bull. Chem. Soc. Jpn. 1985, 58, 3031; e) T. Miyashita, M. Matsuda, Bull. Chem. Soc. Jpn. 1981, 54, 1740; f) A. Deronzier, T. J. Meyer, Inorg. Chem. 1980, 19, 2912.
- [26] For an experiment of monitoring the reaction under light from the beginning, see the Supporting Information.

These are not the final page numbers!