


Synthesis of Pyrido[2,1-*b*]quinazolin-11-ones and Dipyrido[1,2-*a*:2',3'-*d*]pyrimidin-5-ones by Pd/DIBPP-Catalyzed Dearomatizing Carbonylation

Tongyu Xu and Howard Alper*

Centre for Catalysis Research and Innovation, Department of Chemistry, University of Ottawa, 10 Marie Curie, Ottawa, Ontario Canada, K1N 6N5

S Supporting Information



ABSTRACT: N-Fused heterocycles can be easily synthesized by palladium-catalyzed dearomatizing carbonylation using 1,3-bis(diisobutylphosphino)propane (DIBPP) as the ligand. Pyrido[2,1-*b*]quinazolin-11-ones were obtained from *N*-(2-bromophenyl)pyridine-2-amines in up to quantitative yield and dipyrido[1,2-*a*:2',3'-*d*]pyrimidin-5-ones from 3-bromo-*N*-(pyridine-2-yl)pyridine-2-amines in up to 84% yield. The cyclocarbonylation can be also realized without isolation of compound **1** and additional palladium catalyst.

Nitrogen-containing heterocycles are important molecular motifs in natural products, materials and bioactive molecules.¹ The development of efficient and environmental friendly methods for the synthesis of diverse heterocycles is an important challenge in synthetic chemistry.^{2,3} Transition-metal-catalyzed reactions are potentially valuable for the synthesis of a wide range of heterocycles. Dearomatization has become an efficient method to prepare complex nitrogen-containing heterocycles.^{4,5} For example, pyridine derivatives have been used to synthesize different kinds of N-fused heterocycles via transition-metal-catalyzed coupling, cycloisomerization, oxidation, and other reactions.⁵ Carbonylation is a powerful tool to synthesize many valuable carbonyl-containing compounds from a broad scope of substrates.⁶ There are a few reports using dearomatizing carbonylation to synthesize N-fused heterocyclic compounds.⁷ Recently, Beller, Wu and co-workers reported the base-controlled synthesis of fused quinazolinones by a palladium-catalyzed carbonylation/nucleophilic aromatic substitution sequence.^{7a} 11*H*-Pyrido[2,1-*b*]quinazolin-11-one⁸ was also prepared by the palladium-catalyzed C–H carbonylation of *N*-aryl-2-aminopyridines using K₂S₂O₈ as the oxidant in TFA.^{7b} Gevorgyan and co-workers reported a method to prepare indolizines from 2-propargylpyridines by palladium-catalyzed carbonylative cyclization/arylation.^{7c} Tilley and co-workers reported a palladium-catalyzed carbonyl insertion route to pyrido[2,1-*b*]quinazoline, but with only four examples.^{7d} However, the carbonylation of 3-bromo-*N*-(pyridine-2-yl)-pyridine-2-amine to form dipyrido[1,2-*a*:2',3'-*d*]pyrimidin-5-one⁹ was not reported, where two pyridinyl moieties can retard the catalytic activities. Herein we describe the facile synthesis of

both pyrido[2,1-*b*]quinazolin-11-ones and dipyrido[1,2-*a*:2',3'-*d*]pyrimidin-5-ones by palladium-catalyzed dearomatizing carbonylation using 1,3-bis(diisobutylphosphino)propane (DIBPP) as the ligand.

Initially, substrate **1a** was used to optimize the reaction conditions (Table 1). Different ligands were studied when the reaction was carried out in DMA at 120 °C using DBU. Davephos (**L1**) gave 11*H*-pyrido[2,1-*b*]quinazolin-11-one **2a** in 18% isolated yield, and most of **1a** was recovered (entry 1). Product **2a** was obtained in 21–38% yield using A-^{tr}Phos (**L2**), RuPhos (**L3**), or JohnPhos (**L4**) (entries 2–4). When tri-*tert*-butylphosphine (**L5**) was used, the yield of **2a** increased to 41% (entry 5). Tricyclohexylphosphine (**L6**) afforded **2a** in 74% yield (entry 6). The yield of **2a** increased to 92% when THF was used as the solvent (entry 7). Acetonitrile and toluene gave **2a** in 45% and 89% yields, respectively (entries 8 and 9). Different bases were then studied. Inorganic bases potassium carbonate and potassium phosphate afforded **2a** in 89% and 97% yields, respectively, while the organic base triethylamine gave **2a** in 23% yield (entries 10–12). When the temperature was lowered to 100 °C, the yield of **2a** decreased to 33% or 50% using DBU or K₃PO₄ as the base (entries 13 and 14). When a phenyl-substituted phospho-adamantane ligand **L7** was used instead of **L6**, **2a** was isolated in only 15% yield (entry 15). When the bidentate phosphine ligand **L8** (1,3-bis(diisobutylphosphino)propane, DIBPP) was used for the reaction, **2a** was obtained in excellent yield (entry 16). Another

Received: February 12, 2015

Published: March 10, 2015

Table 1. Screening of the Reaction Conditions^a

entry	ligand	base	solvent	temp (°C)	P _{CO} (bar)	yield ^b (%)
1	L1	DBU	DMA	120	20	18
2	L2	DBU	DMA	120	20	38
3	L3	DBU	DMA	120	20	35
4	L4	DBU	DMA	120	20	21
5	L5	DBU	DMA	120	20	41
6	L6	DBU	DMA	120	20	74
7	L6	DBU	THF	120	20	92
8	L6	DBU	MeCN	120	20	45
9	L6	DBU	toluene	120	20	89
10	L6	K ₂ CO ₃	THF	120	20	89
11	L6	K ₃ PO ₄	THF	120	20	97
12	L6	Et ₃ N	THF	120	20	23
13	L6	DBU	THF	100	20	33
14	L6	K ₃ PO ₄	THF	100	20	50
15	L7	K ₃ PO ₄	THF	120	20	15
16	L8	K ₃ PO ₄	THF	120	20	99
17	L9	K ₃ PO ₄	THF	120	20	65
18	L8	K ₃ PO ₄	THF	120	10	99
19	L8	K ₃ PO ₄	THF	100	10	99
20	L8	K ₃ PO ₄	THF	70	10	10
21	L8	K ₃ PO ₄	THF	100	5	99
22 ^c	L8	K ₃ PO ₄	THF	100	5	99
23 ^d	L8	K ₃ PO ₄	THF	100	5	86

^aConditions: 0.2 mmol **1a**, 5 mol % Pd(OAc)₂, Pd/P = 1/2, 0.6 mmol of base, 2 mL of solvent, 15 h. ^bIsolated yield. ^c3 mol % of Pd(OAc)₂. ^d2 mol % of Pd(OAc)₂.

trialkylphosphine **L9** afforded **2a** in 65% yield (entry 17). When the reaction temperature was lowered to 100 °C and the CO pressure to 5 bar, **2a** was obtained in nearly quantitative yield (entries 18–21). The yield did not change when the amount of catalyst was reduced to 3 mol % (entries 22 and 23).

The scope of the reaction was studied using a variety of reactants in the presence of 3 mol % of Pd(OAc)₂/3 mol % of DIBPP, THF as the solvent, and K₃PO₄ as the base at 100 °C and 5 bar of CO (Table 2). Electron-donating or electron-withdrawing substituents on the phenyl moiety have little effect on the reaction. Substrates with electron-donating substituents on the pyridyl moiety smoothly afford **2** (entries 1–3), but electron-withdrawing substituents on the pyridyl moiety retard the reaction. For example, when **1f** with fluoride on the pyridyl moiety was reacted, there was little conversion to **2f**. The carbonylation of **1f** needed higher temperature (120 °C) to give **2f** (entry 6), indicating that electron-withdrawing substituents on the pyridyl moiety could reduce the reactivity of the nitrogen in the pyridyl ring. When pyridine was replaced by quinoline, **2g** was obtained in 90% yield (entry 7). Then the

Table 2. Synthesis of 11H-Pyrido[2,1-b]quinazolin-11-one via Pd-Catalyzed Dearomatizing Carbonylation of N-(2-Bromophenyl)pyridine-2-amine Derivatives^a














entry	substrate	product	yield (%) ^b
1	1a	2a	99
2	1b	2b	98
3	1c	2c	92
4	1d	2d	88
5	1e	2e	84
6	1f	2f	80 ^c
7	1g	2g	90
8	1h	2h	99
9	1i	-	0
10	1j	-	0
11	1k	2k	31

^aConditions: 0.5 mmol of **1**, 3 mol % of Pd(OAc)₂, 3 mol % of DIBPP, 1.5 mmol of base, 5 bar CO, 5 mL of THF, 100 °C, 15 h. ^bIsolated yield. ^c120 °C, 10 bar.

reaction was applied to **1h**, affording **2h** in 99% yield (entry 8). However, when *ortho*-substituted substrates **1i** and **1j** were used, no carbonylation product **2** was isolated, while starting materials disappeared (entries 9 and 10). The product with an *o*-chlorine or methoxy substituent may not be stable. The substrate with an *o*-methyl substituent gave the desired product **2k** in 31% yield (entry 11).

Our method was also applied to bromo-substituted heteroaromatic compounds (Table 3). At 100 °C, most **1l** was not consumed in the presence of 3 mol % of Pd(OAc)₂. 5*H*-Dipyrido[1,2-*a*:2',3'-*d'*]pyrimidin-5-one (**2l**) was obtained in 52% isolated yield at 120 °C under 10 bar of CO in the presence of 5 mol % of Pd(OAc)₂ and 5 mol % of DIBPP. Substituents can have a significant effect on the reactions.

Table 3. Synthesis of 5*H*-Dipyrido[1,2-*a*:2',3'-*d*]pyrimidin-5-one via Pd-Catalyzed Dearomatizing Carbonylation of 3-Bromo-*N*-(pyridine-2-yl)pyridine-2-amine Derivatives^a

			
entry	substrate	product	yield (%) ^b
1			52
2			61
3			80
4			84
5			40 ^c (51) ^d
6			65


^aConditions: 0.5 mmol of **1**, 5 mol % of Pd(OAc)₂, 5 mol % of DIBPP, 1.5 mmol of base, 10 bar CO, 5 mL of THF, 120 °C, 15 h. ^bIsolated yield. ^c120 °C, 72 h. ^d135 °C, 15 h.

Reactants with an electron-donating substituent gave higher yields (61–84%, **2m–o**, entries 2–4). Substrate **1o** with electron-withdrawing substituent (F, entry 5) gave **2p** in 40% yield after 72 h at 120 °C, while the yield was 51% after 15 h at 135 °C. Substrate **1q** with an isoquinoline ring in the molecule also reacted smoothly to give the product **2q** in 65% yield.

The substrates (**1**) for the research were prepared via Pd-catalyzed coupling reactions.¹⁰ The dearomatic cyclocarbonylation reaction was also attempted without the isolation of compound **1** (Scheme 1). Specifically, the reaction mixture of **1a** generated in situ from 1,2-dibromobenzene and 2-aminopyridine in toluene was directly transferred to an autoclave followed by addition of DIBPP and K₃PO₄ (eq 1), affording carbonylation product **2a** in 67% yield with 20% yield of the C–N coupling product **2aa**. When the process was applied to prepare **2l**, the yield was only 22% (eq 2). However, **2n** was obtained in 68% yield from 2-amino-4-methoxypyridine and 2,3-dibromopyridine (eq 3), indicating that the electron-donating substituent could promote the reactivity of nitrogen in the pyridinyl moiety.

In conclusion, the palladium-catalyzed dearomatizing carbonylation reaction to synthesize N-fused heterocycles proceeds efficiently, affording pyrido[2,1-*b*]quinazolin-11-ones and dipyrido[1,2-*a*:2',3'-*d*]pyrimidin-5-ones in moderate to excellent yields.

Scheme 1. Dearomatic Cyclocarbonylation Reaction without Isolation of Intermediate 1



■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and spectral data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: howard.alper@uottawa.ca.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to Cytec Canada and to the Natural Sciences and Engineering Research Council of Canada for support of this research.

■ REFERENCES

- (a) Majumdar, K. C.; Chattopadhyay, S. K. *Heterocycles in Natural Product Synthesis*; Wiley-VCH: Weinheim, 2011. (b) Katritzky, A. R. *Comprehensive Heterocyclic Chemistry III*; Elsevier: Amsterdam, NY, 2008. (c) Lynch, M. A.; Duval, O.; Sukhanova, A.; Devy, J.; MacKay, S. P.; Waigh, R. D.; Nabiev, I. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 2643. (d) Jones, G.; Abarca, B. *Adv. Heterocycl. Chem.* **2010**, *100*, 195. (e) Chittchang, M.; Batsomboon, P.; Ruchirawat, S.; Ploypradith, P. *ChemMedChem* **2009**, *4*, 457. (f) Padmavathi, V.; Radha Lakshmi, T.; Mahesh, K.; Padmaja, A. *Chem. Pharm. Bull.* **2009**, *57*, 1200. (g) Eamvijarn, A.; Gomes, N. M.; Dethoup, T.; Buaruang, J.; Manoch, L.; Silva, A.; Pedro, M.; Marini, I.; Roussis, V.; Kijjoa, A. *Tetrahedron* **2013**, *69*, 8583. (2) (a) Wolfe, J. P., Ed. *Synthesis of Heterocycles via Metal-Catalyzed Reactions that Generate One or More Carbon–Heteroatom Bonds*; Springer: Berlin, 2013. (b) Bolm, C.; Beller, M. *Transition Metals for Organic Synthesis*; Wiley-VCH: Weinheim, 2004; Vols. 1 and 2. (c) Ball, C. J.; Willis, M. C. *Eur. J. Org. Chem.* **2013**, 425. (d) Zhao, C.; Crimmin, M. R.; Toste, F. D.; Bergman, R. G. *Acc. Chem. Res.* **2014**, *47*, 517. (e) Garella, D.; Borretto, E.; Stilo, A. D.; Martina, K.; Cravotto, G.; Cintas, P. *Med. Chem. Commun.* **2013**, *4*, 1323. (f) Eftekhari-Sis, B.; Zirak, M.; Akbari, A. *Chem. Rev.* **2013**, *113*, 2958. (g) Godoi, B.; Schumacher, R. F.; Zeni, G. *Chem. Rev.* **2011**, *111*, 2937. (3) (a) Liao, J.-Y.; Shao, P.-L.; Zhao, Y. *J. Am. Chem. Soc.* **2015**, *137*, 628. (b) Michlik, S.; Kempe, R. *Nat. Chem.* **2013**, *5*, 140. (c) Sherman, E. S.; Chemler, S. R. *Adv. Synth. Catal.* **2009**, *351*, 467. (d) Hummel, J. R.; Ellman, J. A. *J. Am. Chem. Soc.* **2015**, *137*, 490. (e) Du, W.; Gu, Q.

Li, Z.; Yang, D. *J. Am. Chem. Soc.* **2015**, *137*, 1130. (f) Candito, D. A.; Lautens, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 6713. (g) Zhang, W.-Z.; Chu, J. C. K.; Oberg, K. M.; Rovis, T. *J. Am. Chem. Soc.* **2015**, *137*, 553. (h) Chi, Y.; Zhang, W. X.; Xi, Z. *Org. Lett.* **2014**, *16*, 6274.

(4) (a) Zhuo, C.-X.; Zheng, C.; You, S.-L. *Acc. Chem. Res.* **2014**, *47*, 2558. (b) Zheng, C.; Zhuo, C.-X.; You, S.-L. *J. Am. Chem. Soc.* **2014**, *136*, 16251. (c) Wang, P.-F.; Jiang, C.-H.; Wen, X.; Xu, Q.-L.; Sun, H. *J. Org. Chem.* **2015**, *80*, 1155. (d) Romano, C.; Jia, M.; Monari, M.; Manoni, E.; Bandini, M. *Angew. Chem., Int. Ed.* **2014**, *53*, 13854. (e) García-Fortanet, J.; Kessler, F.; Buchwald, S. L. *J. Am. Chem. Soc.* **2009**, *131*, 6676.

(5) (a) Hirayama, T.; Ueda, S.; Okada, T.; Tsurue, N.; Okuda, K.; Nagasawa, H. *Chem.—Eur. J.* **2014**, *20*, 4156. (b) He, Y.; Huang, J.; Liang, D.; Liu, L.; Zhu, Q. *Chem. Commun.* **2013**, *49*, 735. (c) Zhang, X.; Yang, Z.-P.; Huang, L.; You, S.-L. *Angew. Chem., Int. Ed.* **2015**, *54*, 1873. (d) Xiang, L.; Yang, Y.; Zhou, X.; Liu, X.; Li, X.; Kang, X.; Yan, R.; Huang, G. *J. Org. Chem.* **2014**, *79*, 10641. (e) Heller, S. T.; Kihō, T.; Narayan, A. R. H.; Sarpong, R. *Angew. Chem., Int. Ed.* **2013**, *52*, 11129. (f) Lee, D. J.; Han, H. S.; Shin, J.; Yoo, E. J. *J. Am. Chem. Soc.* **2014**, *136*, 11606. (g) Chernyak, D.; Skontos, C.; Gevorgyan, V. *Org. Lett.* **2010**, *12*, 3242. (h) Gao, M.; Tian, J.; Lei, A. *Chem.—Asian J.* **2014**, *9*, 2068. (i) Xia, Y.; Dudnik, A. S.; Li, Y.; Gevorgyan, V. *Org. Lett.* **2010**, *12*, 5538. (j) Chernyak, N.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2010**, *49*, 2743. (k) Bunnelle, E. M.; Smith, C. R.; Lee, S. K.; Singaram, W. S.; Rhodes, A. J.; Sarpong, R. *Tetrahedron* **2008**, *64*, 7008. (l) Wang, H.; Wang, Y.; Peng, C.; Zhang, J.; Zhu, Q. *J. Am. Chem. Soc.* **2010**, *132*, 13217. (m) Hostyn, S.; Van Baelen, G.; Lemièrre, G. L. F.; Maes, B. U. W. *Adv. Synth. Catal.* **2008**, *350*, 2653.

(6) (a) Kollár, L. *Modern Carbonylation Methods*; Wiley-VCH Verlag: Weinheim, 2008. (b) Beller, M. *Catalytic Carbonylation Reactions*; Springer: Berlin, 2006. (c) Gadge, S. T.; Bhanage, B. M. *RSC Adv.* **2014**, *4*, 10367. (d) Sumino, S.; Fusano, A.; Fukuyama, T.; Ryu, I. *Acc. Chem. Res.* **2014**, *47*, 1563. (e) Wu, X.-F.; Neumann, H.; Beller, M. *ChemCatChem* **2013**, *6*, 229. (f) Wu, X.-F.; Neumann, H.; Beller, M. *Chem. Rev.* **2013**, *113*, 1. (g) Wu, X.-F.; Neumann, H.; Beller, M. *ChemCatChem* **2012**, *4*, 447. (h) Liu, Q.; Zhang, H.; Lei, A. *Angew. Chem., Int. Ed.* **2011**, *50*, 10788.

(7) (a) Chen, J.; Natta, K.; Spannenberg, A.; Neumann, H.; Langer, P.; Beller, M.; Wu, X.-F. *Angew. Chem., Int. Ed.* **2014**, *53*, 7579. (b) Liang, D.; He, Y.; Zhu, Q. *Org. Lett.* **2014**, *16*, 2748. (c) Li, Z.; Chernyak, D.; Gevorgyan, V. *Org. Lett.* **2012**, *14*, 6056. (d) Tilley, J. W.; Coffen, D. L.; Schaer, B. H.; Lind, J. J. *J. Org. Chem.* **1987**, *52*, 2469.

(8) (a) Maity, A.; Mondal, S.; Paira, R.; Hazra, A.; Naskar, S.; Sahu, K. B.; Saha, P.; Banerjee, S.; Mondal, N. B. *Tetrahedron Lett.* **2011**, *52*, 3033. (b) Pellón, R. F.; Martínez, A.; Docampo, M. L.; Mesa, M. *Synth. Commun.* **2013**, *43*, 537.

(9) (a) Bentabed-Ababsa, G.; Ely, S. C. S.; Hesse, S.; Nassar, E.; Chevallier, F.; Nguyen, T. T.; Derdour, A.; Mongin, F. *J. Org. Chem.* **2010**, *75*, 839. (b) Sun, J.; Tan, Q.; Yang, W.; Liu, B.; Xu, B. *Adv. Synth. Catal.* **2014**, *356*, 388.

(10) (a) Aluri, B. R.; Kindermann, M. K.; Jones, P. G.; Dix, I.; Heinicke, J. *Inorg. Chem.* **2008**, *47*, 6900. (b) Loones, K. T. J.; Maes, B. U. W.; Meyers, C.; Deruytter, J. *J. Org. Chem.* **2006**, *71*, 260.