Tetrahedron 69 (2013) 2701-2713

Contents lists available at SciVerse ScienceDirect

Tetrahedron

journal homepage: www.elsevier.com/locate/tet

Efficient microwave-assisted one-pot three-component synthesis of 2,3-disubstituted benzofurans under Sonogashira conditions

Nataliya A. Markina^a, Yu Chen^{a,b}, Richard C. Larock^{a,*}

^a Department of Chemistry, Iowa State University, Ames, IA 50011, USA ^b Department of Chemistry and Biochemistry, Queens College of CUNY, Flushing, NY 11367, USA

ARTICLE INFO

Article history: Received 11 December 2012 Received in revised form 30 January 2013 Accepted 1 February 2013 Available online 8 February 2013

Keywords: Benzofuran Sonogashira coupling Three-component reaction Microwave

ABSTRACT

An efficient one-pot method for the synthesis of 2,3-disubstituted benzo[*b*]furans from commercially available 2-iodophenols, terminal acetylenes and aryl iodides has been developed utilizing Sonogashira reaction conditions. After an initial Sonogashira coupling of the 2-iodophenol with the terminal alkyne, cyclization involving the aryl iodide provides the 2,3-disubstituted benzo[*b*]furan in good to excellent yields. The use of microwave irradiation shortens the reaction times and minimizes the side products. This methodology is especially useful for the construction of libraries of highly substituted benzo[*b*]furans and their analogues.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Benzolblfurans have been studied extensively due to the high potential biological and pharmaceutical activity of this ring system.¹ Thus, numerous synthetic methods to access this important scaffold have been developed.² Among others, Pd-catalyzed reactions have proven to be highly efficient and convenient for the synthesis and functionalization of benzo[b]furans. Several review articles and books have been published that summarize previous and recent developments in this area.³ In 1996, Cacchi and coworkers reported that 2-(1-alkynyl)phenols in the presence of vinylic triflates and a palladium catalyst undergo cyclization to the corresponding 2,3-disubstituted benzo[b]furans.⁴ The authors proposed that this process most likely proceeds through a vinylic palladium intermediate, generated in situ from the vinylic triflate via oxidative addition to Pd(0). To access the starting 2-(1-alkynyl) phenols, a three-step route has been employed in most cases. First, the OH group is protected with an appropriate protecting group (e.g., acetyl), then the Sonogashira coupling is conducted, and, following deprotection of the phenol, the desired 2-(1-alkynyl) phenol is obtained, generally in moderate yields. The major problem with the direct Sonogashira reaction between a 2-iodophenol and a terminal alkyne is that the coupling is often inefficient. In addition, if basic reaction conditions are employed, 3H-benzofurans are often formed as products, instead of the desired 2-(1-alkynyl) phenol.

In order to transform Cacchi's process into a three-component process, Flynn and co-workers employed MeMgCl as an additive to mask the phenol oxygen of the iodophenol (1). They were then able to conduct an efficient Sonogashira coupling and subsequent cyclization to a benzofuran (3) without isolating the corresponding 2-(1-alkynyl) phenol (2) (Scheme 1).⁵ A number of highly substituted benzofurans **3** have been obtained in moderate to good yields using this approach. The authors applied their methodology to the synthesis of (\pm) -frondosin B and its analogues, and benzo[*b*]furan-containing inhibitors of tubulin polymerization.⁶ However, this method has not been found suitable for the construction of libraries of 2,3-substituted benzofurans, due to the highly reactive nature of the MeMgCl reagent and its incompatibility with a large number of functional groups.

Thus in 2004, Hu, Fathi, and Yang, in order to access a 210 member library of 2,3-disubstituted benzofurans, optimized the method developed by Cacchi starting from 2-(1-alkynyl)phenols, and, after their optimization studies, found more efficient conditions for the formation of 2,3-diarylsubstituted benzo[*b*]furans from 2-(1-alkynyl)phenols and aryl iodides.⁷ However, considering the three-step route required for synthesis of the 2-(1-alkynyl) phenols, the average yields of the final benzofurans (over 4 steps) were only moderate.

During the course of our own investigations into methodology for the synthesis of 2,3-disubstituted indoles under Sonogashira conditions,⁸ we have found that 2-iodophenols can also participate in an analogous process, providing an efficient and convenient new





Tetrahedror

 $[\]ast$ Corresponding author. Tel.: +1 515 294 4660; fax: +1 515 294 0105; e-mail address: larock@iastate.edu (R.C. Larock).

^{0040-4020/\$ –} see front matter @ 2013 Elsevier Ltd. All rights reserved. http://dx.doi.org/10.1016/j.tet.2013.02.003



Scheme 1. A one-pot approach to benzofurans using MeMgCl.

route to 2,3-disubstituted benzofurans. This finding encouraged us to proceed with our own optimization studies of this process.

2. Results and discussion

2.1. Optimization

2-lodophenol (**4**), phenyl acetylene (**5**), and ethyl 4-iodobenzoate (**6**) have been employed as starting materials under the reaction

Table 1

Optimization of the reaction conditions^a

conditions we developed for the synthesis of indoles,⁸ providing the benzofuran **7** in a 51% yield (Table 1, entry 1). In order to improve the yield, optimization of the reaction conditions has been carried out.

Our initial examination of a number of palladium catalysts (Table 1, entries 1–4) indicated that bis(triphenylphosphine)palladium dichloride provided the best results. Changing the base from triethylamine to diisopropylamine (entries 5 vs 6) did not improve the yield of the desired benzofuran **7**. A study of the effect of various solvents on the second cyclization step revealed that DMF afforded



Entry	Time (min)	Temp (°C) Step 1	Solvent		Ratio 4 : 5:6	Catalyst	Yield 7 ^b (%)
	Step 1		Step 1	Step 2			
1	15	60	Et ₃ N	MeCN	1:1.05:1.1	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % Cul	51
2	15	60	Et₃N	MeCN	1:1.05:1.1	2 mol % Pd(PPh ₃) ₄ 1 mol % CuI	23
3	15	60	Et₃N	MeCN	1:1.05:1.1	2 mol % Pd(dppe) ₂ 1 mol % Cul	10
4	15	60	Et ₃ N	MeCN	1:1.05:1.1	2 mol % Pd(OAc) ₂ 4 mol % PPh ₃ 1 mol % CuI	6
5	15	60	Et ₃ N	MeCN	1:1.05:1.0	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % CuI	53
6	15	60	ⁱ Pr ₂ NH	MeCN	1:1.05:1.0	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % CuI	50
7	15	60	Et₃N	DMF	1:1.05:1.0	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % CuI	34
8	15	60	Et ₃ N	THF	1:1.05:1.0	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % CuI	29
9	15	60	Et ₃ N	Toluene	1:1.05:1.0	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % CuI	15
10	25	80	Et ₃ N	MeCN	1:1.05:1.0	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % CuI	20
11 ^c	15	25	Et ₃ N	MeCN	1:1.05:1.0	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % CuI	53
12	30	25	Et ₃ N	MeCN	1:1.05:1.0	2 mol % PdCl ₂ (PPh ₃) ₂ 1 mol % CuI	73
13	30	25	Et ₃ N	MeCN	1:1.05:1.0	3 mol % PdCl ₂ (PPh ₃) ₂ 2 mol % CuI	86
14 ^d	30	25	NMM/Et ₃ N	MeCN	1:1.2:1.0	3 mol % PdCl ₂ (PPh ₃) ₂ 2 mol % Cul	89
15 ^e	30	25	THF/Et ₃ N	MeCN	1:1.2:1.0	3 mol % $PdCl_2(PPh_3)_2$ 2 mol % Cul	96

^a Unless otherwise noted, all of the reactions were carried out under microwave irradiation on a 1.0 mmol scale in microwave-resistant vials.

^b Isolated yields after column chromatography.

^c When the first step of the reaction was carried out at 25 °C, much cleaner reaction mixtures were obtained than at 60 °C.

^d 0.5 mL of *N*-methylmorpholine (NMM)/1.5 mL of Et₃N using anhydrous solvents under argon.

^e 0.5 mL of THF/1.0 mL of Et₃N, and Cul were added as a solution in 0.5 mL of Et₃N using anhydrous solvents under argon.

product **7** in a 34% yield (entry 7), whereas THF and toluene were less efficient, providing only 29% and 15% yields, respectively (entries 8 and 9). Thus, acetonitrile appeared to be the best solvent for this transformation. An increase in the temperature of the first step of the reaction to 80 °C decreased the yield of the desired product to 20% (entry 10), whereas conducting the first step at room temperature did not affect the yield of the product and the reaction was found to be cleaner based on TLC analysis when compared to the same reaction at 60 °C. Increasing the reaction time of the first step at room temperature to 30 min improved the yield of benzofuran **7** to 73% (entry 12). Slightly increasing the catalyst loading and finding the best Pd/Cu catalyst ratio (3 mol % and 2 mol %, respectively) improved the yield of benzofuran **7** to 86% (entry 13).

After the initial evaluation of the scope of this process had been performed, the solubility of many iodophenols in triethylamine was found to be insufficient. The first step of the process was inefficient, leading to the exclusive formation of coupling products of the aryl iodides with the terminal alkyne and affording only low yields of the cyclized benzofurans. Thus, additional evaluation of the solvents for the first step was performed.

In this methodology, the choice of solvents plays a crucial role. For the first step, the ideal solvent needs to be suitable for an efficient Sonogashira reaction, but should not promote cyclization of the intermediate 2-(1-alkynyl)phenol, which will result in formation of an undesired 3*H*-benzofuran. The second step, on the other hand, requires a solvent, which will promote the cyclization. Based on our observations on the effect of different solvents on the

Table 2

Scope of the reaction

cyclization step (Table 1; entries 5, 7–9), it can be concluded that a more polar solvent generally affords a more efficient cyclization process. The increased polarity of the solvent most likely produces less ion pairing between the phenolate and the Et₃NH⁺ cation, which facilitates a more favorable attack of the phenolate on the alkyne triple bond. We, therefore, tried to add additional reagents to the triethylamine to improve the solubility of the iodophenols. but still retain reaction conditions favorable for the Sonogashira coupling. Solvents, like acetonitrile and DMF, solubilized the iodophenols well, but were not suitable for the first step, since they promote the cyclization subsequent. The best additives were found to be *N*-methylmorpholine (entry 14) and THF. Eventually, the ratio of 3/1 triethylamine/THF afforded the best results and was chosen as our optimized conditions. After we discovered that the reaction was sensitive to the palladium/copper ratio, we again looked at this factor. We subsequently found that only very low amounts of CuI were needed and it was better to add the exact amount of CuI desired as a 7.5 M solution in Et₃N. When running the reaction under an inert atmosphere and anhydrous conditions, this allowed us to increase the yield of the desired benzofuran 7 to 96% (entry 15).

2.2. Evaluation of the scope

After 'optimal' conditions for the formation of the benzo[*b*]furan **7** were found, an evaluation of the reaction scope was performed (Table 2).



Table 2 (continued)



27

Table 2 (continued)

Entry	Iodophenol	Acetylene	Aryl iodide	Product	Yield ^a (%)
11	10	28	13	MeO_2C	52
12	4	S 30	6	CO ₂ Et	100
13	4	Me- _N N=- 32	6	CO ₂ Et	63 ^e
14	4	5	PhI	$ \begin{array}{c} $	87
15	4	5	OMe I 35	OMe OPh 36	53 ^r
16	4	5	OMe I 37	OMe OPh 38	84
17	4	5	CN 1 39	CN CN Ph 40	98
18	4	5	NO ₂ I 41		75 ntinued on next page)

Table 2 (continued)

Entry	Iodophenol	Acetylene	Aryl iodide	Product	Yield ^a (%)
19	4	5	43	O_2N Ph O_2N Ph O	74
20	4	5		$ \begin{array}{c} & \\ & \\ & \\ & \\ & \\ & \\ & \\ & \\ & \\ & $	96
21	4	30	47	48 0 0 0 5 5 5	73
22	4	5	NTs I 49	NTs Ph 50	58
23	4	5	0 1 51	$rac{1}{52}$	43
24	8	Me 53	OHC I	HACK CONCEPTION OF CONCEPTION	65
25	OMe OH Br 56	OAc OMe 57	AcO OAc 58	AcO OAC OMe 0Me 59	60

Table 2 (continued)



^a Isolated yields after column chromatography.

^b This compound was prepared on a large scale and recrystallized, what might have contributed to the lower yield.

^c 1.0 equiv of alkyne was employed.

^d The reaction afforded a complex mixture; for an alternative route to **27**, see Scheme 2.

^e 1.0 equiv of alkyne was employed and the first step of the process was run at 60 °C.

^f The second step of the process was conducted at 80 °C with the addition of 10 mol % Pd(PPh₃)₄.

First, several different iodophenols were investigated under our optimized reaction conditions (Table 2, entries 1-5). The electronrich methoxy-containing phenol 8, and the electron-poor estercontaining phenol **10** afforded the expected benzo[*b*]furans **9** and 11 in 91% and 92% yields, respectively (entries 1 and 2). When 6allyl-2-iodophenol (12) was employed, benzofuran 14 was formed, albeit in only a 53% yield (entry 3). The presence of a bromine atom para to the hydroxy group was well tolerated and the bromo-containing benzo[b]furans 16 and 17 were isolated in 84% and 60% yields, respectively (entries 4 and 5). Even though the addition of THF greatly improved the solubility of many of the starting o-iodophenols, some substrates (e.g., 5-iodovanillin, 7iodo-8-hydroxyquinoline-5-sulfonic acid, and 5-iodouracil) still exhibited insufficient solubility in the Et₃N/THF mixture to afford good results. The poor solubility prevented the first coupling step from proceeding in acceptable yield and thus sharply reducing the yield of the three-component coupling product while increasing the number of side products.

Various additional terminal alkynes have been studied (entries 6–12). Alkynes bearing electron-donating groups, such as alkynes **18**, **20**, and **22**, were well tolerated, providing benzofurans **19**, **21**, and **23** in 94%, 93%, and 83% yields, respectively (entries 6–8). The alkyne **24**, containing an electron-withdrawing aldehyde group in

the position ortho to the alkyne functionality was also tolerated, providing the benzofuran 25 in a 69% yield (entry 9). However, when a stronger electron-withdrawing cyano group (26) was present in the alkyne, no cyclization product was observed (entry 10). Instead a complex reaction mixture, containing the 3H-benzofuran 66, the coupling product of 26 with 6, as well as trace amounts of other products was obtained. This result can be rationalized by examining the nucleophilicity of the alkyne moiety. When an electron-withdrawing group is present, the electron density on the carbon-carbon triple bond is decreased, thus promoting cyclization of the OH group, without interception by the desired intermediate arylpalladium iodide. In the case of 4ethynylbenzonitrile (26), using the corresponding TMS-protected phenol 65 and adding TBAF during the second step of the sequence, we were able to obtain the desired benzo[b]furan 27 in a 53% yield (Scheme 2).

The use of 1-ethynylcyclohexene (**28**) afforded the desired benzo[*b*]furan **29** in a 52% yield (entry 11). The heterocyclecontaining terminal alkynes **30** and **32** were also well tolerated under our reaction conditions, providing thiophenyl- and methylimidazolyl-substituted benzofurans **31** and **33** in 100% and 63% yields, respectively (entries 12 and 13). Unfortunately, aliphatic acetylenes [e.g., 1-pentyne] led to formation of the desired benzo[*b*]



Scheme 2. An alternative route employing 4-ethynylbenzonitrile.

furan in only trace amounts and afforded a complex reaction mixture. Various attempts to modify the reaction conditions failed to improve this reaction.

Finally, we examined the scope of the aryl iodides that can be used in this process under our optimized reaction conditions. Starting with iodobenzene, 2,3-diphenylbenzo[*b*]furan (**34**) was obtained in an 87% yield (entry 14). When 4-iodoanisole (**35**) was employed, only a 22% yield of benzofuran **36** was obtained under

reported.¹⁰ We were interested in knowing if our methodology could provide such a Heck-type transformation to afford 3-(1-alkenyl)benzofurans in a one-pot fashion from 2-iodophenol (**4**). Gratifyingly, by employing butyl acrylate and slightly modifying our reaction conditions for the second step of the process, we were able to obtain the olefinic product **67** in a 56% yield (Scheme 3). The alkyne 1-ethynylcyclohexene (**28**) also afforded the desired product **68** in a 70% yield.



our optimized conditions. However, using a slightly lower temperature (80 °C) and an additional loading of the Pd catalyst in the second step improved the yield of the product **36** to 53% (entry 15). Surprisingly, the presence of an electron-donating methoxy group in the meta-position of the aryl iodide 37 was well tolerated and afforded the desired heterocycle 38 in an 84% yield (entry 16). Aryl iodides with electron-withdrawing groups, such as 4iodobenzonitrile (39) and 4-iodonitrobenzene (41), also afforded the corresponding benzofurans 40 and 42 in 98% and 75% yields, respectively (entries 17 and 18). Placing the nitro group in the position *ortho* to the iodine did not affect the efficiency of the process, providing benzo[b]furan 44 in a 74% yield (entry 19). 4-Chloroiodobenzene (45) afforded the benzo[b]furan 46 in an excellent 96% yield (entry 20). The product 48 has been obtained in a 73% yield by employing 3-ethynylthiophene (30) and p-iodoacetophenone (47) (entry 21). Various heterocyclic aryl and vinylic iodides have been examined under our standard reaction conditions, providing the corresponding heterocyclic products in moderate to good yields (entries 22-24). N-Tosyl 3-iodoindole (49) afforded the desired benzofuran 50 in a 58% yield (entry 22). When employing 2iodochromene (51), the bis-heterocyclic product 52 was obtained, albeit in only a 43% yield (entry 23). To our delight, this methodology proved to be useful for the synthesis of the highly substituted benzofurans 55, 59, and 61 (entries 24-26). 2-Fluoro-3-formyl-4iodopyridine (54) coupled with 6-methoxy-2-iodophenol (8) and 3-tolyl acetylene (53) to afford the highly substituted benzofuran 55 in a 65% yield (entry 24). Bromo-substituted benzofuran 59 was obtained in a 60% yield from the readily accessible starting materials 56, 57, and 58 (entry 25). Compound 61 was prepared in a 63% yield starting from 2-iodophenol 60 and was further converted into the corresponding aldehyde 62 in 94% yield (entry 26). Benzofurans 59 and **62** would appear to be an ideal intermediates for the synthesis of naturally-occurring Gnetuhainin F, a benzofuran from the class of oligostilbenes isolated from Gnetum hainanense species, a traditional Chinese medicine herb.⁹ Finally, the vinylic halide 2-iodo-4,4dimethylcyclohex-2-enone (63) was allowed to react with o-iodophenol and phenyl acetylene to generate the corresponding heterocycle 64, albeit in only a 34% yield (entry 27).

2.3. Additional processes and further diversification

Recently, several processes describing Heck-type Pd(0) or Rh(I)catalyzed reactions of 2-(1-alkynyl)phenols with alkenes have been We have also investigated the possibility of employing aryl boronic acids (analogous to a Suzuki–Miyaura coupling),¹¹ plus aryl halides and terminal alkynes (analogous to a Sonogashira coupling)¹² in a Sonogashira/cyclization/Suzuki–Miyaura sequential process. However, there was no evidence for formation of the expected 2,3-disubstituted benzo[*b*]furans in the two cases we examined using our standard reaction conditions. The major product in both cases was the corresponding 3*H*-benzofuran. This does not mean that there are no reaction conditions under which the desired Sonogashira/cyclization/Suzuki–Miyaura process won't occur.

We have also attempted Pd-catalyzed couplings with the 5bromobenzofurans **16** and **17** prepared by our benzofuran synthesis to illustrate how our benzofurans can be further diversified to provide a large variety of multisubstituted benzofurans for drug testing. Thus, Suzuki–Miyaura¹¹ and Mizoroki–Heck¹² couplings proceeded smoothly, affording the benzofurans **69** and **70** in 83% and 81% yields, respectively (Scheme 4).

3. Conclusions

A novel convenient multicomponent process for the synthesis of 2,3-disubstituted benzo[b]furans under Sonogashira conditions has been developed and the scope of this process studied. Microwave irradiation has been employed, providing higher yields, shorter reaction times and cleaner products. This methodology has proven quite general and should prove a valuable tool in the synthesis of combinatorial libraries of benzofurans.

4. Experimental

4.1. General remarks

All microwave reactions were carried out in sealed oven-dried microwave vials. A Biotage microwave reactor was used for all experiments run at or above 60 °C. A CEM microwave reactor was used for the room temperature microwave reactions. The ¹H and ¹³C NMR spectra were recorded at 300 and 75.5 MHz or 400 and 100 MHz, respectively. The chemical shifts of the ¹H NMR and ¹³C NMR spectra are reported relative to the residual signal of CDCl₃ (δ 7.26 ppm for the ¹H NMR and δ 77.23 ppm for the ¹³C NMR), acetone-*d*₆ (2.05 ppm for the ¹H NMR and δ 29.92 ppm for the ¹³C NMR) or DMSO-*d*₆ (2.50 ppm for the ¹H NMR and δ 39.51 ppm for the ¹³C NMR). All coupling constants (*J*) are reported in Hertz (Hz).



Scheme 4. Pd-catalyzed diversification.

All commercially obtained chemicals were used as received without further purification. Thin layer chromatography was performed using commercially prepared 60-mesh silica gel plates, and visualization was effected with short wavelength UV light (254 nm). All melting points were obtained using an EZ-Melt automated melting point apparatus and are uncorrected. High resolution mass spectra (HRMS) were obtained using an Agilent QTOF 6540 mass spectrometer (ESI at a voltage of 70 eV). All mass spectra (MS) were obtained using a GCT-Agilent 6890 gas chromatograph/mass spectrometer (EI at a voltage of 70 eV). All IR spectra were obtained using a Nicolet 380 FT-IR apparatus.

4.2. Preparation of the starting compounds for the threecomponent coupling

A majority of the starting materials were purchased from commercial sources and used as received. The following compounds were prepared following the procedure described in the literature: iodophenols **8**¹³ and **12**;¹⁴ 3-iodo-*N*-tosylindole (**49**),¹⁵ 3-iodo-4*H*chromen-4-one (**51**),¹⁶ 2-iodo-4,4-dimethylcyclohexenone (**63**).¹⁷

4.2.1. 4-Bromo-2-iodo-6-methoxyphenol (56). Phenol 56 was prepared by a modification of a procedure described for the bromination of guaiacol.¹⁸ 2-Iodo-6-methoxyphenol (**8**, 0.975 g, 3.9 mmol) was dissolved in dry DMF (0.8 mL), the reaction mixture was cooled to 0 °C, and then NBS (0.69 g, 3.9 mmol) in DMF (0.8 mL) was added dropwise. The reaction mixture was stirred for 30 min at 0 °C and then slowly guenched with an ice cold water/ethyl ether mixture at the same temperature. (The yield of the product dropped significantly when the reaction temperature was not kept at or below 0 °C). The organic fraction was separated, washed, and dried over MgSO₄. The reaction mixture was purified using column chromatography and the desired compound 56 was obtained as a brown solid (0.96 g, 75%): mp 77-79 °C; ¹H NMR (300 MHz, CDCl₃) § 3.88 (s, 4H), 6.04 (s, 1H), 6.94 (d, J=2.1 Hz, 1H), 7.43 (d, *I*=2.1 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 56.7, 81.6, 112.5, 114.3, 132.4, 145.3, 146.4.

4.2.2. 4-Ethynyl-2-methoxyphenyl acetate (**57**). Commercially available 4-bromo-2-methoxyphenol (1.02 g, 5.0 mmol) and acetic anhydride (0.71 mL, 7.5 mmol) were dissolved in dichloromethane (10 mL). Then concentrated H_2SO_4 (25 mg) was added and the mixture was stirred for 30 min at rt. The reaction was then subjected to an aqueous work-up analogous to the one described in the literature,¹⁹ resulting in

4-bromo-2-methoxyphenyl acetate, obtained as a colorless solid, 1.21 g (99%): ¹H NMR (300 MHz, CDCl₃) δ 2.31 (s, 3H), 3.82 (s, 3H), 6.90 (d, J=8.6 Hz, 1H), 7.03-7.12 (m, 2H). 4-Bromo-2-methoxyphenyl acetate (1.21 g, 4.9 mmol), Pd(OAc)₂ (53.8 mg, 0.24 mmol), CuI (23 mg, 0.12 mmol). and tris(*tert*-butyl)phosphoniumtetrafluoroborate (69.6 mg, 0.24 mmol) were dissolved in diisopropylamine (10 mL) and purged with argon. Trimethylsilyl acetylene (1.38 mL, 9.8 mmol) was added and the reaction mixture was stirred at 40 °C for 2 h. The workup was conducted analogous to a procedure described in the literature.²⁰ The desired alkyne was obtained as a colorless solid (1.28 g, 100%). The resulting product was dissolved in THF (29 mL) and water (3.5 mL) and a 1 M solution of TBAF in THF (5.7 mL) was added at 0 °C. The reaction mixture was allowed to warm to room temperature and stirred for an additional 1 h. The volatile solvents were evaporated and the aqueous layer was extracted with ethyl acetate. The organic fractions were dried and evaporated, affording 0.74 g (80%) of the desired compound **57** as a colorless solid: mp 77–79 °C; ¹H NMR (400 MHz, CDCl₃) δ 2.31 (s, 3H), 3.07 (s, 1H), 3.82 (s, 3H), 6.99 (d, J=8.0 Hz, 1H), 7.06–7.12 (m, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 20.7, 56.0, 77.3, 83.2, 116.0, 120.8, 123.0, 125.1, 140.5, 150.9, 168.8.

4.2.3. 3,5-Diacetoxyiodobenzene (**58**). 3,5-Dihydroxyiodobenzene (0.27 g, 1.16 mmol) was dissolved in dichloromethane (3 mL); Ac₂O (0.33 mL, 3.49 mmol) and H₂SO₄ (1.2 mg) were added and the mixture was stirred at rt for 1 h. Then concentrated aq NaHCO₃ solution was added at 0 °C and the mixture was allowed to warm up to room temperature. The organic phase was collected, dried (MgSO₄) and evaporated. Compound **58** was obtained as a colorless solid (0.34 g, 91%) and used without further purification: mp 77–80 °C; ¹H NMR (400 MHz, CDCl₃) δ 2.28 (s, 3H), 6.92 (t, *J*=2.0 Hz, 1H), 7.36 (d, *J*=2.0 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 21.3, 69.5, 92.7, 115.6, 128.4, 151.3, 168.8; HRMS calcd for C₁₀H₉IO₄ [M+Na]⁺ 342.9438, found 342.9441.

4.2.4. 4-(1,3-Dioxolan-2-yl)-2-iodo-6-methoxyphenol (**60**). Compound **60** was prepared following a procedure described for an analogous reaction.²¹ 5-Iodovanillin (1.0 mmol) and ethylene glycol (5.0 mmol) were dissolved in toluene. Then acidic aluminum oxide was added and the resulting mixture was refluxed for 24 h. After cooling, the mixture was filtered, washed with dichloromethane/ water and the organic phase was dried (MgSO₄) and evaporated. Column chromatography using ethyl acetate/hexanes (1:3) as the eluent afforded 196 mg (62%) of product **60** as a colorless oil: ¹H NMR (400 MHz, CDCl₃) δ 3.91 (s, 3H), 3.97–4.06 (m, 2H), 4.08–4.16 (m, 2H), 5.69 (s, 1H), 6.16 (s, 1H), 6.96 (s, 1H), 7.42 (s, 1H); ^{13}C NMR (101 MHz, CDCl₃) δ 56.5, 65.5, 80.9, 95.7, 103.0, 108.9, 129.1, 131.6, 146.2, 146.6.

4.3. General procedure for the one-pot, three-component Sonogashira/Cacchi type coupling to synthesize benzofurans

The 2-iodophenol (0.5 mmol) and dichlorobis(triphenylphosphine)palladium (10.5 mg, 3 mol %) were placed in a 5 mL microwave vial and purged with argon. Dry THF (0.5 mL) was added and the reaction mixture was stirred until the iodophenol completely dissolved. Then dry triethylamine (1.0 mL) and a 3.8 M solution of Cul in dry triethylamine (0.5 mL) were added and the mixture allowed to stir for 10 min. Then 1.2 equiv of the corresponding alkyne was added; the vial was capped, purged with argon and placed in the microwave reactor for 30 min at 25 °C. The corresponding aryl iodide (0.5 mmol) and dry acetonitrile (2 mL) were added and the reaction mixture was heated in the microwave reactor at 100 °C for 25 min. After cooling, the solvents were evaporated and column chromatography using ethyl acetate/ hexane as the eluent afforded the desired products.

4.3.1. Ethyl 4-(2-phenylbenzofuran-3-yl)benzoate (**7**). Yellow solid, 164.2 mg (96%): mp 100–103 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.42 (t, *J*=7.1 Hz, 3H), 4.41 (q, *J*=7.1 Hz, 2H), 7.21–7.26 (m, 1H), 7.26–7.36 (m, 4H), 7.49 (d, *J*=7.3 Hz, 1H), 7.52–7.66 (m, 5H), 8.14 (d, *J*=8.1 Hz, 2H); ¹³C NMR (101 MHz, CDCl₃) δ 14.6, 61.3, 111.4, 116.7, 119.9, 123.3, 125.1, 127.4, 128.7, 128.9, 129.8, 129.8, 130.4, 131.6, 131.9, 137.9, 151.3, 154.2, 166.6; HRMS calcd for C₂₃H₁₈O₃ [M+H]⁺ 342.1256, found 342.1260.

4.3.2. Ethyl 4-(7-methoxy-2-phenylbenzofuran-3-yl)benzoate (**9**). Bright yellow solid, 169.6 mg (91%): mp 125–127 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.44 (t, *J*=7.1 Hz, 3H), 4.08 (s, 3H), 4.43 (q, *J*=7.1 Hz, 2H), 6.87 (d, *J*=7.9 Hz, 1H), 7.10 (d, *J*=7.8 Hz, 1H), 7.19 (t, *J*=7.9 Hz, 1H), 7.28–7.36 (m, 3H), 7.59 (d, *J*=8.0 Hz, 2H), 7.61–7.70 (m, 2H), 8.14 (d, *J*=8.0 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 14.6, 56.4, 61.3, 107.4, 112.3, 117.0, 124.0, 127.5, 128.7, 128.9, 129.8, 129.9, 130.3, 130.3, 131.4, 137.9, 143.6, 145.6, 151.6, 166.6; HRMS calcd for C₂₄H₂₀O₄ [M+H]⁺ 372.1362, found 373.1438.

4.3.3. *Methyl* 3-[4-(*ethoxycarbonyl*)*phenyl*]-2-*phenylbenzofuran*-5*carboxylate* (**11**). Colorless solid, 162.3 mg (92%): mp 186–187 °C; ¹H NMR (300 MHz, CDCl₃) δ 1.44 (t, J=7.1 Hz, 3H), 3.92 (s, 3H), 4.44 (q, J=7.1 Hz, 2H), 7.29–7.47 (m, 3H), 7.46–7.73 (m, 5H), 8.09 (d, J=8.6 Hz, 1H), 8.17 (d, J=8.2 Hz, 2H), 8.20 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 14.6, 52.4, 61.4, 111.4, 117.1, 122.4, 125.8, 126.9, 127.4, 128.9, 129.4, 129.9, 129.9, 130.2, 130.6, 137.2, 152.7, 156.8, 166.5, 167.3; HRMS calcd for C₂₅H₂₀O₅ [M+H]⁺ 401.1384, found 401.1395.

4.3.4. Ethyl 4-(7-allyl-2-phenylbenzofuran-3-yl)benzoate (**14**). Colorless oil, 97.6 mg (53%): ¹H NMR (300 MHz, CDCl₃) δ 1.38 (t, *J*=7.1 Hz, 3H), 3.79 (d, *J*=6.7 Hz, 2H), 4.39 (q, *J*=7.1 Hz, 2H), 5.11–5.33 (m, 2H), 5.06–5.38 (m, 2H), 6.16 (ddt, *J*=16.8, 10.0, 6.6 Hz, 1H), 7.11–7.40 (m, 6H), 7.53 (t, *J*=7.7 Hz, 1H), 7.59–7.73 (m, 3H), 8.10 (dt, *J*=7.8, 1.5 Hz, 1H), 8.23 (t, *J*=1.8 Hz, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 14.5, 29.9, 30.2, 34.2, 61.3, 116.5, 117.0, 118.1, 123.5, 123.9, 125.1, 127.2, 128.7, 129.0, 129.3, 130.0, 130.7, 131.0, 131.6, 133.7, 134.5, 136.1, 150.9, 152.6, 166.6; HRMS calcd for C₂₆H₂₂O₃ [M+H]⁺ 382.1563, found 382.1558.

4.3.5. *Ethyl* 4-(5-*bromo-2-phenylbenzofuran-3-yl)benzoate* (**16**). Colorless solid, 176.8 mg (84%): mp 135–137 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.44 (t, *J*=7.1 Hz, 3H), 4.43 (q, *J*=7.1 Hz, 2H), 7.29–7.37 (m, 3H), 7.44 (d, *J*=1.0 Hz, 2H), 7.56 (d, *J*=8.3 Hz, 2H), 7.58–7.64 (m, 3H), 8.15 (d, *J*=8.2 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 14.6, 61.4, 112.9, 116.3, 116.6, 122.7, 127.5, 128.0, 128.9, 129.4, 129.8, 129.9, 130.2, 130.6, 131.9, 137.2,

152.6, 153.0, 166.5; HRMS calcd for $C_{23}H_{17}Br$ $O_3\ [M+H]^+$ 421.0361, found 421.0434.

4.3.6. *Ethyl* 3-(5-bromo-2-phenylbenzofuran-3-yl)benzoate (**17**). Colorless solid: mp 103–104 °C; ¹H NMR (300 MHz, CDCl₃) δ 1.39 (t, *J*=7.1 Hz, 3H), 4.40 (q, *J*=7.1 Hz, 2H), 7.28–7.35 (m, 3H), 7.44 (d, *J*=1.2 Hz, 2H), 7.50–7.71 (m, 5H), 8.12 (dt, *J*=7.6, 1.4 Hz, 1H), 8.17 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 14.5, 61.4, 112.9, 116.2, 116.5, 122.7, 126.4, 127.2, 127.9, 128.8, 128.9, 129.2, 129.4, 129.5, 130.0, 130.9, 131.8, 132.3, 132.8, 134.4, 152.4, 152.9, 166.5; HRMS calcd for C₂₃H₁₈BrO₃ [M+H]⁺ 421.0434, found 421.0440.

4.3.7. Ethyl 4-[2-(4-methoxyphenyl)benzofuran-3-yl]benzoate (**19**). Yellow solid, 175.0 mg (94%): mp 125–127 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.41 (t, *J*=7.1 Hz, 3H), 3.78 (s, 3H), 4.41 (q, *J*=7.1 Hz, 2H), 6.83 (d, *J*=8.8 Hz, 2H), 7.22 (t, *J*=7.5 Hz, 1H), 7.29 (t, *J*=7.7 Hz, 1H), 7.47 (d, *J*=7.7 Hz, 1H), 7.52–7.61 (m, 5H), 8.13 (d, *J*=8.2 Hz, 2H); ¹³C NMR (101 MHz, CDCl₃) δ 14.6, 55.4, 61.2, 111.2, 114.2, 115.2, 119.6, 122.9, 123.2, 124.6, 128.9, 129.5, 129.8, 130.3, 134.1, 138.2, 151.5, 154.1, 160.1, 166.6; HRMS calcd for C₂₄H₂₀O₄ [M+H]⁺ 372.1362, found 372.1372.

4.3.8. *Ethyl* 4-[2-(3,5-*dimethoxyphenyl*)*benzofuran*-3-*yl*]*benzoate* (**21**). Cream colored solid, 186.3 mg (93%): mp 102–104 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.45 (t, *J*=7.1 Hz, 3H), 3.70 (s, 6H), 4.44 (q, *J*=7.1 Hz, 2H), 6.45 (s, 1H), 6.81 (s, 2H), 7.27 (t, *J*=7.5 Hz, 1H), 7.37 (t, *J*=7.7 Hz, 1H), 7.49 (d, *J*=7.8 Hz, 1H), 7.58 (d, *J*=8.2 Hz, 1H), 7.63 (d, *J*=8.0 Hz, 2H), 8.18 (dd, *J*=8.2, 1.5 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 14.6, 55.5, 61.3, 101.6, 101.6, 105.2, 111.4, 117.1, 119.9, 123.3, 125.2, 129.8, 129.9, 130.3, 131.9, 137.9, 150.9, 154.0, 160.8, 166.5; HRMS calcd for C₂₅H₂₂O₅ [M+H]⁺ 403.1540, found 403.1549.

4.3.9. *Ethyl* 4-[2-(4-(*dimethylamino*)*phenyl*)*benzofuran*-3-*yl*]*benzoate* (**23**). Yellow-green oil, 160.1 mg (83%): ¹H NMR (300 MHz, CDCl₃) δ 1.47 (t, *J*=7.2 Hz, 3H), 2.99 (s, 6H), 4.47 (q, *J*=7.1 Hz, 2H), 7.29 (dt, *J*=17.4, 7.0 Hz, 2H), 6.66 (d, *J*=8.5 Hz, 2H), 7.29 (dt, *J*=17.4, 7.0 Hz, 2H), 6.66 (d, *J*=8.5 Hz, 2H), 7.29 (dt, *J*=17.4, 7.0 Hz, 2H), 7.48-7.61 (m, 4H), 7.67 (d, *J*=8.0 Hz, 2H), 8.19 (d, *J*=8.0 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 14.5, 40.3, 61.1, 111.1, 111.9, 113.7, 117.9, 119.2, 123.0, 124.1, 128.5, 129.2, 129.8, 130.1, 130.2, 138.8, 150.6, 152.7, 153.9, 166.7; HRMS calcd for C₂₅H₂₃NO₃ [M+H]⁺ 386.1751, found 386.1759.

4.3.10. Ethyl 4-[2-(2-formylphenyl)benzofuran-3-yl]benzoate (**25**). Yellow oil, 128.6 mg (69%): ¹H NMR (400 MHz, CDCl₃) δ 1.36 (t, *J*=7.2 Hz, 3H), 4.36 (q, *J*=7.2 Hz, 2H), 7.36 (t, *J*=7.5 Hz, 1H), 7.43 (t, *J*=7.6 Hz, 2H), 7.47–7.64 (m, 5H), 7.73 (d, *J*=7.8 Hz, 1H), 8.01 (t, *J*=7.6 Hz, 2H), 8.14 (s, 1H), 10.07 (s, 1H); ¹³C NMR (101 MHz, CDCl₃) δ 14.5, 61.3, 111.8, 117.6, 120.5, 120.7, 123.8, 125.8, 128.3, 128.3, 129.0, 129.3, 129.8, 130.5, 131.3, 131.6, 131.9, 132.9, 133.8, 134.3, 148.6, 155.1, 166.2, 191.0; HRMS calcd for C₂₄H₁₈O₄ [M+H]⁺ 371.1278, found 371.1276.

4.3.11. Ethyl 4-[2-(4-cyanophenyl)benzofuran-3-yl]benzoate (**27**). Colorless solid, 97.5 mg (53%): mp 135–137 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.45 (t, *J*=7.1 Hz, 3H), 4.45 (q, *J*=7.1 Hz, 2H), 7.29 (t, *J*=7.5 Hz, 1H), 7.41 (t, *J*=7.8 Hz, 1H), 7.48 (d, *J*=7.8 Hz, 1H), 7.57 (t, *J*=8.1 Hz, 5H), 7.73 (d, *J*=8.1 Hz, 2H), 8.19 (d, *J*=8.0 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 14.6, 56.9, 61.4, 111.6, 111.9, 118.8, 119.7, 120.5, 123.8, 126.3, 127.3, 129.6, 129.8, 130.5, 130.7, 132.5, 134.7, 136.9, 148.7, 154.4, 166.4; HRMS calcd for C₂₄H₁₇NO₃ [M+H]⁺ 367.1208, found 367.1212.

4.3.12. Ethyl 4-[2-(cyclohex-1-en-1-yl)benzofuran-3-yl]benzoate (**29**). Yellow oil, 104.1 mg (52%): ¹H NMR (400 MHz, CDCl₃) δ 1.40 (t, J=7.1 Hz, 3H), 1.62 (t, J=3.2 Hz, 4H), 2.12 (br s, 2H), 2.18 (br s, 2H), 3.88 (s, 3H), 4.40 (q, *J*=7.1 Hz, 2H), 6.41–6.48 (m, 1H), 7.47 (d, *J*=8.5 Hz, 1H), 7.54 (t, *J*=7.7 Hz, 1H), 7.61–7.66 (m, 1H), 8.01 (dd, *J*=10.8, 1.5 Hz, 2H), 8.09 (d, *J*=7.8 Hz, 1H), 8.13 (d, *J*=1.9 Hz, 1H); 13 C NMR (100 MHz, CDCl₃) δ 14.5, 21.9, 22.6, 25.9, 26.6, 52.2, 61.3, 110.8, 115.5, 121.9, 125.2, 126.4, 127.9, 128.9, 128.9, 130.6, 131.0, 131.1, 131.2, 133.2, 134.5, 154.5, 156.0, 166.6, 167.4; HRMS calcd for C₂₅H₂₅O₅ [M+H]⁺ 405.1697, found 405.1706.

4.3.13. Ethyl 4-[2-(thiophen-3-yl)benzofuran-3-yl]benzoate (**31**). Yellow oil, 174.2 mg (100%): ¹H NMR (300 MHz, CDCl₃) δ 1.45 (t, *J*=7.2 Hz, 3H), 4.44 (q, *J*=7.1 Hz, 2H), 7.21 (t, *J*=5.2 Hz, 1H), 7.24–7.30 (m, 2H), 7.35 (t, *J*=7.6 Hz, 1H), 7.47 (d, *J*=7.2 Hz, 1H), 7.55 (d, *J*=8.1 Hz, 1H), 7.58–7.70 (m, 3H), 8.18 (d, *J*=8.4 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 14.6, 61.3, 111.3, 115.9, 119.8, 123.4, 123.8, 125.0, 126.1, 126.2, 129.7, 130.0, 130.3, 131.6, 137.7, 148.3, 154.0, 166.6; HRMS calcd for C₂₁H₁₆O₃S [M+H]⁺ 349.0893, found 349.0900.

4.3.14. Ethyl 4-[2-(1-methyl-1H-pyrazol-5-yl)benzofuran-3-yl]benzoate (**33**). Green amorphous solid, 109.4 mg (63%): ¹H NMR (400 MHz, CDCl₃) δ 1.41 (t, *J*=7.1 Hz, 3H), 3.63 (s, 3H), 4.40 (q, *J*=7.1 Hz, 2H), 7.31 (t, *J*=7.3 Hz, 1H), 7.38 (t, *J*=7.3 Hz, 1H), 7.42–7.49 (m, 1H), 7.55 (t, *J*=6.8 Hz, 3H), 7.60–7.72 (m, 2H), 8.11 (d, *J*=7.7 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 14.6, 61.3, 111.6, 120.2, 123.7, 125.5, 128.1, 128.6, 128.7, 129.1, 129.9, 130.4, 132.1, 132.2, 132.3, 136.8, 154.8, 166.4 (N–CH₃ does not show up); HRMS calcd for C₂₁H₁₈N₂O₃ [M+H]⁺ 347.1390, found 347.1398.

4.3.15. 2,3-Diphenylbenzofuran (**34**).²² Yellow-green solid, 117.6 mg (87%): mp 116–119 °C [lit. mp 123 °C];²³ ¹H NMR (300 MHz, CDCl₃) δ 7.20–7.36 (m, 5H), 7.46 (m, 6H), 7.56 (d, *J*=8.2 Hz, 1H), 7.66 (d, *J*=7.3 Hz, 2H); ¹³C NMR (75 MHz, CDCl₃) δ 111.3, 117.7, 120.3, 123.1, 124.9, 127.2, 127.8, 128.6, 128.6, 129.2, 129.9, 130.5, 130.9, 133.1, 150.7, 154.2.

4.3.16. 3-(4-Methoxyphenyl)-2-phenylbenzofuran (**36**).²⁴ Yellow amorphous solid, 79.7 mg (53%): ¹H NMR (400 MHz, CDCl₃) δ 3.89 (s, 3H), 6.98–7.05 (m, 2H), 7.23–7.27 (m, 1H), 7.28–7.36 (m, 4H), 7.41–7.45 (m, 2H), 7.50 (d, *J*=8.5 Hz, 1H), 7.56 (d, *J*=8.2 Hz, 1H), 7.69 (dd, *J*=8.2, 1.6 Hz, 2H).

4.3.17. 3-(3-*Methoxyphenyl*)-2-*phenylbenzofuran* (**38**).⁷ Yellow oil, 125.7 mg (84%): ¹H NMR (400 MHz, CDCl₃) δ 3.83 (s, 3H), 6.98–7.03 (m, 1H), 7.13 (dd, *J*=11.5, 4.9 Hz, 2H), 7.24–7.45 (m, 6H), 7.56–7.63 (m, 2H), 7.75 (dd, *J*=8.1, 1.5 Hz, 2H).

4.3.18. 3-(4-Cyanophenyl)-2-phenylbenzofuran (**40**). Yellow solid, 140.1 mg (98%): mp 113–115 °C, ¹H NMR (400 MHz, CDCl₃) δ 7.27 (t, *J*=7.5 Hz, 1H), 7.32–7.39 (m, 4H), 7.48 (d, *J*=7.7 Hz, 1H), 7.59 (dt, *J*=11.4, 4.6 Hz, 5H), 7.69–7.78 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 111.4, 111.6, 115.9, 118.9, 119.6, 123.6, 125.4, 127.5, 128.9, 129.2, 129.3, 130.0, 130.5, 132.9, 138.3, 151.8, 154.3; HRMS calcd for C₂₁H₁₃NO [M+H]⁺ 296.107, found 296.1073.

4.3.19. 3-(4-Nitrophenyl)-2-phenylbenzofuran (**42**).⁷ Yellow solid, 114.5 mg (73%): mp 135–138 °C; ¹H NMR (300 MHz, CDCl₃) δ 7.26–7.28 (m, 1H), 7.32 (d, *J*=7.6 Hz, 1H), 7.36–7.39 (m, 4H), 7.52 (d, *J*=7.8 Hz, 1H), 7.60 (d, *J*=7.2 Hz, 3H), 7.70 (d, *J*=8.2 Hz, 2H), 8.33 (dd, *J*=8.7, 1.7 Hz, 2H).

4.3.20. 3-(2-Nitrophenyl)-2-phenylbenzofuran (**44**). Yellow crystals, 116.6 mg (74%): mp 119–122 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.17–7.25 (m, 2H), 7.28–7.39 (m, 4H), 7.51 (dd, *J*=7.5, 1.7 Hz, 1H), 7.52–7.60 (m, 3H), 7.62 (td, *J*=7.7, 1.6 Hz, 1H), 7.68 (td, *J*=7.5, 1.5 Hz, 1H), 8.13 (dd, *J*=8.0, 1.4 Hz, 1H); ¹³C NMR (101 MHz, CDCl₃) δ 111.6, 113.4, 119.5, 123.5, 125.2, 125.2, 126.9, 128.2, 128.9, 128.9, 129.4,

129.8, 130.1, 133.6, 133.6, 149.9, 151.4, 154.0; HRMS calcd for $C_{20}H_{13}NO_3$ [M+H]⁺ 316.0968, found 316.0977.

4.3.21. 3-(4-Chlorophenyl)-2-phenylbenzofuran (**46**). Yellow solid, 145.6 mg (96%): mp 100–101 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.24–7.28 (m, 1H), 7.35 (dd, *J*=10.2, 5.7 Hz, 4H), 7.45 (s, 4H), 7.47 (d, *J*=7.8 Hz, 1H), 7.57 (d, *J*=8.2 Hz, 1H), 7.64 (dd, *J*=7.9, 1.7 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 111.4, 116.5, 119.9, 123.3, 125.1, 127.3, 128.7, 128.8, 129.5, 130.0, 130.5, 131.3, 131.5, 133.7, 143.7, 150.9; HRMS calcd for C₂₀H₁₃ClO [M+H]⁺ 305.0728, found 305.0731.

4.3.22. 1-[4-(2-(Thiophen-3-yl)benzofuran-3-yl)phenyl]ethanone(**48**). Cream colored solid, 116.1 mg (73%): mp 172–174 °C; ¹H NMR (400 MHz, CDCl₃) δ 2.68 (s, 3H), 7.20 (d, *J*=5.1 Hz, 1H), 7.27 (dd, *J*=7.3, 5.5 Hz, 2H), 7.35 (t, *J*=7.7 Hz, 1H), 7.47 (d, *J*=7.8 Hz, 1H), 7.55 (d, *J*=8.1 Hz, 1H), 7.65 (dd, *J*=8.6, 5.6 Hz, 3H), 8.09 (d, *J*=8.0 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 26.9, 111.4, 115.8, 119.8, 123.4, 123.9, 125.1, 126.1, 126.3, 129.1, 129.6, 130.2, 131.6, 136.5, 138.1, 148.3, 154.0, 197.9; HRMS calcd for C₂₀H₁₄O₂S [M+H]⁺ 318.0715, found 318.0731.

4.3.23. 3-(2-Phenylbenzofuran-3-yl)-1-tosyl-1H-indole (**50**). Yellow amorphous solid, 134.1 mg (58%): ¹H NMR (400 MHz, CDCl₃) δ 2.40 (s, 3H), 7.14 (t, *J*=7.5 Hz, 1H), 7.25 (ddd, *J*=21.5, 15.1, 8.0 Hz, 7H), 7.37 (dd, *J*=15.9, 8.5 Hz, 3H), 7.57–7.64 (m, 3H), 7.76 (s, 1H), 7.85 (d, *J*=8.2 Hz, 2H), 8.13 (d, *J*=8.3 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 21.9, 108.1, 111.5, 114.1, 114.9, 120.3, 121.4, 123.2, 123.7, 125.1, 125.3, 125.4, 126.9, 127.1, 128.6, 128.7, 130.0, 130.2, 130.5, 130.6, 135.4, 135.7, 145.3, 152.1, 154.3; HRMS calcd for C₂₉H₂₁NO₃S [M+H]⁺ 463.1242, found 463.1315.

4.3.24. 3-(2-Phenylbenzofuran-3-yl)-4H-chromen-4-one (**52**). Cream colored solid, 73.4 mg (43%): mp 183–184 °C; ¹H NMR (400 MHz, CDCl₃) δ 7.22–7.28 (m, 1H), 7.29–7.40 (m, 4H), 7.40–7.45 (m, 1H), 7.50 (ddd, *J*=8.2, 7.2, 1.1 Hz, 1H), 7.53–7.59 (m, 2H), 7.73–7.78 (m, 3H), 8.01 (s, 1H), 8.36 (dd, *J*=8.0, 1.7 Hz, 1H); ¹³C NMR (101 MHz, CDCl₃) δ 107.8, 111.5, 118.3, 118.5, 120.5, 123.3, 124.5, 125.0, 125.8, 126.7, 127.3, 128.9, 128.9, 130.4, 130.5, 134.1, 153.1, 154.3, 155.4, 156.7, 176.2; HRMS calcd for C₂₃H₁₄O₃ [M+H]⁺ 339.1016, found 339.1015.

4.3.25. 2-Fluoro-4-[7-methoxy-2-(m-tolyl)benzofuran-3-yl]nicotinaldehyde (**55**). Yellow solid, 116.0 mg (65%): mp 177–180 °C; ¹H NMR (300 MHz, CDCl₃) δ 2.33 (s, 3H), 4.09 (s, 3H), 6.83 (d, *J*=7.9 Hz, 1H), 6.90 (d, *J*=8.0 Hz, 1H), 7.19 (t, *J*=3.2 Hz, 4H), 7.36 (d, *J*=5.1 Hz, 1H), 7.47 (s, 1H), 8.46 (d, *J*=5.1 Hz, 1H), 10.05 (s, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 21.7, 56.5, 108.0, 111.5, 124.6, 124.9, 125.2, 125.2, 127.9, 128.9, 129.0, 130.7, 131.0, 139.1, 140.5, 143.6, 145.8, 149.3, 152.3, 152.6, 153.6, 161.4, 164.7, 187.1, 187.1 (extra peaks due to ¹³C–¹⁹F coupling); HRMS calcd for C₂₂H₁₆FNO₃ [M+H]⁺ 361.1114, found 362.1187.

4.3.26. 2-(4-Acetoxy-3-methoxyphenyl)-5-bromo-3-(3,5-diacetoxyphenyl)-7-methoxybenzofuran (**59**). Compound **59** was synthesized following the general procedure for a three-component Sonogashira/Cacchi type cyclization and was obtained as a brown amorphous solid (340.6 mg, 60%): ¹H NMR (400 MHz, CDCl₃) δ 2.29 (s, 6H), 2.31 (s, 3H), 3.67 (s, 3H), 4.03 (s, 3H), 6.95 (s, 1H), 7.00 (s, 1H), 7.03 (d, *J*=8.3 Hz, 1H), 7.09 (d, *J*=1.2 Hz, 2H), 7.15 (s, 1H), 7.21 (s, 1H), 7.35 (d, *J*=8.3 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 20.9, 21.3, 55.9, 56.6, 110.9, 111.0, 111.1, 114.9, 115.6, 115.7, 116.6, 119.8, 120.6, 123.3, 128.2, 132.6, 134.2, 140.4, 142.2, 145.7, 151.2, 151.5, 151.8, 168.9; HRMS calcd for C₂₈H₂₄BrO₉ [M+H]⁺ 583.0525, found 583.0598.

4.3.27. 2-(4-Acetoxy-3-methoxyphenyl)-3-(3,5-diacetoxyphenyl)-5-(1,3-dioxolan-2-yl)-7-methoxybenzofuran (**61**). Compound **61** was synthesized following the general procedure for the threecomponent Sonogashira/Cacchi type cyclization and was obtained as a bright yellow oil (340.6 mg, 63%): ¹H NMR (400 MHz, CDCl₃) δ 2.28 (s, 6H), 2.30 (s, 3H), 3.67 (s, 3H), 4.00–4.05 (m, 2H), 4.07 (s, 3H), 4.14–4.23 (m, 2H), 5.84 (s, 1H), 6.97–7.05 (m, 3H), 7.12 (d, *J*=2.2 Hz, 2H), 7.19 (d, *J*=13.7 Hz, 2H), 7.37 (dd, *J*=8.3, 2.0 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 20.8, 21.3, 55.9, 56.4, 65.5, 104.2, 105.5, 110.9, 111.1, 115.4, 116.5, 117.6, 119.7, 120.7, 123.2, 128.6, 131.2, 134.1, 134.8, 140.2, 143.8, 145.4, 151.1, 151.7, 168.9, 168.9; HRMS calcd for C₃₁H₂₉O₁₁ [M+H]⁺ 577.1704, found 577.1711.

4.3.28. 2-(4-Acetoxy-3-methoxyphenyl)-3-(3,5-diacetoxyphenyl)-5formyl-7-methoxybenzofuran (**62**). Compound **61** (85.4 mg, 0.155 mmol) was dissolved in THF (0.3 mL), 10% aq HCl (62 μ L) was added at 0 °C, and the reaction mixture was stirred for 20 min. An aqueous work-up was conducted according to a procedure described in the literature.²⁵ Compound **62** was obtained as a yellow oil (75.8 mg, 92%): ¹H NMR (400 MHz, CDCl₃) δ 2.29 (s, 6H), 2.30 (s, 3H), 3.67 (s, 3H), 4.08 (s, 3H), 6.99–7.06 (m, 2H), 7.14 (d, *J*=2.1 Hz, 2H), 7.18 (s, 1H), 7.37 (dd, *J*=8.3, 1.8 Hz, 1H), 7.40 (s, 1H), 7.61 (s, 1H), 9.96 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 20.9, 21.3, 55.9, 56.4, 105.1, 111.1, 115.7, 116.7, 118.5, 119.8, 120.6, 123.4, 127.9, 131.5, 133.9, 133.9, 140.6, 146.2, 146.8, 151.3, 151.9, 152.2, 168.9, 191.8, 191.9; HRMS calcd for C₂₉H₂₅O₁₀ [M+H]⁺ 533.1442, found 533.1443.

4.3.29. 4,4-Dimethyl-2-(2-phenylbenzofuran-3-yl)cyclohex-2-enone (**64**). Yellow solid, 54.3 mg (34%): mp 100–106 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.27 (s, 6H), 2.08 (t, *J*=6.8 Hz, 2H), 2.73 (t, *J*=6.8 Hz, 2H), 6.82 (s, 1H), 7.23 (t, *J*=7.5 Hz, 1H), 7.31 (d, *J*=8.8 Hz, 3H), 7.33–7.48 (m, 2H), 7.52 (d, *J*=8.1 Hz, 1H), 7.72 (d, *J*=7.3 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 27.8, 33.9, 35.3, 36.3, 111.3, 112.5, 115.5, 120.1, 122.9, 124.7, 127.0, 128.6, 130.2, 130.3, 130.9, 151.9, 154.0, 160.7, 197.2; HRMS calcd for C₂₂H₂₀O₂ [M+H]⁺ 339.1356, found 339.1348.

4.4. General procedure for the synthesis of benzofurans 67 and 68 by a three-component Sonogashira/Heck-type coupling

2-Iodophenol (0.5 mmol) and bis(triphenylphosphine)palladium dichloride (10.5 mg, 3 mol %) were placed in a 5 mL microwave vial and purged with argon. Dry THF (0.5 mL) was added and the reaction mixture was stirred until the iodophenol completely dissolved. Then dry triethylamine (1.0 mL) and a 3.8 M solution of CuI in dry triethylamine (0.5 mL) were added and the mixture allowed to stir for 10 min. Then 1.2 equiv of the corresponding alkyne was added; the vial was capped, purged with argon and placed in the microwave reactor for 30 min at 25 °C. The corresponding alkene (2.5 mmol), dry acetonitrile (2 mL), benzoquinone (0.5 mmol), and anhydrous KOAc (1.5 mmol) were added and the reaction mixture was heated in a microwave reactor at 60 °C for 25 min. After allowing the reaction to cool and a standard aqueous work-up, the reaction mixture was subject to column chromatography using ethyl acetate/hexane as the eluent to afford the desired products.

4.4.1. *n*-Butyl (*E*)-3-(2-phenylbenzofuran-3-yl)acrylate (**67**). Orange amorphous solid, 88.9 mg (56%): ¹H NMR (400 MHz, CDCl₃) δ 1.00 (t, *J*=7.4 Hz, 3H), 1.48 (h, *J*=7.3 Hz, 2H), 1.67–1.76 (m, 2H), 4.26 (t, *J*=6.6 Hz, 2H), 4.14–4.41 (m, 2H), 6.71 (s, 1H), 7.31–7.41 (m, 2H), 7.52 (dt, *J*=13.4, 6.9 Hz, 4H), 7.78 (d, *J*=8.0 Hz, 2H), 7.90 (d, *J*=7.5 Hz, 1H), 8.05 (d, *J*=16.0 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 14.0, 19.5, 31.0, 64.6, 111.8, 112.8, 115.5, 119.4, 121.2, 123.9, 125.5, 126.9, 128.7, 129.1, 129.9, 136.0, 154.7, 157.7, 167.6; HRMS calcd for C₂₁H₂₁O₃ [M+H]⁺ 321.1485, found 321.1491.

4.4.2. *n*-Butyl (*E*)-3-[2-(cyclohex-1-en-1-yl)benzofuran-3-yl]acrylate (**68**). Yellow amorphous solid, 113.6 mg (70%): ¹H NMR (400 MHz, CDCl₃) δ 0.99 (t, *J*=7.4 Hz, 3H), 1.45 (dd, *J*=14.8, 7.4 Hz,

2H), 1.68–1.82 (m, 6H), 2.32 (s, 2H), 2.52 (s, 2H), 4.24 (t, *J*=6.6 Hz, 2H), 6.33 (s, 1H), 6.56 (d, *J*=16.1 Hz, 1H), 7.26–7.31 (m, 2H), 7.45 (s, 1H), 7.81 (d, *J*=8.0 Hz, 1H), 7.95 (d, *J*=16.0 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 13.9, 19.4, 21.9, 22.5, 26.2, 26.6, 31.0, 64.4, 111.4, 111.8, 117.6, 120.9, 123.6, 124.9, 126.8, 128.4, 134.9, 136.7, 153.9, 160.6, 167.9; HRMS calcd for C₂₁H₂₄O₃ [M+H]⁺ 324.1725, found 325.1799.

4.5. Elaboration of the bromo-containing benzofurans by Pdcatalyzed couplings

4.5.1. Ethyl 4-[5-(4-methoxyphenyl)-2-phenylbenzofuran-3-yl]benzoate (69). In a 2 mL microwave vial, compound 16 (42.2 mg, 0.1 mmol), 4-methoxyphenylboronic acid (18.2 mg, 0.12 mmol), and tetrakis(triphenylphosphine)palladium (5.8 mg) were dissolved in a 1:1 mixture of EtOH/DMF (1.6 mL), then 1 M aq Cs₂CO₃ (0.25 mL) was added and the mixture was heated in a microwave reactor at 120 °C for 20 min. The mixture was diluted with saturated aq Na₂SO₄ and extracted with ethyl acetate $(3 \times 15 \text{ mL})$, dried (MgSO₄), and evaporated. Column chromatography using ethyl acetate/hexanes (1:10) as the eluent afforded 35.6 mg (80%) of benzofuran 69 as a colorless solid: mp 127–129 °C; ¹H NMR (400 MHz, CDCl₃) δ 1.44 (dd, *I*=7.4, 6.8 Hz, 3H), 3.85 (d, *I*=0.5 Hz, 3H), 4.44 (q, *I*=7.1 Hz, 2H), 6.97 (d, J=8.2 Hz, 2H), 7.30-7.38 (m, 3H), 7.53 (t, J=8.1 Hz, 3H), 7.63 (dt, J=17.1, 5.9 Hz, 6H), 8.17 (d, J=7.7 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 14.6, 55.6, 61.3, 111.5, 114.4, 116.9, 117.9, 124.5, 127.4, 128.6, 128.8, 128.9, 129.8, 129.9, 130.3, 130.4, 130.5, 134.2, 136.8, 137.9, 151.9, 153.6, 159.1, 166.6; HRMS calcd for C₃₀H₂₄O₄ [M+H]⁺ 449.1747, found 449.1675.

4.5.2. Ethyl 3-[5-((E)-3-butoxy-3-oxoprop-1-en-1-yl)-2-phenylbenzofuran-3-yl]benzoate (70). In a 2 mL vial, compound 17 (41.5 mg, 0.1 mmol), *n*-butyl acrylate (17.9 µL, 0.12 mmol), palladium acetate (0.6 mg) and SPhos (4.1 mg) were dissolved in DMF (0.5 mL). Then triethylamine (0.12 mL) was added and the mixture was heated at 100 °C for 24 h. The mixture was diluted with brine and extracted with ethyl ether (3×15 mL), dried (MgSO₄), and evaporated. Column chromatography using ethyl acetate/hexanes (1:10) as the eluent afforded 37.1 mg (81%) of benzofuran **70** as a cream colored solid: mp $77-80 \circ C$; ¹H NMR (400 MHz, CDCl₃) δ 0.96 (t, J=7.3 Hz, 3H), 1.42 (m, 5H), 1.69 (m, 2H), 4.20 (t, J=6.7 Hz, 2H), 4.41 (q, J=7.1 Hz, 2H), 6.41 (d, *I*=15.9 Hz, 1H), 7.29–7.36 (m, 3H), 7.50–7.68 (m, 7H), 7.76 (d, *J*=16.0 Hz, 1H), 8.14 (d, *J*=7.7 Hz, 1H), 8.21 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) § 13.9, 14.5, 19.4, 31.0, 61.4, 64.6, 111.9, 116.7, 117.4, 120.1, 125.2, 127.2, 128.8, 129.1, 129.3, 129.5, 130.0, 130.2, 130.9, 131.8, 132.9, 134.4, 144.9, 152.2, 155.2, 166.4, 167.4; HRMS calcd for C₃₀H₂₈O₅ [M+H]⁺ 468.1937, found 469.2010.

Acknowledgements

We gratefully acknowledge the National Science Foundation and the National Institutes of Health Kansas University Center of Excellence in Chemical Methodology and Library Development (P50 GM069663) for their generous financial support of this project. We would also like to thank the research groups of Professors Nicola Pohl and George Kraus for allowing us to use their CEM microwave reactors for the room temperature experiments.

Supplementary data

Copies of ¹H NMR and ¹³C NMR spectra of all novel compounds are available. Supplementary data related to this article can be found at http://dx.doi.org/10.1016/j.tet.2013.02.003.

References and notes

- For recent selected examples, see: (a) Boto, A.; Alvarez, L. *Heterocycles Nat. Prod.* Synth. 2011, 99–152; (b) Ashwood, V. A.; Field, M. J.; Horwell, D. C.; Julien-Larose, C.; Lewthwaite, R. A.; McCleary, S.; Pritchard, M. C.; Raphy, J.; Singh, L. J. Med. Chem. 2001, 44, 2276–2285.
- For some recent comprehensive reviews on the synthesis of benzofurans, see: (a) Cacchi, S.; Fabrizi, G.; Goggiamani, A. Org. Biomol. Chem. 2011, 9, 641–652; (b) Yeung, K.-S.; Peng, S.-X.; Wu, J.; Hou, X.-L. Prog. Heterocycl. Chem. 2011, 23, 195–229; (c) Wu, J. Mod. Heterocycl. Chem. 2011, 593–633; (d) Mentwally, M. A.; Abdel-Wahab, B. F.; Al-Hiti, G. A. Curr. Org. Chem. 2010, 14, 48–64; (e) Miyata, O.; Takeda, N.; Naito, T. Heterocycles 2009, 78, 843–871; (f) De Luca, L.; Nieddu, G.; Porcheddu, A.; Giacomelli, G. Curr. Med. Chem. 2009, 16, 1–20 and references therein.
- (a) Majumdar, K. C.; Chattopadhyay, B.; Maji, P. K.; Chattopadhyay, S. K.; Samanta, S. *Heterocycles* **2010**, *81*, 517–584; (b) Cacchi, S.; Fabrizi, G.; Goggiamani, A. *Curr. Org. Chem.* **2006**, *10*, 1423–1455; (c) Dudnik, A. S.; Gevorgyan, V. *Catal. Carbon-Heteroat. Bond Form.* **2011**, 317–410; (d) For examples of Pd-catalyzed C–H activation leading to benzofuran formation, see: Ferreira, E. M.; Zhang, H.; Stoltz, B. M. *Tetrahedron* **2008**, *64*, 5987–6001.
- Arcadi, A.; Cacchi, S.; Del Rosario, M.; Fabrizi, G.; Marinelli, F. J. Org. Chem. 1996, 61, 9280–9288.
- 5. Chaplin, J. H.; Flynn, B. L. Chem. Commun. 2001, 1594-1595.
- 6. Flynn, B. L.; Hamel, E.; Jung, M. K. J. Med. Chem. 2002, 45, 2670-2673.
- Hu, Y.; Nawoschik, K. J.; Liao, Y.; Ma, J.; Fathi, R.; Yang, Z. J. Org. Chem. 2004, 69, 2235–2239.
- 8. Chen, Y.; Markina, N. A.; Larock, R. C. Tetrahedron 2009, 65, 8908-8915.
- (a) Huang, K.-S.; Wang, Y.-H.; Li, R.-L.; Lin, M. Phytochemistry 2000, 54, 875–881;
 (b) Li, W.-L.; He, K.-K.; Li, Y.; Hou, Z.-J. Acta Chim. Sin. 2005, 63, 1607–1612.

- (a) Martinez, C.; Alvarez, R.; Aurrecoechea, J. M. Org. Lett. 2009, 11, 1083–1086;
 (b) Boyer, A.; Isono, N.; Lackner, S.; Lautens, M. Tetrahedron 2010, 66, 6468–6482; (c) Alvarez, R.; Martínez, C.; Madich, Y.; Denis, J. G.; Aurrecoechea, J. M.; de Lera, A. L. Chem.—Eur. J. 2010, 16, 12746–12753.
- For selected recent reviews, see: (a) Bubnov, Y. N. *Heterocycles* 2010, 80, 1–5; (b) McGlacken, G. P.; Bateman, L. M. *Chem. Soc. Rev.* 2009, 38, 2447–2464.
- For the most recent reviews, see: (a) Chinchilla, R.; Najera, C. Chem. Soc. Rev. 2011, 40, 5084–5121; (b) Chinchilla, R.; Najera, C. Chem. Rev. 2007, 107, 874–922.
- 13. Weeratunga, G.; Jaworska-Sobiesiak, A.; Horne, S.; Rodrigo, R. Can. J. Chem. 1987, 65, 2019–2023.
- 14. Larock, R. C.; Lee, N. H. J. Org. Chem. 1991, 56, 6253-6254.
- 15. Amjad, M.; Knight, D. W. Tetrahedron Lett. 2004, 45, 539-541.
- 16. Gammill, R. B. Synthesis **1979**, 901–903.
- Jan, N.-W.; Liu, H.-J. Org. Lett. 2006, 8, 151–153.
 Fujikawa, N.; Ohta, T.; Yamaguchi, T.; Fukuda, T.; Ishibashi, F.; Iwao, M. Tetrahedron 2006, 62, 594–604.
- Boudet, N.; Sase, S.; Sinha, P.; Liu, C.-Y.; Krasovskiy, A.; Knochel, P. J. Am. Chem. Soc. 2007, 129, 12358–12359.
- 20. Novák, Z.; Timári, G.; Kotschy, A. Tetrahedron 2003, 59, 7509-7513.
- Kamitori, Y.; Hojo, M.; Masuda, R.; Yoshida, T. Tetrahedron Lett. 1985, 26, 4767–4770
 - 22. Chiong, H. A.; Daugulis, O. Org. Lett. **2007**, 9, 1449–1451.
 - 23. Kamlet, M. J.; Dacons, J. C. J. Org. Chem. **1961**, 26, 220–225.
 - Ohba, H.; Ikeda, T.; Kobayashi, S.; Taniguchi, H. J. Chem. Soc., Chem. Commun. 1980, 988-989.
 - 25. Fetter, J.; Bertha, F.; Poszávácz, L.; Simig, G. J. Heterocycl. Chem. 2005, 42, 137–139.