



A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

GDCh

International Edition

www.angewandte.org

Accepted Article

Title: Visible-Light-Induced Selective Defluoroborylation of Polyfluoroarenes, gem-Difluoroalkenes, and Trifluoromethylalkenes

Authors: Wengang Xu, Heming Jiang, Jing Leng, Han-Wee Ong, and Jie Wu

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *Angew. Chem. Int. Ed.* 10.1002/anie.201911819
Angew. Chem. 10.1002/ange.201911819

Link to VoR: <http://dx.doi.org/10.1002/anie.201911819>
<http://dx.doi.org/10.1002/ange.201911819>

RESEARCH ARTICLE

Visible-Light-Induced Selective Defluoroborylation of Polyfluoroarenes, *gem*-Difluoroalkenes, and Trifluoromethylalkenes

Wengang Xu, Heming Jiang, Jing Leng, Han-Wee Ong, and Jie Wu*


Abstract: Fluorinated organoboranes serve as versatile synthetic precursors for the preparation of value-added fluorinated organic compounds. Recent progress has been mainly focused on the transition-metal catalyzed defluoroborylation. Herein, we report a photocatalytic defluoroborylation platform via direct B–H activation of *N*-heterocyclic carbene boranes, through the synergistic merger of a photoredox catalyst and a hydrogen atom transfer catalyst. This atom-economic and operationally simple protocol has enabled defluoroborylation of an extremely broad scope of multifluorinated substrates including polyfluoroarenes, *gem*-difluoroalkenes, and trifluoromethylalkenes in a highly selective fashion. Intriguingly, the defluoroborylation protocol can be transition-metal free, and the regioselectivity obtained is complementary to the reported transition-metal-catalysis in many cases.

Introduction


Approximately 20% of all pharmaceuticals and 30% of all agrochemicals contain carbon–fluorine bonds.^[1] There is currently an increasing demand for introduction of fluorinated building blocks into organic frameworks, and compounds of this sort have found a wide range of applications in the development of novel pharmaceutical agents,^[2] insecticides,^[3] catalysts,^[4] and materials.^[5] However, fluoroaromatic and vinyl fluoride compounds do not exist naturally and must be synthesized.^[6] Despite spectacular advances in single-site catalytic fluorination, synthetic access to polyfluorinated compounds in a selective fashion is still challenging.^[7] In this context, fluorinated organic boranes can serve as versatile synthetic precursors to obtain

distinct organic fluorides through a wide spectrum of established and reliable derivatization reactions.^[8] The most straightforward way to access fluorinated organic borane building blocks is selective C–F bond borylation of polyfluorinated organic compounds. However, this is inherently challenging for several reasons: 1) the C–F bond is among the most unreactive functional groups; 2) the high bond energy of metal–fluorine intermediates can lead to sluggish catalytic turnover in transition-metal-catalyzed C–F activation; and 3) boron reagents are fluorophilic and serve as fluorine scavengers in C–F derivatization.


A. Defluoroborylation of polyfluoroarenes and alkenes via transition-metal-catalysis



B. Photomediated C–F functionalization for C–C bond formation



C. Photomediated selective defluoroborylation of polyfluoroarenes and alkenes (this work)



Scheme 1. C–F bond activation of fluoroarenes and alkenes. TM = transition-metal catalyst; pinB-Bpin = bis(pinacolato)diboron; PC = photocatalyst; HAT = hydrogen atom transfer; LED = light-emitting diode.

Most success in catalytic defluoroborylation thus far has relied on catalysis by transition-metal, including Rh,^[9] Ni,^[10] Cu,^[11] Co,^[12] and Fe-based^[13] catalysis (Scheme 1a). However, higher temperatures are generally required in these reactions, and

[*] Dr. W. Xu, J. Leng, H.-W. Ong, Dr. J. Wu
Department of Chemistry, National University of Singapore
3 Science Drive 3, Singapore 117543, Republic of Singapore
E-mail: chmjie@nus.edu.sg
H. Jiang
Laboratory of Computational Chemistry & Drug Design, State Key
Laboratory of Chemical Oncogenomics, Peking University
Shenzhen Graduate School, Shenzhen, 518055, People's Republic
of China
J. Leng
State Key Laboratory of Silicate Materials for Architectures, Wuhan
University of Technology, 122 Luoshi Road, Wuhan, Hubei, 430070,
People's Republic of China
Dr. W. Xu
College of New Energy, Institute of New Energy, State Key
Laboratory of Heavy Oil Processing, China University of Petroleum
(East China), Qingdao, 266580, People's Republic of China
Dr. J. Wu
National University of Singapore (Suzhou) Research Institute, 377
Lin Quan Street, Suzhou Industrial Park, Suzhou, Jiangsu, 215123,
People's Republic of China

Supporting information for this article is given via a link at the end of
the document. CCDC 1894872 and 1937072.

RESEARCH ARTICLE


stoichiometric metallic additives are often employed to scavenge the fluoride ions generated in situ. Photocatalysis has witnessed dramatic developments over the past decade which have enabled previously inaccessible synthetic transformations.^[14] In particular, photolytically-induced borylation of C–F bonds in electron-rich monofluoroarenes has been realized by Li^[15] and by Larionov^[16] though the formation of triplet aryl cations by the heterolysis of C(sp²)–F bonds using strong ultraviolet (UV) light. Very recently, the group of Guo, Radius, Steffen, and Marder demonstrated a visible-light-promoted C–F borylation protocol that employs a Rh photosensitizer to accelerate the difficult transmetalation step in the nickel-catalyzed C–F activation.^[17]

Polyfluorinated arenes and alkenes have been applied to C–C bond formation via photocatalytic C–F bond functionalization. Weaver et al. have achieved a variety of C–C couplings of perfluoroarenes with alkenes, alkynes, arenes, and prenyl reagents by photoredox catalysis.^[18] This coupling was initiated by polyfluoroaryl radicals generated through light-promoted single electron reduction of polyfluoroarenes followed by extrusion of fluoride (Scheme 1b). Remarkably, Xie, Hashmi and co-workers recently demonstrated a photo-mediated mono-defluoroalkylation of *gem*-difluoroalkenes using *N*-aryl amines by a radical-radical coupling pathway.^[19a] They subsequently expanded the scope to polyfluoroarenes for controllable defluoroalkylation (Scheme 1b).^[19b] The Molander group achieved a fascinating synthesis of *gem*-difluoroalkene moieties by photoredox-induced carbon radical addition to α -trifluoromethylalkenes.^[20]

On the other hand, the development of boron-centered radical chemistry has recently gained increasing momentum, and is conceptually appealing because it may offer regio- or stereoselectivities different from those of transition-metal-catalysis.^[21] Elegant work from Fensterbank, Lacôte, Malacria, and Curran has shown that *N*-heterocyclic carbene (NHC)-boryl radicals can be generated by hydrogen atom abstraction from NHC-BH₃ complexes,^[21a,22] which are powerful intermediates useful in building a wide range of value-added boron compounds. For instance, Wang has developed radical borylation/cyclization cascade of 1,6-enynes with NHC-BH₃,^[21c] and Curran and Taniguchi reported the Sato-Myers cyclization triggered by NHC-boryl radicals.^[22d] Consistent with our continuing interest in development of atom-economic and redox-neutral transformations through hydrogen atom transfer (HAT)-based photocatalysis,^[23] we envisioned that a photo-mediated catalytic HAT process with NHC-BH₃ might deliver the boryl radical in a mild pathway, which could react with polyfluoroarenes and alkenes to introduce synthetically valuable fluorinated organic boranes (Scheme 1c). Notably, Lacôte and Lalevée developed visible-light induced NHC-boryl radical formation through a HAT process with thiyl radicals.^[22c] During the course of our investigation, the group of Xie and Zhu has reported an elegant study on photocatalytic version of this reaction to realize the inverse hydroboration of imines.^[24]

Results and Discussion

Table 1. Condition optimization for photo-mediated defluoroborylation of polyfluoroarenes




entry	deviation	Yield of 3a ^[a]
1	none	94 ^[b]
2	Ir[dF(CF ₃)ppy] ₂ (dtbbpy)PF ₆ (2.0 mol%)	93
3	Ir(ppy) ₃ instead of PC1	0
4	Ir(ppy) ₂ (dtbbpy)PF ₆ instead of PC1	88
5	Ir(dFppy) ₂ (dtbbpy)PF ₆ instead of PC1	87
6	4CzIPN (2 mol%) instead of PC1	76
7 ^[c]	4CzIPN (2 mol%) instead of PC1	90
8	without light or PC1	0
9	without S1	16
10	without base	68

Reaction conditions: hexafluorobenzene (**1a**, 0.20 mmol), NHC-borane (**2a**, 0.24 mmol), Ir[dF(CF₃)ppy]₂(dtbbpy)PF₆ (**PC1**, 1 mol%), **S1** (10 mol%), NaOAc (0.10 mmol), THF (4 mL), 18W blue LED irradiation, room temperature, 12 h. [a] Yields were determined by analysis of the crude ¹H NMR spectra using 1,3,5-trimethoxybenzene as an internal standard. [b] Isolated yields. [c] **S2** was used instead of **S1**. THF = tetrahydrofuran.

Our investigation was initiated with examination of the defluoroborylation of hexafluorobenzene (**1a**) with NHC-borane **2a** as the model substrates in the presence of a photoredox catalyst and a HAT catalyst under blue LED irradiation. After extensive evaluation (Table 1 and Table S1), we established that a combination of Ir[dF(CF₃)ppy]₂(dtbbpy)PF₆ (**PC1**) (1 mol%), thiol (**S1**) (10 mol%), and NaOAc (0.5 equiv) in THF (0.05 M) at ambient temperature was optimal and produced the desired defluoroborylation product (**3a**) in 94% isolated yield (Table 1, entry 1). Increasing the catalyst loading to 2 mol% gave a similar yield (entry 2). Ir(ppy)₃, which was previously employed in Weaver's photocatalytic hydrodefluorination,^[25] showed no catalytic activity in the defluoroborylation (entry 3). Other Ir photocatalysts, such as Ir(ppy)₂(dtbbpy)PF₆ and Ir(dFppy)₂(dtbbpy)PF₆ exhibited much lower efficiency (entries 4 and 5). Moderate yield could be obtained by using the organic dye, 2,4,5,6-tetra(9H-carbazol-9-yl)isophthalonitrile (4CzIPN) as the

RESEARCH ARTICLE



Scheme 2. Scope of photo-mediated selective defluoroborylation of polyfluoroarenes.^[a] [a] All isolated yields. Product was produced as a single regioisomer unless otherwise noted. Condition A was employed unless otherwise noted. [b] Condition B was employed. [c] **2a** (2.4 equiv) was used.

photocatalyst, which possesses redox properties similar to those of **PC1** (entry 6).^[26] The yield can be improved to 90% by replacing the thiol HAT catalyst **S1** with **S2**, enabling an effective transition-metal-free photocatalytic defluoroborylation protocol (entry 7). Control experiments demonstrated that light, photocatalyst, thiol and base were all essential for efficient defluoroborylation (entries 8-10).


With the optimal photo-induced defluoroborylation conditions in hand, we explored the substrate scope using the Ir photocatalyst-based protocol (Condition A) or the transition-metal-free protocol (Condition B) (Scheme 2). A series of pentafluoroarene derivatives underwent regioselective defluoroborylation efficiently (**1b-1l**). Notably, the transition-metal-free protocol (Condition B) afforded products (**3b-3e**) in yields comparable to those obtained

RESEARCH ARTICLE

with the Ir protocol (Condition A), demonstrating a practical strategy to prepare fluorinated organic boranes in a transition-metal-free manner. Functionalities such as a trifluoromethyl group (**3e**), ketone (**3g**), ester (**3h**), boronic ester (**3i**), pyridine (**3j**), and phosphine (**3k**, **3l**) were well-tolerated. Diborylation could be selectively achieved in good yields using 2.4 equivalents of NHC-borane (**3d'**, **3l'**). 1,2,3,4-Tetrafluorobenzene was defluoroborylated in excellent regioselectivity to deliver product **3m** in 85% yield. Tetrafluorobenzenes bearing functional groups such as aryl bromides and benzyl ethers also participated in the defluoroborylation to afford **3n** and **3o** in 45% and 53% yield respectively. Tetrafluorobenzenes containing an amine substituent underwent defluoroborylation selectively at the *ortho*-position to the amino group (**3p-3r**). The regioselectivity was confirmed by crystallographic analysis of the product **3q**.^[27] Defluoroborylation of ester-substituted tetrafluoroarenes occurred exclusively at the position *ortho* to the ester group (**3s**). Importantly, the substrate scope could be successfully expanded to trifluoroarenes and even difluoroarenes substituted with electron-deficient functionalities such as ketone (**3u**), ester (**3v-3x**), and nitrile (**3z** and **3ab**) in a highly selective fashion. Surprisingly, hydroborylation occurred to produce dearomatization products (**4t**, **4x**, **4y**, and **4aa**) in good yields as a single diastereomer if the *para*-position of an electron-withdrawing functional group (e.g. ester and nitrile) was unsubstituted in the polyfluoroarenes. The stereo-configuration of **4y** was identified by X-ray crystallographic analysis, illustrating a *trans* hydroborylation,^[28] and the structures of other dearomatization products were assigned by analogy. The scope for NHC-boranes was subsequently evaluated. Various NHC-boranes more sterically hindered than **2a** were good candidates, giving products **3ac-3ag** with 73% to 96% yields.

The most intriguing part of this method is probably the excellent regioselectivity for C–F bond activation achieved with polyfluoroarene substrates. Even though a general prediction mode cannot be established at the current stage, and may require sophisticated kinetic studies and computational calculations, the following trends can be observed and may be instructional with respect to the other radical-based C–F functionalization of polyfluoroarenes. 1) Polyfluoroarenes with electron-deficient substituents are generally more reactive and selective than those with electron-rich substituents (**1e** vs **1f**). 2) For pentafluoroarenes, the fluoride substitution took place at the most electronegative site of the radical anion intermediates derived from pentafluoroarenes (**1b-1l**).^[29] 3) For tetrafluoroarene (**1s**) and trifluoroarene (**1u-1x**, **1z**) with an electron-withdrawing substituent (e.g. ester, ketone, nitrile), the C–F activation reactivity trend was *ortho* > *para* > *meta*. 4) In the case of fluoroarenes containing an amine substituent (**1p-1r**), the reaction was directed by intramolecular hydrogen bonding and occurred at the *ortho* position of the amine substituent. The hydrogen bond might facilitate the fragmentation step and direct the regioselectivity to overcome the innate electronics of the fluorinated arene substrates.^[30] 5) Hydroborylation may occur to produce dearomatization products if a proton was located at the *para*-position of an electron-withdrawing group (**1t**, **1x**, **1y**, and **1aa**). Notably, in some cases, such as **3c** and **3j**, the regioselectivity is different from that obtained with the transition-metal-catalyzed processes where *ortho* defluoroborylation was

observed,^[9c,10b,17] providing an orthogonal strategy for regioselective defluoroborylation of polyfluoroarenes.




Scheme 3. Photo-mediated defluoroborylation of *gem*-difluoroalkenes.^[a] [a] Isolated yields. *E/Z* ratios were determined by the analysis of ¹H NMR spectra. DIPEA = *N,N*-diisopropylethylamine.

We subsequently attempted to extend this methodology to *gem*-difluoroalkenes, which are readily available building blocks.^[31] The optimal conditions for defluoroborylation of *gem*-difluoroalkenes were defined using a combination of $[\text{Ir}(\text{dF}(\text{CF}_3)\text{ppy})_2(5,5'\text{-dFbpy})]\text{PF}_6$,^[32] thiol (**S2**) and DIPEA in THF at room temperature under blue LED irradiation for 12 h. As illustrated in Scheme 3, symmetrical *gem*-difluoroalkenes participated well to afford mono-fluoroalkenylboranes in moderate to good yields (**6a-6d**). In the case of unsymmetrical *gem*-difluoroalkenes, mixtures of *E/Z* isomers (**6e-6p**) were obtained. Electron-withdrawing substituents such as chloride (**6g**) and bromide (**6h**), as well as electron-rich substituents including methyl sulfide (**6i**), phenol (**6j**), amine (**6k**), and methyl ether (**6l**) were well-tolerated. *meta*-Substituted aryl (**6l**, **6m**) and 1- or 2-naphthyl (**6n**, **6o**) *gem*-difluoroalkenes delivered the defluoroborylation products smoothly. A fluoroalkene containing a pyridyl substituent was also a good candidate for this transformation (**6p**).

gem-Difluoroallylboranes are important synthons for preparation of bioactive fluorinated compounds.^[13] We speculated that the HAT-induced nucleophilic NHC-boryl radical^[33] could add to an α -trifluoromethyl alkene, and a subsequent single electron reduction^[20] would promote an E1cB-type fluoride elimination^[34] to access such boranes. Indeed, after slight modifications of the photo-induced defluoroborylation protocol, this transformation was achieved efficiently under the transition-metal free conditions, with 4-CzIPN as the photocatalyst. The scope of trifluoromethylalkenes is illustrated in Scheme 4. The reactions proceeded smoothly with *para*-, *ortho*-, and *meta*-substituted α -trifluoromethylstyrenes to deliver *gem*-difluoroallylboranes in moderate to good yields (**8a-8j**). Styryl systems bearing a phenyl


RESEARCH ARTICLE

ether (**8c**), a trifluoromethyl substituent (**8d**), a bromide (**8e**), an alkene (**8f**), a fluoride (**8g** and **8i**), and a chloride (**8j**) were all effectively transformed into their corresponding *gem*-difluoroallylboranes. Naphthyl-substituted trifluoromethyl alkene also underwent successful defluoroborylation (**8k**).



Scheme 4. Photo-mediated defluoroborylation of trifluoromethylalkenes.^[a] [a] Isolated yields.

A series of control experiments were conducted to further elucidate the reaction mechanism (Scheme 5). No product was observed in the presence of 2,2,6,6-tetramethyl-1-piperidyl-oxyl (TEMPO), indicating a radical-based pathway (Scheme 5a). Stern-Volmer quenching studies illustrated that the thiol can quench the excited photocatalyst, whereas the polyfluoroarene, *gem*-difluoroalkene, and trifluoromethylalkene cannot (see the Supporting Information). Under similar conditions, the hydroboration of alkynes proceeded in high efficiency (Scheme 5b), which suggested the presence of boryl radical intermediates.^[21e] Hydrodefluorination could be achieved with the same photocatalyst (Scheme 5c), indicating that a single electron reduction of the fluoroarene to fluoroaryl radical anion might occur.^[25] The reaction of hexafluorobenzene with deuterated borane (**2a-D**) proceeded efficiently under the standard conditions to give deuterated product (**3a-D**) in 90% yield (Scheme 5d). The kinetic isotope effect (KIE) was determined to be 1.1 by measuring the initial rates of parallel reactions of **2a** and **2a-D** with hexafluorobenzene (see the Supporting Information), which suggested that the HAT of NHC-borane may not be involved in the rate-determining step. A radical chain-based mechanism was unlikely as the measured quantum yield was 0.43.^[35]




Scheme 5. Control experiments to elucidate reaction mechanisms.

Plausible mechanistic pathways for the defluoroborylation were proposed in light of all the experimental data. As illustrated in Scheme 6a, after photoexcitation of Ir(III) catalyst, the Ir(III)* ($E_{1/2}^{Ir(III)/Ir(II)} = +1.21$ V vs. saturated calomel electrode (SCE)) undergoes a single electron transfer with a thiol ($E_{1/2}^{S^{\cdot-}/S} = +0.77$ V vs. SCE) through a proton-coupled electron transfer to produce an Ir(II) species and thiyl radical **I** in the presence of base. A kinetically favored polarity-matched HAT^[22] between electrophilic thiyl radical **I** and NHC-BH₃ (BDE of B-H = 72.8 kcal/mol based on DFT calculations, slightly lower than the previous calculated value in the range of 74–80 kcal/mol^[21a,36]) gives NHC-boryl radical **II** and recovers the thiol catalyst (BDE of S-H = 76.1 kcal/mol). The defluoroborylation of polyfluoroarenes and *gem*-difluoroalkenes can be achieved by three possible pathways involving the NHC-boryl radical **II** and the fluoro substrate. 1) The generated Ir(II) species ($E_{1/2}^{Ir(II)/Ir(III)} = -1.37$ V vs. SCE) can be oxidized by fluoroarene ($E_{1/2}^{red} = -2.38$ V vs. SCE for **1a**) or *gem*-difluoroalkene ($E_{1/2}^{red} = -1.04$ V vs. SCE for **5a**)^[19a] to reproduce the Ir(III) catalyst and a radical anion **III**. This process is supported by the control experiment (Scheme 5c) and may be accelerated by the stabilization induced by the interaction between fluoroaryl radical anion **III** and the positively charged Ir(III) catalyst.^[19] Radical recombination occurs between the fluoroaryl radical anion **III** and NHC boryl radical **II** followed by the fluoride fragmentation to give the desired product **3** or **6**. 2) Fluoride extrusion from radical anion **III** may occur before the radical recombination. 3) Direct addition of NHC-boryl radical **II** to the fluoroarene or *gem*-difluoroalkene and subsequent single-electron reduction by the Ir(II) species, followed by fluoride extrusion delivers the final product. When a proton is located at the *para*-position of an electron-withdrawing group in the polyfluoroarene, protonation of intermediate **IV** occurs to generate

RESEARCH ARTICLE


the dearomatized hydroborylation product **4** (Scheme 6b), which may exclude the second possible pathway. Detailed coupling mechanistic investigations to distinguish between these pathways and gain a better appreciation of the source of selectivity are currently under progress in our laboratory. With trifluoromethylalkene substrates, the in situ formed NHC-boryl radical **II** will undergo radical addition to the alkene followed by a SET reduction by the reduced state of 4-CzIPN to generate a carbanion **VI**.^[20] The following E1cB-type fluoride elimination will provide the *gem*-difluoroallylborane product **8** (Scheme 6c).



Scheme 6. Proposed plausible mechanisms.

The synthetic value of this method was further demonstrated by diversification of the defluoroborylation product **3a** and *gem*-difluoroallylborane **8e** (Scheme 7). (Perfluoro-phenyl)boronic acid **9** was produced in high yield by a simple bromination/hydrolysis cascade, which has been widely utilized in cross-coupling reactions to deliver valuable polyfluoroarene compounds.^[8] Treatment of borane **3a** with phenyl bromide under palladium-catalyzed cross-coupling conditions led to mono- (**10a**) and diarylation (**10b**) of the NHC part of **3a** in 17% and 35% yield, respectively. The mono-defluoroborylation product **3a** could undergo further defluoroborylation to achieve 1,4-diborane product **11** in good yield. The reaction between **3a** and Selectfluor afforded borane difluoride **12** in high yield, which could act as a boronic acid equivalent in further diversifications.^[37] The borane difluoride **12** could also undergo further defluoroborylation, leading to the diborylated compound **13** with two distinguishable boron substituents. *gem*-Difluoroallylboranes **8e** can be directly transformed to *gem*-difluoroallyl alcohol **14** under oxidative

conditions or *gem*-difluoroallylboronates **15** with pinacol under acidic conditions.



Scheme 7. Further transformations of fluoroarylborene (**3a**) and *gem*-difluoroallylborane (**8e**)

Conclusion

In summary, we have developed a practical and selective defluoroborylation of polyfluoroarenes, *gem*-difluoroalkenes, and trifluoromethylalkenes in an atom- and redox-economic manner through photoredox and HAT-induced B–H activation. The selectivity is generally high, and can be orthogonal to transition-metal-catalyzed defluoroborylation. Fluorinated organoboranes with different degrees of fluorination and a wide range of functionalities were accessed by operationally simple and mild protocols, even in a transition-metal-free manner, and will likely find broad applications for the synthesis of value-added organofluorine compounds.

Experimental Section

Typical procedure for defluoroborylation of fluoroarenes: In a 10 mL Schlenk tube with a magnetic stir bar were placed **PC1** (1 mol%) or 4-CzIPN (2 mol%), NaOAc (8.4 mg, 0.1 mmol, 50 mol%), and NHC-BH₃ (**2**, 0.24 mmol, 1.2 equiv). Under nitrogen atmosphere, fluoroarenes (**1**, 0.2 mmol, 1 equiv), **S1** or **S2** (0.02 mmol, 10 mol%), THF (4 mL) were subsequently added. The resulting mixture was sealed and degassed via freeze-pump-thaw for three times. Then, the reaction was placed under blue LED (2-meter strips, 18 W) and irradiated for 12 h at room temperature. The solvent was removed under vacuum. Silica gel chromatography (eluent: hexane and ethyl acetate) of the crude product mixture afforded the desired compound.

Acknowledgements

We appreciate the insightful discussion from Prof. Zhang Xinhao (Peking University Shenzhen Graduate School) and Prof. Luo Sanzhong (Tsinghua University). We are grateful for the financial

RESEARCH ARTICLE

support provided by the Ministry of Education (MOE) of Singapore (MOE2017-T2-2-081), GSK-EDB (R-143-000-687-592), NUS (Suzhou) Research Institute, and National Natural Science Foundation of China (Grant No. 21702142, 21871205).

Keywords: polyfluoroarene • boryl radical • hydrogen atom transfer • photocatalysis • defluoroborylation

- [1] J. D. Weaver, S. Senaweera, *Tetrahedron* **2014**, 70, 7413-7428.
- [2] a) S. Purser, P. R. Moore, S. Swallow, V. Gouverneur, *Chem. Soc. Rev.* **2008**, 37, 320-330; b) Y. Zhou, J. Wang, Z. Gu, S. Wang, W. Zhu, J. L. Aceña, V. A. Soloshonok, K. Izawa, H. Liu, *Chem. Rev.* **2016**, 116, 422-518.
- [3] a) T. Mori, K. Ujihara, O. Matsumoto, K. Yanagi, N. Matsuo, *J. Fluorine Chem.* **2007**, 128, 1174-1181; b) P. Jeschke, *Pest Manag. Sci.* **2017**, 73, 1053-1066.
- [4] For selected studies, see: a) T. Ritter, M. W. Day, R. H. Grubbs, *J. Am. Chem. Soc.* **2006**, 128, 11768-11769; b) R. Shintani, N. Misawa, T. Tsuda, R. Iino, M. Fujii, K. Yamashita, K. Nozaki, *J. Am. Chem. Soc.* **2017**, 139, 3861-3867.
- [5] M. L. Tang, Z. Bao, *Chem. Mater.* **2011**, 23, 446-455.
- [6] a) A. D. Sun, J. A. Love, *Dalton Trans.* **2010**, 39, 10362-10374; b) T. Fujita, K. Fuchibe, J. Ichikawa, *Angew. Chem. Int. Ed.* **2019**, 58, 390-402; *Angew. Chem.* **2019**, 131, 396-408.
- [7] M. G. Campbell, T. Ritter, *Chem. Rev.* **2015**, 115, 612-633.
- [8] a) N. Miyaura, A. Suzuki, *Chem. Rev.* **1995**, 95, 2457-2483; b) T. Kinzel, Y. Zhang, S. L. Buchwald, *J. Am. Chem. Soc.* **2010**, 132, 14073-14075; c) S. Handa, Y. Wang, F. Gallou, B. H. Lipshutz, *Science* **2015**, 349, 1087-1091; d) L. Chen, D. R. Sanchez, B. Zhang, B. P. Carrow, *J. Am. Chem. Soc.* **2017**, 139, 12418-12421; e) D. Bulfield, S. M. Huber, *J. Org. Chem.* **2017**, 82, 13188-13203; f) L. Chen, H. Francis, B. P. Carrow, *ACS Catal.* **2018**, 8, 2989-2994.
- [9] a) T. Braun, M. Ahijado Salomon, K. Altenhöner, M. Teltewskoi, S. Hinze, *Angew. Chem. Int. Ed.* **2009**, 48, 1818-1822; *Angew. Chem.* **2009**, 121, 1850-1854; b) M. Teltewskoi, J. A. Panetier, S. A. Macgregor, T. Braun, *Angew. Chem. Int. Ed.* **2010**, 49, 3947-3951; *Angew. Chem.* **2010**, 122, 4039-4043; c) W.-H. Guo, Q.-Q. Min, J.-W. Gu, X. Zhang, *Angew. Chem. Int. Ed.* **2015**, 54, 9075-9078; *Angew. Chem.* **2015**, 127, 9203-9206.
- [10] a) X.-W. Liu, J. Echavarren, C. Zarate, R. Martin, *J. Am. Chem. Soc.* **2015**, 137, 12470-12473; b) J. Zhou, M. W. Kuntze-Fechner, R. Bertermann, U. S. D. Paul, J. H. J. Berthel, A. Friedrich, Z. Du, T. B. Marder, U. Radius, *J. Am. Chem. Soc.* **2016**, 138, 5250-5253.
- [11] a) T. Niwa, H. Ochiai, Y. Watanabe, T. Hosoya, *J. Am. Chem. Soc.* **2015**, 137, 14313-14318; b) H. Sakaguchi, Y. Uetake, M. Ohashi, T. Niwa, S. Ogoshi, T. Hosoya, *J. Am. Chem. Soc.* **2017**, 139, 12855; c) P. Gao, C. Yuan, Y. Zhao, Z. Shi, *Chem* **2018**, 4, 2201-2211; (d) R. Kojima, S. Akiyama, H. Ito, *Angew. Chem. Int. Ed.* **2018**, 57, 7196-7199; *Angew. Chem.* **2018**, 130, 7314-7317.
- [12] S. Lim, D. Song, S. Jeon, Y. Kim, H. Kim, S. Lee, H. Cho, B. C. Lee, S. E. Kim, K. Kim, E. Lee, *Org. Lett.* **2018**, 20, 7249-7252.
- [13] Y. Liu, Y. Zhou, Y. Zhao, J. Qu, *Org. Lett.* **2017**, 19, 946-949.
- [14] a) J. M. R. Narayanam, C. R. J. Stephenson, *Chem. Soc. Rev.* **2011**, 40, 102-113; b) J. Xuan, W.-J. Xiao, *Angew. Chem. Int. Ed.* **2012**, 51, 6828-6838; *Angew. Chem.* **2012**, 124, 6934-6944; c) C. K. Prier, D. A. Rankic, D. W. C. MacMillan, *Chem. Rev.* **2013**, 113, 5322-5363; d) K. L. Skubi, T. R. Blum, T. P. Yoon, *Chem. Rev.* **2016**, 116, 10035-10074; e) D. Ravelli, S. Protti, M. Fagnoni, *Chem. Rev.* **2016**, 116, 9850-9913; f) N. A. Romero, D. A. Nicewicz, *Chem. Rev.* **2016**, 116, 10075-10166; g) L. Marzo, S. K. Pagire, O. Reiser, B. König, *Angew. Chem. Int. Ed.* **2018**, 57, 10034-10072; *Angew. Chem.* **2018**, 130, 10188-10228.
- [15] K. Chen, M. S. Cheung, Z. Lin, P. Li, *Org. Chem. Front.* **2016**, 3, 875-879.
- [16] A. M. Mfuh, J. D. Doyle, B. Chhetri, H. D. Arman, O. V. Larionov, *J. Am. Chem. Soc.* **2016**, 138, 2985-2988.
- [17] Y.-M. Tian, X.-N. Guo, M. W. Kuntze-Fechner, I. Krummenacher, H. Braunschweig, U. Radius, A. Steffen, T. B. Marder, *J. Am. Chem. Soc.* **2018**, 140, 17612-17623.
- [18] a) A. Singh, J. J. Kubik, J. D. Weaver, *Chem. Sci.* **2015**, 6, 7206-7212; b) A. Singh, C. J. Fennell, J. D. Weaver, *Chem. Sci.* **2016**, 7, 6796-6802; c) S. Senaweera, J. D. Weaver, *J. Am. Chem. Soc.* **2016**, 138, 2520-2523; d) S. Priya, J. D. Weaver, *J. Am. Chem. Soc.* **2018**, 140, 16020-16025.
- [19] a) J. Xie, J. Yu, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Angew. Chem. Int. Ed.* **2016**, 55, 9416-9421; *Angew. Chem.* **2016**, 128, 9563-9568; b) J. Xie, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Angew. Chem. Int. Ed.* **2017**, 56, 7266-7270; *Angew. Chem.* **2017**, 129, 7372-7376.
- [20] S. B. Lang, R. J. Wiles, C. B. Kelly, G. A. Molander, *Angew. Chem. Int. Ed.* **2017**, 56, 15073-15077; *Angew. Chem.* **2017**, 129, 15269-15273.
- [21] a) S.-H. Ueng, M. Makhlof Brahmi, É. Derat, L. Fensterbank, E. Lacôte, M. Malacria, D. P. Curran, *J. Am. Chem. Soc.* **2008**, 130, 10082-10083; b) G. Duret, R. Quinlan, P. Bissleret, N. Blanchard, *Chem. Sci.* **2015**, 6, 5366-5382; c) S.-C. Ren, F.-L. Zhang, J. Qi, Y.-S. Huang, A.-Q. Xu, H.-Y. Yan, Y.-F. Wang, *J. Am. Chem. Soc.* **2017**, 139, 6050-6053; d) Y. Su, R. Kinjo, *Coord. Chem. Rev.* **2017**, 352, 346-378; e) M. Shimoi, T. Watanabe, K. Maeda, D. P. Curran, T. Taniguchi, *Angew. Chem. Int. Ed.* **2018**, 57, 9485-9490; *Angew. Chem.* **2018**, 130, 9629-9634; f) W. Dai, T. R. McFadden, D. P. Curran, H. A. Früchtl, J. C. Walton, *J. Am. Chem. Soc.* **2018**, 140, 15868-15875; g) S.-C. Ren, F.-L. Zhang, A.-Q. Xu, Y. Yang, M. Zheng, X. Zhou, Y. Fu, Y.-F. Wang, *Nat. Commun.* **2019**, 10, 1934; h) A.-Q. Xu, F.-L. Zhang, T. Ye, Z.-X. Yu, Y.-F. Wang, *CCS Chem.* **2019**, 1, 504-512.
- [22] a) M.-A. Tehfe, M. Makhlof Brahmi, J.-P. Fouassier, D. P. Curran, M. Malacria, L. Fensterbank, E. Lacôte, J. Lalevée, *Macromolecules* **2010**, 43, 2261-2267; b) X. Pan, E. Lacôte, J. Lalevée, D. P. Curran, *J. Am. Chem. Soc.* **2012**, 134, 5669-5674; c) S. Telitel, S. Schweizer, F. Morlet-Savary, B. Graff, T. Tschamber, N. Blanchard, J. P. Fouassier, M. Lelli, E. Lacôte, J. Lalevée, *Macromolecules* **2013**, 46, 43-48; d) T. Watanabe, D. Hirose, D. P. Curran, T. Taniguchi, *Chem. Eur. J.* **2017**, 23, 5404-5409.
- [23] a) R. Zhou, Y. Y. Goh, H. W. Liu, H. R. Tao, L. H. Li, J. Wu, *Angew. Chem. Int. Ed.* **2017**, 56, 16621-16625; *Angew. Chem.* **2017**, 129, 16848-16852; b) H. P. Deng, X. Z. Fan, Z. H. Chen, Q. H. Xu, J. Wu, *J. Am. Chem. Soc.* **2017**, 139, 13579-13584; c) J. Hou, A. Ee, H. Cao, H.-W. Ong, J.-H. Xu, J. Wu, *Angew. Chem. Int. Ed.* **2018**, 57, 17220-17224; *Angew. Chem.* **2018**, 130, 17466-17470; d) X. Z. Fan, J. W. Rong, H. L. Wu, Q. Zhou, H. P. Deng, J. Da Tan, C. W. Xue, L. Z. Wu, H. R. Tao, J. Wu, *Angew. Chem. Int. Ed.* **2018**, 57, 8514-8518; *Angew. Chem.* **2018**, 130, 8650-8654; e) H. P. Deng, Q. Zhou, J. Wu, *Angew. Chem. Int. Ed.* **2018**, 57, 12661-12665; *Angew. Chem.* **2018**, 130, 12843-12847; f) J. Li, Y. Luo, H. W. Cheo, Y. Lan, J. Wu, *Chem* **2019**, 5, 192-203.
- [24] N. Zhou, X.-A. Yuan, Y. Zhao, J. Xie, C. Zhu, *Angew. Chem. Int. Ed.* **2018**, 57, 3990-3994; *Angew. Chem.* **2018**, 130, 4054-4058.
- [25] S. M. Senaweera, A. Singh, J. D. Weaver, *J. Am. Chem. Soc.* **2014**, 136, 3002-3005.
- [26] J. Luo, J. Zhang, *ACS Catal.* **2016**, 6, 873-877.
- [27] CCDC 1894872 for **3q** contains the supplementary crystallographic data.
- [28] CCDC 1937072 for **4y** contains the supplementary crystallographic data. A kinetically controlled radical-based *trans*-hydroboration of alkynes with *N*-heterocyclic carbene boranes has been reported, see Ref.21e.
- [29] See the Supporting Information for details.
- [30] M. B. Khaled, R. K. Mokadem, J. D. Weaver, *J. Am. Chem. Soc.* **2017**, 139, 13092-13101.
- [31] M. Hu, Z. He, B. Gao, L. Li, C. Ni, J. Hu, *J. Am. Chem. Soc.* **2013**, 135, 17302-17305.
- [32] G. J. Choi, Q. Zhu, D. C. Miller, C. J. Gu, R. R. Knowles, *Nature* **2016**, 539, 268-271.
- [33] C. Wu, X. Hou, Y. Zheng, P. Li, D. Lu, *J. Org. Chem.* **2017**, 82, 2898-2905.


RESEARCH ARTICLE

- [34] S. Alunni, F. De Angelis, L. Ottavi, M. Papavasileiou, F. Tarantelli, *J. Am. Chem. Soc.* **2005**, *127*, 15151-15160.
- [35] M. A. Cismesia, T. P. Yoon, *Chem. Sci.* **2015**, *6*, 5426-5434.
- [36] J. Hioe, A. Karton, J. M. L. Martin, H. Zipse, *Chem. Eur. J.* **2010**, *16*, 6861-6865.
- [37] S. Nerkar, D. P. Curran, *Org. Lett.* **2015**, *17*, 3394-3397.

RESEARCH ARTICLE

Entry for the Table of Contents

RESEARCH ARTICLE



Wengang Xu, Heming Jiang, Jing Leng,
Han-Wee Ong, and Jie Wu*

Page No. – Page No.

**Visible-Light-Induced Selective
Defluoroborylation of
Polyfluoroarenes, *gem*-
Difluoroalkenes, and
Trifluoromethylalkenes**

A practical and selective defluoroborylation of polyfluoroarenes, *gem*-difluoroalkenes, and trifluoromethylalkenes was developed through photoredox and hydrogen atom transfer-induced B–H activation to synthesize fluorinated organoboranes as the building blocks for value-added organofluorine compounds.