

Synthesis and ^{31}P and ^{195}Pt NMR characterisation of the first binuclear platinum(I) complex $[\text{Pt}_2\text{Cl}_2(\mu\text{-dppm})_2(\mu\text{-Bu}^t\text{CP})]$ containing a μ -parallel ligated phospho-alkyne

Saud I. Al-Resayes* and John F. Nixon**

School of Chemistry and Molecular Sciences, University of Sussex, Brighton BN1 9QJ, Sussex (UK)

(Received February 5, 1993)


Abstract

The first example of a μ -parallel ligated phospho-alkyne is reported in the complex $[\text{Pt}_2\text{Cl}_2(\mu\text{-dppm})_2(\mu\text{-Bu}^t\text{CP})]$ whose structure has been elucidated by ^{31}P and ^{195}Pt NMR studies.

Introduction


In recent years extensive work has been carried out on a variety of binuclear transition metal complexes of dppm [1]. The greatest interest has focussed on studies of $[\text{MM}'\text{X}_2(\mu\text{-dppm})_2]$ complexes $\text{M}=\text{M}'=\text{Pd}$, Pt or Rh ; $\text{M}=\text{Pd}$, $\text{M}'=\text{Pt}$ [2–14]. In these complexes the addition of small molecules (e.g. CH_2 , CO , CNR , C_2R_2 , CS_2 , H_2 , S and SO_2) across the metal–metal bond have been thoroughly studied to give the so-called ‘A-frame’ complexes [15–33].

Of particular interest in connection with the work described in this paper concerns the interaction of these dinuclear complexes with alkynes, since the resulting products adopt a *cis*-dimetallated olefin μ -parallel bonding mode shown below as evidenced by NMR studies, single crystal X-ray diffraction studies [15, 22, 25] and rationalised by MO calculations [32] rather than the more common perpendicular bridging mode found in other systems.



Phospha-alkynes, $\text{RC}\equiv\text{P}$, are known to behave like alkynes in their coordination complexes of transition

metals [34], but to date interactions with dimetallic centres have resulted in μ -perpendicular complexes of the type [35–38]



It was therefore of interest to investigate analogous reactions of $[\text{M}_2\text{Cl}_2(\mu\text{-dppm})_2]$ ($\text{M}=\text{Pt}$) with the phospho-alkyne Bu^tCP to see if it also undergoes a μ -parallel bonding mode in accord with the Hoffmann's predictions based on MO calculations [32].

Results and discussion

Treatment of a solution of $[\text{Pt}_2\text{Cl}_2(\mu\text{-dppm})_2]$ in dichloromethane with an excess of Bu^tCP at ambient temperature afforded a high yield of the yellow microcrystalline complex $[\text{Pt}_2\text{Cl}_2(\mu\text{-dppm})_2(\text{Bu}^t\text{CP})]$ (**1**) which was characterised by elemental analysis, IR spectroscopy and its structure suggested by $^{31}\text{P}\{^1\text{H}\}$ NMR and ^{195}Pt NMR spectroscopy (*vide infra*).

The $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of **1** recorded at 32.4 and 145.8 MHz are shown in Figs. 1 and 2 and coupling constant data are listed in Table 1. Analysis of the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum recorded at 32.4 MHz was complicated by the overlapping of the low field ^{195}Pt satellites of the dppm resonance with that of the $\mu\text{-Bu}^t\text{CP}$ ligand, but the 145.8 MHz spectrum proved more amenable and observed and simulated resonances for P^{A} and P^{B} are shown using data listed in Table 1.

*Present address: Chemistry Department, King Saud University, Riyadh, Saudi Arabia.

**Author to whom correspondence should be addressed.




Fig. 1. $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of $[\text{Pt}_2\text{Cl}_2(\mu\text{-dppm})_2(\text{Bu}'\text{CP})]$ (**1**) at 32.4 MHz.





Fig. 2. (a) $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of $[\text{Pt}_2\text{Cl}_2(\mu\text{-dppm})_2(\text{Bu}'\text{CP})]$ (**1**) at 145.8 MHz (b) Simulated (top) and experimental $^{31}\text{P}\{^1\text{H}\}$ NMR spectra for the A and B spectra of the central $[\text{AA}'\text{BB}'\text{M}]$ resonance.



Data for related μ -alkyne diplatinum complexes are listed for comparison. No change was observed in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **1** over the temperature range -60 to 30°C .

The $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **1** results from the three non-equivalent phosphorus nuclei (P^{A} , P^{B} and P^{M}), and therefore consists of three super-imposed spectra which arise from the structures shown below, which reflect the fact that ^{195}Pt ($I = 1/2$) occurs in 33.8%

TABLE 1. $^{31}\text{P}\{^1\text{H}\}$ NMR data for **1** and related diplatinum complexes

Compound	$\delta\text{P}^{\text{A}}$ ^a	$\delta\text{P}^{\text{B}}$ ^a	$^1J(\text{Pt}^{\text{X}}\text{P}^{\text{A}})^{\text{b}}$ $=^1J(\text{Pt}^{\text{X}}\text{P}^{\text{A}})$	$^3J(\text{Pt}^{\text{X}}\text{P}^{\text{B}})^{\text{b}}$ $=^3J(\text{Pt}^{\text{X}}\text{P}^{\text{B}})$	$^1J(\text{Pt}^{\text{X}}\text{P}^{\text{B}})^{\text{b}}$ $=^1J(\text{Pt}^{\text{X}}\text{P}^{\text{B}})$	$^3J(\text{Pt}^{\text{X}}\text{P}^{\text{A}})^{\text{b}}$ $=^3J(\text{Pt}^{\text{X}}\text{P}^{\text{A}})$	$^2J(\text{P}^{\text{A}}\text{P}^{\text{B}})^{\text{b}}$ $=^2J(\text{P}^{\text{A}}\text{P}^{\text{B}})$	Ref.
 (1)	+3.2	-0.4 ^c +332 ^d	3605 ^e 420 ^f	+122.1	3557	+180.6	22.1 ^g 22.5 ^h 6.8 ⁱ 14.8 ^j	this work
 R ¹ =R ² =CF ₃ R ³ =H R ⁴ =Me ₂ CO	+11.88	-0.15	3110	90	3055	100	20	24
 R ¹ =R ² =CF ₃ R ³ =H R ⁴ =Cl	+11.3	+0.06	3232	50	3193	70	25	24
 R ¹ =CF ₃ R ² =H R ³ =R ⁴ =CCCCF ₃	+0.9	-0.206	3020	100	3250	100	17	24


^aIn ppm relative to 85% H₃PO₄. ^bIn Hz. ^c $\delta\text{P}^{\text{B}}$. ^d $\delta\text{P}^{\text{M}}$. ^e $^1J(\text{Pt}^{\text{X}}\text{P}^{\text{A}})=^1J(\text{Pt}^{\text{X}}\text{P}^{\text{A}})$. ^f $^1J(\text{Pt}^{\text{X}}\text{P}^{\text{B}})$. ^g $^2J(\text{P}^{\text{A}}\text{P}^{\text{B}})=^2J(\text{P}^{\text{A}}\text{P}^{\text{B}})$. ^h $^2J(\text{P}^{\text{A}}\text{P}^{\text{M}})=^2J(\text{P}^{\text{A}}\text{P}^{\text{M}})$. ⁱ $^4J(\text{P}^{\text{A}}\text{P}^{\text{B}})=^4J(\text{P}^{\text{B}}\text{P}^{\text{A}})$. ^j $^3J(\text{P}^{\text{B}}\text{P}^{\text{M}})=^3J(\text{P}^{\text{B}}\text{P}^{\text{M}})$. P⁺P⁻=dppm.

abundance. The three spin systems are thus [AA'BB'M] for (i), [AA'BB'MX] for (ii) and [AA'BB'MXX'] for (iii), (A=B=M, ^{31}P $I=1/2$ 100%; X= ^{195}Pt $I=1/2$ 33.8%), from which the coupling constants given in Table 1 were obtained.

The $^{195}\text{Pt}\{^1\text{H}\}$ NMR spectrum (Fig. 3) though much poorer resolved than the $^{31}\text{P}\{^1\text{H}\}$ NMR spectra shows the expected two non-equivalent Pt nuclei both appearing as 'triplets' but only the latter showing the extra $J(\text{PtP}^{\text{M}})$ coupling (420 Hz).

The NMR data thus confirm the presence of an η^2 -ligated phospho-alkyne adopting the same μ -parallel ligating behaviour as in alkyne diplatinum complexes. This bonding type for the phospho-alkyne is in fact peculiar to the dinuclear platinum systems since in related work affording the dirhodium complex $[\text{Rh}_2\text{Cl}_2(\mu\text{-dppm})_2(\text{Bu}^i\text{CP})]$ the phospho-alkyne adopts a different ligating mode in which the two metal atoms

are chemically identical both in solution and in the solid state as shown below [39].



Experimental

All manipulations were carried out under an atmosphere of dry dinitrogen or argon, and all reactions were handled by high-vacuum and Schlenk tube techniques. Solvents were dried by standard methods and freshly distilled under dinitrogen before use unless otherwise stated.

IR spectra were recorded on a Perkin-Elmer 457 spectrometer (4000–250 cm⁻¹) or Perkin-Elmer 1430 spectrometer (4000–200 cm⁻¹).

Elemental analyses were carried out by Mrs A.G. Olney of this School.

$^{31}\text{P}\{^1\text{H}\}$ NMR spectra were obtained using either Bruker WP80 (32.4 MHz) or Bruker WM360 (145.8 MHz) spectrometers both operating in the FT mode. Trimethyl phosphite P(OMe)₃ (TMP) was used as an external standard and chemical shifts were converted relative to 85% H₃PO₄ by using the relationship $\delta\text{p}(\text{P}(\text{OMe})_3) = +141$ ppm. ^{195}Pt NMR spectra were recorded on a WM360 spectrometer (77.1 MHz) and referenced to K₂PtCl₄.




Fig. 3. ^{195}Pt NMR spectrum of $[\text{Pt}_2\text{Cl}_2(\mu\text{-dppm})_2(\text{Bu}^i\text{CP})]$.

Preparation of $[Pt_2Cl_2(\mu-dppm)_2]$

$[Pt_2Cl_2(\mu-dppm)_2]$ was prepared by a variation of synthetic routes reported elsewhere [40, 41] involving either (a) or (b).

(a) Reaction of $[Pt(PPh_3)_4]$ with $[PtCl_2(SEt_2)_2]$ and dppm

To a solution of $[Pt(PPh_3)_4]$ (0.920 g, 0.739 mmol) (slight excess) in dry benzene (30 cm³) was added dppm (0.517 g, 1.344 mmol) under dinitrogen. $[PtCl_2(SEt_2)_2]$ (0.30 g, 0.672 mmol) was added and the mixture was refluxed under dinitrogen for 1 h. The mixture was left to cool to room temperature and the yellow product was filtered off, washed with benzene (10 cm³), and dried *in vacuo* for 3 h to give the yellow microcrystalline complex $[Pt_2Cl_2(\mu-dppm)_2]$ (0.560 g, 67%). *Anal.* Found: C, 48.2; H, 3.4. *Calc.* for $C_{50}H_{44}P_4Cl_2Pt_2$: C, 48.83; H, 3.61%. The $^31P\{^1H\}$ NMR and IR spectra are similar to those described in the literature.

(b) Reaction of $[Pt(PPh_3)_4]$ with $[PtCl_2(COD)]$ and dppm

$[Pt(PPh_3)_4]$ (1.254 g, 1.008 mmol) and dppm (0.704 g, 1.832 mmol) in dry benzene (30 cm³) were refluxed with $[PtCl_2(COD)]$ (0.343 g, 0.916 mmol) to afford $[Pt_2Cl_2(\mu-dppm)_2]$ (0.45 g, 40%), identical with the above sample.

Preparation of $[Pt_2Cl_2(\mu-dppm)_2(Bu'CP)]$ (1)

To a solution of $[Pt_2Cl_2(\mu-dppm)_2]$ (0.40 g, 0.325 mmol) in dry dichloromethane (15 cm³) was added Bu'CP (0.050 g, 0.50 mmol) in dichloromethane (5 cm³). The mixture was stirred for 1 h at room temperature during which time its colour changed from yellow to orange. The volatile materials were removed under reduced pressure and the yellow solid washed several times with a mixture of dichloromethane and petroleum ether (60–80 °C) (1:1). The product was dried *in vacuo* to give the yellow microcrystalline complex $[1,2\text{-dichloro-}\mu\text{-bis(diphenylphosphino)-methane-}P,P']\{\mu\text{-2,2-dimethylpropylidynephosphine-(}P,C)\}$ diplatinum(I) (Pt^1 , Pt^2) (0.37 g, 85%). *Anal.* Found: C, 48.95; H, 4.0. *Calc.* for $C_{55}H_{53}P_5Cl_2Pt_2$: C, 49.67; H, 4.02%. IR spectrum (KBr): 3050w, 2958m, 2905w, 2830w, 1573w, 1560m, 1483s, 1455sh, 1450vw, 14w, vs, 1382w, 1358m, 1332w, 1310w, 1261vs, 1223vw, 1192m, 1160vw, 1146sh, 1101vs, 1025vs, br, 1008sh, 990sh, 830w, br, 803vs, 790sh, 775s, 756sh, 742vs, 722s, 700vs, 678sh, 625w, 600sh, 503vs, 473s, 423w, 40m, 387s, 362w, 347w cm⁻¹.

Acknowledgements

We thank the Saudi Arabian government for financial support (to S.I.AI-R.) and Dr Tony Avent for help with the spectroscopic analysis.

References

- 1 R.J. Puddephatt, *Chem. Soc. Rev.*, 12 (1983) 99
- 2 A.L. Balch, *J. Am. Chem. Soc.*, 98 (1976) 8049
- 3 A.L. Balch and B. Tulyathan, *Inorg. Chem.*, 16 (1977) 2840.
- 4 A.L. Balch, J.W. Labadie and G. Delker, *Inorg. Chem.*, 18 (1979) 1224.
- 5 J.T. Mague and S.H. De Vries, *Inorg. Chem.*, 19 (1980) 3743.
- 6 R. Colton, M.J. McCormick, R.G. Holloway and B.R. Penfold, *J. Chem. Soc., Chem. Commun.*, (1976) 485.
- 7 M.P. Brown, R.J. Puddephatt and M. Rashidi, *Inorg. Chim. Acta*, 19 (1976) L33
- 8 M.P. Brown, R.J. Puddephatt, M. Rashidi and K.R. Seddon, *J. Chem. Soc., Dalton Trans.*, (1977) 951.
- 9 Lj. Manojlovic-Muir, K.W. Muir and T. Solomun, *Acta Crystallogr., Sect. B*, 35 (1979) 1237.
- 10 R. Colton, M.J. McCormick and C.D. Pannan, *Aust. J. Chem.*, 31 (1978) 1425.
- 11 P.G. Pringle and B.L. Shaw, *J. Chem. Soc., Dalton Trans.*, (1983) 889.
- 12 J.R. Batson, M.C. Grossel, R.P. Moulding and K.R. Seddon, *J. Organomet. Chem.*, 304 (1986) 391
- 13 R. Colton, R.H. Farthing and M.J. McCormick, *Aust. J. Chem.*, 26 (1973) 2607.
- 14 L.S. Benner and A.L. Balch, *J. Am. Chem. Soc.*, 100 (1978) 6094.
- 15 M. Cowie and T.G. Southern, *J. Organomet. Chem.*, 193 (1980) C46.
- 16 B.P. Brown, J.R. Fisher, S.J. Franklin and R.J. Puddephatt, *J. Chem. Soc., Chem. Commun.*, (1978) 749.
- 17 M.P. Brown, R.J. Puddephatt, M. Rashidi and K.R. Seddon, *J. Chem. Soc., Dalton Trans.*, (1978) 1540.
- 18 A.L. Balch, L.S. Benner and M.M. Olmstead, *Inorg. Chem.*, 18 (1979) 2996.
- 19 M.P. Brown, J.R. Fisher, R.J. Puddephatt and K.R. Seddon, *Inorg. Chem.*, 18 (1979) 2808.
- 20 M.P. Brown, S.J. Cooper, A.A. Frew, Lj. Manojlovic-Muir, K.W. Muir, R.J. Puddephatt and M.A. Thomson, *J. Chem. Soc., Dalton Trans.*, (1982) 299.
- 21 R. Colton, M.J. McCormick and C.D. Pannan, *J. Chem. Soc., Chem. Commun.*, (1977) 823.
- 22 A.L. Balch, C.-L. Lee, C.H. Lindsay and M.M. Olmstead, *J. Organomet. Chem.*, 177 (1979) C22.
- 23 C.-L. Lee, C.T. Hunt and A.L. Balch, *Inorg. Chem.*, 20 (1981) 2498.
- 24 R.J. Puddephatt and M.A. Thomson, *Inorg. Chem.*, 21 (1982) 725.
- 25 M. Cowie and R.S. Dickson, *Inorg. Chem.*, 20 (1981) 2682
- 26 M.P. Brown, R.J. Puddephatt, M. Rashidi and K.R. Seddon, *J. Chem. Soc., Dalton Trans.*, (1978) 516.
- 27 T.S. Cameron, P.A. Gardner and K.R. Grundy, *J. Organomet. Chem.*, 212 (1981) C19.
- 28 A.D. Rattray and D. Sutton, *Inorg. Chim. Acta*, 27 (1978) L85.
- 29 L.S. Benner, M.M. Olmstead, H. Hope and A.L. Balch, *J. Organomet. Chem.*, 153 (1978) C31.
- 30 M.M. Olmstead, H. Hope, L.S. Benner and A.L. Balch, *J. Am. Chem. Soc.*, 99 (1977) 5502.
- 31 M.P. Brown, A.N. Keigh, Lj. Manojlovic-Muir, K.W. Muir, R.J. Puddephatt and K.R. Seddon, *Inorg. Chim. Acta*, 34 (1979) L223.
- 32 D.M. Hoffman and R. Hoffman, *Inorg. Chem.*, 20 (1981) 3543.
- 33 M. Cowie, R.S. Dickson and B.W. Hames, *Organometallics*, 3 (1984) 1879

- 34 J.F. Nixon, *Chem. Rev.*, **88** (1988) 1327.
- 35 J.C.T.R. Burchett St. Laurent, P.B. Hitchcock, H.W. Kroto, M.F. Meidine and J.F. Nixon, *J Organomet Chem.*, **238** (1982) C82.
- 36 R. Bartsch, P.B Hitchcock, M.F. Meidine and J.F. Nixon, *J Organomet Chem*, **266** (1984) C41.
- 37 R. Bartsch, J.F. Nixon and N. Sarjudeen, *J Organomet. Chem*, **294** (1985) 267.
- 38 M.F. Meidine, C.J. Meir, S. Morton and J.F. Nixon, *J Organomet. Chem*, **297** (1985) 255.
- 39 S.I. Al-Resayes, C Jones, M.J. Maah and J.F. Nixon, *J. Organomet Chem*, in press.
- 40 M.P. Brown, R.J. Puddephatt, M. Rashidi and K.R Seddon, *J Chem Soc., Dalton Trans*, (1977) 951.
- 41 P.G. Pringle and B.L. Shaw, *J Chem. Soc , Dalton Trans*, (1983) 889