

CHEM MED CHEM

CHEMISTRY ENABLING DRUG DISCOVERY

Accepted Article

Title: Resolving Binding Events on the Multifunctional Human Serum Albumin

Authors: Stefan Petry, Lea Wenskowsky, Michael Wagner, Johannes Reusch, Herman Schreuder, Hans Matter, and Till Opatz

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: *ChemMedChem* 10.1002/cmdc.202000069

Link to VoR: <http://dx.doi.org/10.1002/cmdc.202000069>

WILEY-VCH

www.chemmedchem.org

A Journal of



Resolving Binding Events on the Multifunctional Human Serum Albumin

Lea Wenskowsky,^[b] Michael Wagner,^[a] Johannes Reusch,^[c] Herman Schreuder,^[a] Hans Matter,^[a] Till Opatz,^{[b]*} and Stefan Matthias Petry^{*[a]}

Abstract: Physiological processes rely on initial recognition events between cellular components and other molecules or modalities. Biomolecules can have multiple sites or mode of interaction with other molecular entities, so that a resolution of the individual binding events in terms of spatial localization as well as association and dissociation kinetics is required for a meaningful description. Here we describe a trichromatic fluorescent binding- and displacement assay for simultaneous monitoring and localization of three individual binding sites in the important transporter and binding protein human serum albumin. Independent investigations of binding events by X-ray crystallography and time-resolved dynamics measurements (switchSENSE technology) confirm the validity of the assay, the localization of binding sites and furthermore reveal conformational changes associated with ligand binding. The described assay system allows for the detailed characterization of albumin-binding drugs and is therefore well-suited for prediction of drug-drug and drug-food interactions. Moreover, conformational changes, usually associated with binding events, can also be analyzed.

Human serum albumin (HSA) is an abundant plasma protein, which binds and transports hydrophobic endogenous ligands like fatty acids. It also interacts with a wide range of drug and drug-like molecules.^[1] Several X-ray crystal structures of ligands in complex with HSA revealed seven individual fatty acid (FA) binding sites in this biomolecule.^[2]

Earlier, we addressed the specificity of these HSA binding sites with respect to ligand binding. This led to the identification of a single high affinity fatty acid binding site by introducing and applying the novel molecular probe NDB-FA.^[3]

Under physiological conditions, an albumin molecule accommodates only 0.1–2 molecules of fatty acids.^[4] In contrast to standard crystallization conditions using an excess of free fatty acids to stabilize the albumin conformation, we obtained an X-ray crystal structure (PDB 6ezq, resolution 2.4 Å) under conditions close to those employed for K_D determination for albumin complexes (NBD-FA:HSA 6:1) in our previous study.^[3]

To obtain more insight into the gradual occupancy of binding

sites in albumin as well as their relevance to physiology and pharmacokinetics, we then sought to develop a rapid assay system with optical readout. This system should allow the simultaneous monitoring of different albumin binding sites, when interacting with ligands. Therefore, it provides significantly more information for understanding and optimizing these interactions than the current “general albumin binding” assay-readout.

In addition to the previously reported 7-nitrobenzo-2-oxa-1,3-diazole (NBD) labelled fatty acid (NBD-FA) as molecular probe, more site-specific fluorescent ligands are then needed as additional probes to selectively label the ibuprofen binding site in the albumin “Sudlow-site II” and the warfarin binding site (Sudlow-site-I) respectively. These new labels should exhibit only minimal spectral overlap with the existing probe NBD-FA to allow for selective optical detection and possess a similar binding affinity. Their saturation curve should show a clear plateau region, thus indicating specific binding to the corresponding albumin binding site. Moreover a clear 1:1 binding stoichiometry must be revealed by Job-plots^[5] (see supplemental).

While such behavior is indicative for specific binding and release, effects like quenching, reorganization or rebinding cannot be ultimately ruled out.

Finally, competition experiments of newly identified molecular probes with known albumin site-specific binders need to be performed to clearly assign these probes to known albumin binding sites. These new probe molecules should be able to release the known fluorescent ligands in a concentration-dependent fashion.


Recently, several publications reported syntheses and applications of BODIPY (4,4-difluoro-4-bora-3a,4a-diaza-s-indacene) dyes exhibiting similar “switch-on” characteristics upon binding^[6] as observed for NBD-FA and other NBD-labelled ligands.^[3, 7] BODIPY derivatives cover a wide range of wavelengths of their fluorescence maxima which strongly depend on their chemical substitution patterns. They have been described as polarity-based fluorescence sensors,^[6] as sensors for hydrophobic protein surfaces,^[8] as products of diversity-oriented library synthesis for unbiased screening^[9] of hydrophobic protein surfaces or in combination with click-chemistry.^[10] Kim et al.^[11] recently described NIR-labelling with 8-amino BODIPY derivatives, further increasing the application range of these dyes.

In several reports,^[6, 8–11] the affinity of BODIPY derivatives to bovine serum albumin (BSA) has been described. Er et al.^[12] have published a mixture of two fluorescent dyes for dual drug site mapping on HSA using a BODIPY derivative selected from a product library as a label for Sudlow-site II. Competition

- [a] Dr. M. Wagner, Dr. H. Schreuder, Dr. H. Matter, Dr. S. M. Petry
Sanofi-Aventis Deutschland GmbH,
Industriepark Höchst, 65926 Frankfurt am Main, Germany
E-mail: StefanMatthias.Petry@sanofi.com
- [b] Dr. L. Wenskowsky, Prof. Dr. T. Opatz
Institute of Organic Chemistry, Johannes Gutenberg-University
Duesbergweg 10–14, 55128 Mainz, Germany
E-mail: opatz@uni-mainz.de
- [c] J. Reusch
Dynamic Biosensors GmbH
Lochhamer Straße 15, 82152 Martinsried/Planegg, Germany

Supporting information for this article is given via a link at the end of the document

experiments with ibuprofen proved the identity of the binding site. However, these authors chose dansylamide as the second label which we found to exhibit a 4:1^[3] instead of the required 1:1 binding stoichiometry, which is also in agreement with results from X-ray crystallography by Curry^[13] et al. These studies led us to the design and synthesis^[12, 14] of a small, directed library of BODIPY derivatives **5a–c** for targeting Sudlow-site II (Scheme 1, see Supplemental Experimental Procedures).



Scheme 1. Synthesis of BODIPY derivatives **5a–c**.

First, we investigated these compounds for unbiased HSA binding. Saturation curves were recorded and the respective K_D values were determined as described earlier.^[3]

Compounds with suitable fluorescence emission wavelength that showed clear saturation in a 1:1 HSA:BODIPY stoichiometry were further investigated in competition experiments with ibuprofen (**1**). BODIPY derivative **5a** was finally selected as an ibuprofen (**1**) competitive probe showing the desired 1:1 release stoichiometry (Figure 1C).





Figure 1. A) Absorption and emission spectra (both normalized) of dimethylamino-BODIPY **5a** (40 μM ethanol). λ_{exc} = 625 nm. B) Titration of dimethylamino-BODIPY **5a** (0–50 μM) to HSA (12.5 μM) (K_D = 4.6 ± 0.8 μM, R^2 = 0.9176). λ_{exc} = 615 nm. λ_{em} = 690 nm. C) Determination of the binding stoichiometry of dimethylamino-BODIPY **5a** to HSA (12.5 μM, Job plot). λ_{exc} =

615 nm. λ_{em} = 690 nm. D) Competition experiment of ibuprofen (**1**) against dimethylamino-BODIPY **5a** on HSA (12.5 μM each). λ_{exc} = 615 nm. λ_{em} = 690 nm.

We then intended to use a third fluorescent ligand in order to directly label the warfarin binding site (Sudlow-site I) as well. Again, binding affinity in the same range as for NBD-FA and **5a** as well as a 1:1 binding stoichiometry were the selection criteria for the optimal ligand.



Scheme 2. Synthesis of warfarin derivatives **8a–c**.

Warfarin (**2**) itself showed suitable fluorescence intensity but unfortunately shows absorption at the excitation wavelength of dimethylamino-BODIPY **5a** (Figure S5). Thus, a set of coumarin derivatives **8a–c** was synthesized^[15] and screened as described for compound **5a** (Scheme 2). The warfarin derivative **8a** fulfilled the aforementioned criteria (Figure 2A–D). Furthermore, its fluorescence emission was substantially different from those of NBD-FA and BODIPY **5a**, so that the simultaneous selective quantification of all three ligands should be possible in an unprecedented three-color fluorescent assay.




Figure 2. A) Absorption and emission spectra (both normalized) of warfarin derivative **8a** (40 μM ethanol). λ_{exc} = 340 nm. B) Titration of warfarin derivative

8a (0–400 μM) to HSA (25 μM) ($K_D = 26.5 \pm 3.0 \mu\text{M}$, $R^2 = 0.9607$). $\lambda_{\text{exc}} = 330 \text{ nm}$, $\lambda_{\text{em}} = 384 \text{ nm}$. C) Determination of the binding stoichiometry of warfarin derivative **8a** to HSA (25 μM , Job plot). $\lambda_{\text{exc}} = 330 \text{ nm}$, $\lambda_{\text{em}} = 384 \text{ nm}$. D) Competition experiment of iophenoxic acid (**3**) against warfarin derivative **8a** on HSA (25 μM each). $\lambda_{\text{exc}} = 330 \text{ nm}$, $\lambda_{\text{em}} = 384 \text{ nm}$.

Since we have determined the K_D values of our probes exclusively by titration experiments, we considered the independent validation of these values with a different method.

To this end, independent characterization of all three ligands has been performed using the switchSENSE technology^[16] (Dynamic Biosensors, Munich). This fluorescence-based biosensor allows for the biophysical characterization of biomolecules determining kinetic rate constants of association and dissociation (k_{ON} and k_{OFF}), and the dissociation constant (K_D) via fluorescence proximity sensing. Furthermore, switchSENSE enables the user to detect intramolecular conformational changes by determination of the hydrodynamic friction of surface-bound complexes using a complementary measurement mode (molecular dynamics mode).^[17]

Here, switchSENSE was used for K_D determination via kinetic rate constants and via binding equilibrium as well as analyzing conformational changes of HSA upon binding of different compounds (induced fit). For interaction analysis, HSA was immobilized on a gold microelectrode via short, single stranded DNA nanolevers using standard thiol coupling chemistry. The complementary DNA strand, covalently attached to the gold microelectrodes via Au-S bonds, carries a fluorescent dye (oxazine derivative) on the top end for signal detection. For the measurements of HSA with NBD-FA, dimethylamino-BODIPY **5a** and warfarin derivative **8a**, K_D values in the micromolar range were determined (NBD-FA: $K_D = 27.5 \mu\text{M}$, BODIPY **5a**: $K_D = 5.2 \mu\text{M}$, warfarin derivative **8a**: $K_D = 25.5 \mu\text{M}$) (Figures S8–11). In the molecular dynamics mode, HSA showed a significant expansion upon NBD-FA binding. Due to a higher hydrodynamic radius in solution (measured in dynamic response units), the molecular dynamics of HSA slows down (Figure S10). This effect was neither observed for BODIPY **5a** nor for warfarin derivative **8a** (for further details see supplemental).

To check the suitability of the chosen trichromatic dye cocktail for simultaneous detection (for a graphical representation of the fluorescent properties of all three labels, see Figure 3), finally a mixture of 0.3 equivalents of each label was added to a solution of HSA (25 μM). This stoichiometry does not address differences in K_D values, since this setup is used for the localization of binding. For determination of K_D we generally use and propose independent measurements with a single dye. Since the affinity of our labels is high enough to ensure specific binding we could reduce the equivalents added. We found that 0.3 equivalents of each label delivered the highest assay sensitivity and reduces the possibility of unspecific rebinding.




Figure 3. Absorption spectra (normalized) of warfarin derivative **8a** (blue), NBD-FA (green) and dimethylamino-BODIPY **5a** (red). Excitation wavelengths for each dye were drawn.

Gratifyingly, all three labels also behaved well in the chosen combination and in each case permitted the selective competition in the respective binding site. Figures 4–5 show typical displacement




Figure 4. Competition experiments against NBD-FA/BODIPY **5a**/coumarin **8a** (8 μM each) on HSA. • NBD-FA. ♦ BODIPY **5a**. ■ warfarin derivative **8a**. A) ibuprofen (**1**). B) ketoprofen (**4**). C) indomethacin (**5**). D) iophenoxic acid (**3**).

experiments. Commercial drugs were added to HSA loaded with the trichromatic cocktail at the concentrations given. The assay was performed in parallel in 96-well-microtiter plates, which were analyzed with a fluorescent plate reader preset to the wavelengths characteristic for the three reporter dyes (NBD-FA: $\lambda_{\text{exc}} = 435 \text{ nm}$, $\lambda_{\text{em}} = 535 \text{ nm}$. BODIPY **5a**: $\lambda_{\text{exc}} = 615 \text{ nm}$, $\lambda_{\text{em}} = 690 \text{ nm}$. warfarin derivative **8a**: $\lambda_{\text{exc}} = 330 \text{ nm}$, $\lambda_{\text{em}} = 384 \text{ nm}$). Ibuprofen (**1**), ketoprofen (**4**) and indomethacin (**5**) selectively displaced dimethylamino-BODIPY **5a** indicating selective binding to the albumin ibuprofen binding site (Sudlow-site II) (Figures 4A–C), Indomethacin (**5**) in addition to **5a** also displaces the warfarin derivative **8a**.^[18] At high concentrations, both NSAIDs (non-steroidal anti-inflammatory drugs) **1** and **4** lead to

decreased fluorescence, indicating secondary binding sites.^[19] lophenoxic acid (**3**) is competitive to coumarin derivative **8a**^[20] as well as to dimethylamino-BODIPY **5a**.^[21]




Figure 5. Competition experiments against NBD-FA/BODIPY **5a**/coumarin **8a** (8 μ M each) on HSA. \bullet NBD-FA. \blacklozenge BODIPY **5a**. \blacksquare warfarin derivative **8a**. A) sulfasalazine (**6**). B) balsalazide (**7**).

For sulfasalazine (**6**), a marketed drug for treatment of e.g. rheumatoid arthritis, ulcerative colitis, and Crohn's disease, the fluorescence signal decreases for NBD-FA and coumarin **8a**,^[3, 22] which indicates release of the site-specific probe-ligands while BODIPY **5a** in (Sudlow-site II) is not released (Figure 5A). The same behavior is observed for the structurally related drug balsalazide (**7**) (Figure 5B).^[3]

Thus, NBD-FA as well as coumarin **8a** are effectively competed by sulfasalazine (**6**). This interesting observation can be explained by direct competition between ligand and molecular probes in two separate albumin binding sites, by neighboring sites, or by an allosteric binding event. The similar shape of displacement curves for NBD-FA and warfarin derivative **8a** suggests a simultaneous release of both probes by only a single sulfasalazine (**6**) molecule rather than binding of two sulfasalazine molecules to these sites. Independent kinetic analysis using switchSENSE technology revealed a K_D of 1.31 μ M, with a remarkably fast K_{ON} ($5.76 \cdot 10^{-1} \text{ M}^{-1}\text{s}^{-1}$). As expected, only one binding event could be observed.

A comparable situation was described by F. Yang et al.^[23] They have observed independent binding of three different compounds (cinnamic acid, lamivudine, indomethacin) within this hydrophobic binding site, while binding is associated with a conformational change of albumin.

To further investigate this interesting binding behavior, we were able to solve an X-ray crystal structure of sulfasalazine (**6**) in complex with HSA, obtained by co-crystallization. The anisotropic crystal diffracted to 2.2 Å in the best and to 2.8 Å in the worst direction. The structure was solved by molecular replacement and the resulting electron density maps were clear. The conformation of HSA is more similar to its fatty-acid free conformation^[18d] (RMSD 2.5 Å to the PDB structure 1uor) than to the fatty-acid bound structure^[2a] (RMSD 5.5 Å to the myristate-bound structure 1e7g). This might be due to our crystallization conditions in the absence of fatty acids. In total, three molecules of sulfasalazine (**6**) are bound to different HSA binding sites. As expected, one of them is located in the Sudlow-site I with clear electron density. Two other molecules showing lower occupancy are bound to either HSA domain IIIb or to the interface between domains IIIa and IIIb (see the Supporting Information). These

additional interactions might be attributed to high sulfasalazine (**6**) concentration during crystallization (16 mM **6**, 2 mM HSA). K_D values for the two additional sites were estimated from the fractional occupancies to be 0.9 mM and 3.4 mM, respectively (see Supporting Information).




Figure 6. X-ray crystal structure of HSA in complex with sulfasalazine (**6**) (PDB ID 6R7S resolution 2.2 Å). A) Binding of sulfasalazine (**6**) (orange carbon atoms) to Sudlow-site I plus Fo-Fc omit map contoured at 3 σ . Key amino acids and water oxygen atoms are indicated. B) Binding site analysis of the HSA-**6** complex using SiteMap. Red regions indicate favorable potential H-bond acceptor interactions; blue regions indicate favorable H-bond donor interactions and yellow regions indicate favorable hydrophobic interactions. C) Superposition of sulfasalazine (**6**) with the enantiomers of warfarin (**2**) from PDB 1ha2 (S-warfarin, dark-blue) and 1h9z (R-warfarin, light-blue), both at 2.5 Å resolution.^[24] D) Superposition of sulfasalazine with NBD-FA (green carbon atoms) from PDB 6ezq^[3] at 2.5 Å resolution.

Figure 6A shows experimentally determined interactions of sulfasalazine bound to Sudlow-site I. While its location in this binding site agrees with previous docking studies, the molecule turned out to be flipped by 180°.^[3] The salicylate carboxylate group is engaged in hydrogen-bond interactions and a salt-bridge to Arg257 and Ser287 side chains. The central sulfasalazine aromatic ring is involved in a cation- π interaction to Arg222. Both attached sulfonamide oxygen atoms interact via bridging water molecules with either Trp214 or Lys199 side chains. The distal pyridine aromatic ring is stacked on top of the indole ring of Trp214 with both nitrogen atoms likely arranged close together (distance 3.2 Å). The indole NH and the pyridine N are interacting with bridging water molecules.

The binding site was then analyzed using SiteMap^[25] (Figure 6B) to identify favorable hydrophobic (yellow), hydrogen-bond donor (blue) and acceptor interactions (red), which are in agreement to ligand interactions. A comparison with HSA X-ray crystal structures with (*R*)- or (*S*)-warfarin (Figure 6C, PDB 1ha2 for (*S*)-warfarin and 1h9z for (*R*)-warfarin, both 2.5 Å resolution) shows that the sulfasalazine salicylate and its central aromatic ring occupy parts of the warfarin binding site. This finding

explains the competitive binding observed for sulfasalazine (**6**) and warfarin (**2**).

A comparison with the HSA-NBD-FA complex (Figure 6D) reveals that the pyrimidine ring overlaps with the NBD group, both being stacked on top of the indole ring of Trp214. Also the remaining parts of both ligands overlap, again explaining the strong competition of both ligands for binding to HSA. Although a significant excess of sulfasalazine (**6**) was used for crystallization (vide supra), the X-ray crystal structure of its HSA complex revealed the bridging of the fatty acid binding site and Sudlow-site I by only a single sulfasalazine molecule.

The trichromatic ligand cocktail was then employed for the analysis of the site-specific binding behaviour of a series of commercial drugs. We also applied it for the analysis of biomolecules which carry albumin-binding moieties, such as fatty acids. As an example we describe the kinetic analysis of HSA-binding of two new antidiabetic peptides. Table 1 summarizes the results, while respective titration curves are given in the Supporting Information.

Table 1. Localization of binding site and determination of K_D of fatty acid modified GLP1 agonists.

Compound	Duration of action	Albumin binding site	K_D ^[b]
Liraglutide	Once daily	NBD-FA ^[a] binding site	8.9 μ M
Semaglutide	Once weekly	NBD-FA binding site and FA3	1.9 μ M

[a] NBD-FA: (7-nitrobenz-2-oxa-1,3-diazol-4-yl)-C₁₂ fatty acid.

[b] K_D : dissociation constant.

Interestingly, these binding events were also associated with changes in HSA's hydrodynamic diameter.

The results obtained with the trichromatic assay are in accordance to literature^[3, 18b, 18c, 19-20, 22] and to X-ray crystal structures of HSA-ligand complexes.^[18a, 18d, 21] The advantage of the present method over individual analysis of binding sites with only a single fluorescent ligand at a time is the reduction of the number of experiments required as well as of the overall protein consumption. The simultaneous analysis of displacement patterns also permits the rapid characterization of HSA binding behaviour for larger (bio)-molecules displaying polyvalent binding characteristics.

We propose this general concept for the investigation of further biomolecules containing different binding sites. In the case of HSA, the developed assay system is useful for the initial characterization and differentiation of albumin-binding drugs and is of potential relevance for the prediction of drug-drug and drug-food interactions (in particular involving fatty acids and cholesterol). It can also be used for nanobodies or the newly described small molecule-based HSA-binders^[26] employed to prolong plasma half-life.

Experimental Section

For experimental details, crystallographic data and further biochemical data, see the Supporting Information.

Acknowledgements

We thank Dr. Michael Kurz (Frankfurt) and Dr. J. Liermann (Mainz) for NMR spectroscopy, Dr. N. Hanold (Mainz, deceased) for mass spectrometry, Dr. Dieter Schollmeyer (Mainz) for X-Ray crystal structure analysis of BODIPY derivative **5a**, Alexander Liesum (Frankfurt) for help with crystallization and mounting of the HSA crystal, Joachim Diez (Frankfurt) for data collection, Petra Loenze (Frankfurt) for help with data processing, and Pia Freisewinkel (Frankfurt) for competition experiments with fatty acid modified GLP1 agonists.

Keywords: Multicolor Assay • switchSENSE Technology • kinetic investigation half-life extension • Drug interaction • albumin binding

- [1] a) K. Oettl, R. E. Stauber, *Brit. J. Pharmacol.* **2007**, *151*, 580-590; b) K. J. Fehske, W. E. Müller, U. Wollert, *Biochem. Pharmacol.* **1981**, *30*, 687-692; c) A. A. Spector, E. C. Santos, J. D. Ashbrook, J. E. Fletcher, *Ann. N. Y. Acad. Sci.* **1973**, *226*, 247-258; d) A. A. Spector, *J. Lipid Res.* **1975**, *16*, 165-179; e) M. Fasano, S. Curry, E. Terreno, M. Galliano, G. Fanali, P. Narciso, S. Notari, P. Ascenzi, *IUBMB Life* **2005**, *57*, 787-796; f) G. Fanali, A. di Masi, V. Trezza, M. Marino, M. Fasano, P. Ascenzi, *Mol. Aspects Med.* **2012**, *33*, 209-290.
- [2] a) A. A. Bhattacharya, T. Grüne, S. Curry, *J. Mol. Biol.* **2000**, *303*, 721-732; b) S. Curry, H. Mandelkow, P. Brick, N. Franks, *Nat. Struct. Biol.* **1998**, *5*, 827-835; c) I. Petitpas, T. Grüne, A. A. Bhattacharya, S. Curry, *J. Mol. Biol.* **2001**, *314*, 955-960; d) S. Curry, *Drug Metab. Pharmacokinet.* **2009**, *24*, 342-357.
- [3] L. Wenskowsky, H. Schreuder, V. Derdau, H. Matter, J. Volkmar, M. Nazaré, T. Opatz, S. Petry, *Angew. Chem.* **2018**, *130*, 1056-1060.
- [4] a) D. S. Fredrickson, R. S. Gordon, *J. Clin. Invest.* **1958**, *37*, 1504-1515; b) T. Peters Jr., *All about albumin: biochemistry, genetics, and medical applications*, Academic press, New York, **1995**.
- [5] P. Job, *Ann. Chim. Appl.* **1928**, *9*, 113-203.
- [6] H. Sunahara, Y. Urano, H. Kojima, T. Nagano, *J. Am. Chem. Soc.* **2007**, *129*, 5597-5604.
- [7] a) J. Rohacova, M. L. Marin, M. A. Miranda, *J. Phys. Chem. B* **2010**, *114*, 4710-4716; b) K. L. Diehl, M. A. Ivy, S. Rabidoux, S. M. Petry, G. Müller, E. V. Anslyn, *Proc. Natl. Acad. Sci. U.S.A.* **2015**, *112*, E3977-E3986; c) A. Chattopadhyay, *Chem. Phys. Lipids* **1990**, *53*, 1-15.
- [8] N. Dorh, S. Zhu, K. B. Dhungana, R. Pati, F.-T. Luo, H. Liu, A. Tiwari, *Sci. Rep.* **2015**, *5*, 18337.
- [9] H. KyuáKim, *Chem. Commun.* **2011**, *47*, 2339-2341.
- [10] L. P. Jameson, N. W. Smith, O. Annunziata, S. V. Dzyuba, *Phys. Chem. Chem. Phys.* **2016**, *18*, 14182-14185.
- [11] D. Kim, D. Ma, M. Kim, Y. Jung, N. H. Kim, C. Lee, S. W. Cho, S. Park, Y. Huh, J. Jung, *J. Fluoresc.* **2017**, *27*, 2231-2238.
- [12] J. C. Er, M. Vendrell, M. K. Tang, D. Zhai, Y.-T. Chang, *ACS Comb. Sci.* **2013**, *15*, 452-457.
- [13] A. J. Ryan, J. Ghuman, P. A. Zunszain, C.-W. Chung, S. Curry, *J. Struct. Biol.* **2011**, *174*, 84-91.
- [14] J.-S. Lee, N.-y. Kang, Y. K. Kim, A. Samanta, S. Feng, H. K. Kim, M. Vendrell, J. H. Park, Y.-T. Chang, *J. Am. Chem. Soc.* **2009**, *131*, 10077-10082.


- [15] a) P.-Y. Chen, Y.-H. Wu, M.-H. Hsu, T.-P. Wang, E.-C. Wang, *Tetrahedron* **2013**, *69*, 653-657; b) N. Halland, T. Hansen, K. A. Jørgensen, *Angew. Chem.* **2003**, *115*, 5105-5107.
- [16] A. Cléry, T. J. Sohler, T. Welte, A. Langer, F. H. Allain, *Methods* **2017**, *118*, 137-145.
- [17] a) A. Langer, P. A. Hampel, W. Kaiser, J. Knezevic, T. Welte, V. Villa, M. Maruyama, M. Svejda, S. Jähner, F. Fischer, *Nat. Commun.* **2013**, *4*, 2099; b) A. Langer, W. Kaiser, M. Svejda, P. Schwertler, U. Rant, *J. Phys. Chem. B* **2014**, *118*, 597-607.
- [18] a) J. Ghuman, P. A. Zunszain, I. Petitpas, A. A. Bhattacharya, M. Otagiri, S. Curry, *J. Mol. Biol.* **2005**, *353*, 38-52; b) G. Sudlow, D. Birkett, D. Wade, *Mol. Pharmacol.* **1976**, *12*, 1052-1061; c) H. Watanabe, S. Tanase, K. Nakajou, T. Maruyama, U. Kragh-Hansen, M. Otagiri, *Biochem. J.* **2000**, *349*, 813-819; d) X. M. He, D. C. Carter, *Nature* **1992**, *358*, 209-215.
- [19] a) F. Li, D. Zhou, X. Guo, *J. Chromatogr. Sci.* **2003**, *41*, 137-141; b) A. Kober, I. Sjöholm, *Mol. Pharmacol.* **1980**, *18*, 421-426.
- [20] G. Sudlow, D. J. Birkett, D. N. Wade, *Clin. Exp. Pharmacol. Physiol.* **1975**, *2*, 129-140.
- [21] A. J. Ryan, C.-w. Chung, S. Curry, *BMC Struct. Biol.* **2011**, *11*, 18.
- [22] J. Równicka-Zubik, A. Sulkowska, J. Pozycka, K. Gazdzicka, B. Bojko, M. Maciazek-Jurczyk, W. Sulkowski, *J. Mol. Struct.* **2009**, *924*, 371-377.
- [23] F. Yang, J. Yue, L. Ma, Z. Ma, M. Li, X. Wu, H. Liang, *Mol. Pharmaceutics* **2012**, *9*, 3259-3265.
- [24] I. Petitpas, A. A. Bhattacharya, S. Twine, M. East, S. Curry, *J. Biol. Chem.* **2001**, *276*, 22804-22809.
- [25] T. A. Halgren, *J. Chem. Inf. Model.* **2009**, *49*, 377-389.
- [26] a) E. M. Bech, S. L. Pedersen, K. J. Jensen, *ACS Med. Chem. Lett.* **2018**, *9*, 577-580; b) D. Sleep, J. Cameron, L. R. Evans, *Biochim. Biophys. Acta, Gen. Subj.* **2013**, *1830*, 5526-5534; c) S. B. van Witteloostuijn, S. L. Pedersen, K. J. Jensen, *ChemMedChem* **2016**, *11*, 2474-2495; d) P. Kurtzhals, S. Havelund, I. Jonassen, B. Kiehr, U. Larsen, U. Ribbel, J. Markussen, *Biochem. J.* **1995**, *312*, 725-731; e) R. E. Kontermann, *BioDrugs* **2009**, *23*, 93-109; f) R. E. Kontermann, *Curr. Opin. Biotechnol.* **2011**, *22*, 868-876; g) M. F. Koehler, K. Zobel, M. H. Beresini, L. D. Caris, D. Combs, B. D. Paasch, R. A. Lazarus, *Bioorg. Med. Chem. Lett.* **2002**, *12*, 2883-2886.

Entry for the Table of Contents (Please choose one layout)

Layout 1:

COMMUNICATION

A trichromatic fluorescent binding and displacement assay for simultaneous monitoring and localization of three individual binding sites on HSA has been developed using highly specific binding molecules. Whereas localization is confirmed by X-ray, kinetic data are independently confirmed by switchSENSE technology, revealing important data on conformational changes. The novel tool permits the detailed characterization of albumin-binding drugs and prediction of drug-drug and drug-food interactions.



Lea Wenskowsky, Michael Wagner, Johannes Reusch, Herman Schreuder, Hans Matter, Till Opatz, and Stefan Matthias Petry**

Page No. – Page No.

Resolving Binding Events on the Multifunctional Human Serum Albumin