

Communications to the Editor

[Chem. Pharm. Bull.]
30(7)2641-2643(1982)


ENANTIOSELECTIVE SYNTHESIS OF (-)-PHYSOSTIGMINE

Seiichi Takano,* Emiko Goto, Michiyasu Hiram, and Kunio Ogasawara
Pharmaceutical Institute, Tohoku University, Aobayama, Sendai 980,
Japan

The first enantioselective synthesis of (-)-physostigmine (1), a potent cholinesterase-inhibiting alkaloid from *Physostigma venenosum* Balf., has been accomplished using (S)-(-)-benzyl 2,3-epoxypropyl ether (5) as a chiral synthon.

KEYWORDS—chiral synthesis; chiral synthon;
(-)-physostigmine; (-)-eseroline; (-)-esermethole; cholinesterase inhibitor

(-)-Physostigmine (1),¹⁾ a major alkaloid of the fruit (Calabar bean) of *Physostigma venenosum* Balf., is a long-established compound and is being currently used as a cholinesterase inhibitor. Because of the medicinal importance of the alkaloid, a number of syntheses²⁾ have been reported since the first one by Julian and Pikel³⁾ in 1935. However, we know of no report to date of an enantioselective approach to this alkaloid. We report here the first enantioselective synthesis of (-)-physostigmine (1) in satisfactory optical purity using a readily accessible chiral synthon, (S)-(-)-benzyl 2,3-epoxypropyl ether (5).



Scheme

(S)-(-)-Benzyl 2,3-epoxypropyl ether (5),^{4,5,6} obtained from (D)-mannitol-1,2:5,6-diacetonide, was treated with the carbanion prepared *in situ* from 3-methoxybenzyl cyanide and lithium diisopropylamide (LDA) in tetrahydrofuran (THF) (-78°C room temperature) to give the cyano-alcohol (6) as a mixture of epimers. Hydrolysis of the crude mixture with hot ethanolic potassium hydroxide (7h) afforded the γ -lactone (7)⁷ as a mixture of epimers after acid work-up (dil. HCl). Overall yield of (7) from (5) was 63%. Alkylation of the mixture (7) with methyl iodide in the presence of LDA in THF (-78°C room temperature) led to a stereoselective introduction of the methyl group from the less hindered face of the molecule to give 80% yield of the desired compound (8), $[\alpha]_D -42.1^\circ$ (c=1.34, MeOH), along with 11% yield of the undesired epimer after separation employing silica-gel column chromatography. The benzyl group of (8) was removed by hydrogenolysis under an atmospheric pressure of hydrogen in the presence of 10% palladized charcoal in methanol containing a trace of perchloric acid to give the hydroxy-lactone (9) in quantitative yield. Upon sequential one-pot treatment, *viz.*, hydrolysis with methanolic potassium hydroxide, oxidative cleavage with sodium periodate, and reduction with sodium borohydride, followed by acid work-up (dil. HCl), the compound (9) furnished the lactone (10), $[\alpha]_D -72.1^\circ$ (c=1.01, MeOH) in 96% overall yield. Condensation of the lactone (10) with 40% aqueous methylamine at 180°C using a sealed tube (12h) gave the lactam (11), $[\alpha]_D -79.9^\circ$ (c=1.22, MeOH) in 82% yield.

Although we were unable to find optimal conditions for the regioselective introduction of the nitrogen group on the aromatic ring, the desired compound (13) was obtained from (11) in a pure state in 27% overall yield. Thus, treatment of (11) with cupric nitrate trihydrate in acetic anhydride⁸ (5-10°C, 30min) furnished a mixture of three regio-isomeric compounds from which 19% of the 4-nitro isomer (12:4-NO₂) and 72% of a mixture of the 2- and 6-isomers were separated through silica-gel column chromatography. The latter mixture was then hydrogenated under an atmospheric pressure of hydrogen in the presence of platinum catalyst in methanol to give a mixture of the amino derivatives which could be separated by using a silica-gel column to give the desired 6-amino-lactam (13) in 38% yield accompanied by 30% yield of the unnecessary 2-amino derivative. Reduction of the lactam (13) with a 1.5 equivalent molar excess of lithium aluminum hydride in THF at room temperature (1h) afforded the tricyclic aminal (4), mp 68-69°C, $[\alpha]_D -101^\circ$ (c=0.388, CHCl₃), in 60% yield. Reductive N-methylation with formalin and sodium cyanoborohydride in acidic media (pH 5-6) at room temperature gave (-)-esermethole (3),⁹ $[\alpha]_D -123^\circ$ (c=0.412, benzene) in 44% yield. Its spectroscopic data (IR, ¹H-NMR, MS) as well as chromatographic behavior (tlc on silica-gel and alumina plates) were completely identical with those of an authentic material (3), $[\alpha]_D -129^\circ$ (c=0.326, benzene), prepared from a natural physostigmine (1).¹⁰ O-De-methylation of (3) with boron tribromide in methylene chloride at -78 ~ 10°C afforded (-)-eseroline (2), mp 175-178°C, $[\alpha]_D -82.1^\circ$ (c=0.524, CHCl₃) (authentic (-)-eseroline (2),¹⁰ mp 178-182°C, $[\alpha]_D -88.7^\circ$ (c=0.284, CHCl₃) (mixed mp 178-182°C) in 76% yield. Since (-)-eseroline (2) has been nicely converted into (-)-physostigmine (1) in one step,¹¹ this is the first enantioselective synthesis.

REFERENCES AND NOTES

- 1) Pertinent reviews: (a) E. Coxworth, "The Alkaloids," R.H.F. Manske, Edn., Vol. VIII, Academic Press, New York, 1965, p. 1. (b) B. Robinson, "The Alkaloids," R.H.F. Manske, Edn., Vol. X, Academic Press, New York, 1968, p. 383.
- 2) Recent racemic synthesis: J.B.P.A. Wijnberg and W.N. Speckamp, *Tetrahedron*, 34, 2399(1978).
- 3) P.L. Julian and J. Pikl, *J. Am. Chem. Soc.*, 57, 755(1935).
- 4) A.K.M. Anisuzzaman and L.N. Owen, *J. Chem. Soc. (C)*, 1967, 1021.
- 5) S. Takano, E. Goto, M. Hirama, and K. Ogasawara, *Heterocycles*, 16, 381(1981).
- 6) The ether (5) was used as a chiral synthon for the first enantioselective synthesis of (-)-mesembrine: see, S. Takano, Y. Imamura, and K. Ogasawara, *Tetrahedron Lett.*, 22, 4479(1981).
- 7) Satisfactory spectral (IR, ¹H-NMR, MS) and analytical (combustion and high resolution MS) data were obtained for all new compounds.
- 8) Cf. F.A. Carey and R.M. Giuliano, *J. Org. Chem.*, 46, 1366(1981).
- 9) Recent synthesis of racemic esermethole (3): M. Ikeda, S. Matsugashita, and Y. Tamura, *J. Chem. Soc. Perkin Trans. 1*, 1977, 1166.
- 10) Natural physostigmine (1) was hydrolyzed with ethanolic potassium hydroxide to give (-)-eseroline (2) which gave (-)-esermethole (3) on treatment with diazomethane.
- 11) M. Polonovski and C. Nitzberg, *Bull. Soc. Chim. Fr.*, [iv], 19, 33(1916).

(Received June 1, 1982)