Tetrazine-*trans*-cyclooctene ligation for the rapid construction of ¹⁸F labeled probes[†]

Zibo Li,*^{*a*} Hancheng Cai, \ddagger^{a} Matthew Hassink, \ddagger^{b} Melissa L. Blackman,^{*b*} Richard C. D. Brown,^{*c*} Peter S. Conti^{*a*} and Joseph M. Fox*^{*b*}

Received 5th August 2010, Accepted 6th September 2010 DOI: 10.1039/c0cc03078c

A radiolabeling method for bioconjugation based on the Diels–Alder reaction between 3,6-diaryl-s-tetrazines and an ¹⁸F-labeled *trans*-cyclooctene is described. The reaction proceeds with exceptionally fast rates, making it an effective conjugation method within seconds at low micromolar concentrations.

Positron emission tomography (PET) is a non-invasive imaging modality that utilizes positron-emitting radionuclides (C-11, N-13, O-15 and F-18).¹ F-18 PET has a number of attributes that make it clinically attractive, including near 100% positron efficiency, a very high specific radioactivity, and a short half-life of ~110 min.² However, the short half-life of F-18 and the poor nucleophilicity of fluoride render it difficult to incorporate F-18 in complex molecules. Currently, radiochemistry is a major limiting factor for the field of PET.³ Despite recent advances,^{4,5} challenges exist for improving F-18 incorporation with respect to reaction rates, efficiency, and selectivity.

Recently, we introduced the tetrazine–*trans*-cyclooctene ligation ('TTCO ligation', Fig. 1) as a method of bioconjugation that proceeds with fast reaction rates without the need for catalysis.^{6,7} *trans*-Cyclooctene derivatives are readily prepared from *cis*-cyclooctenes using a photochemical flow-reaction that we developed.⁸ While a variety of *s*-tetrazine derivatives were known to react with strained alkenes,⁹ we have found



Fig. 1 Tetrazine-trans-cyclooctene ligation ('TTCO-ligation').

- ^a Molecular Imaging Center, Department of Radiology, University of Southern California, Los Angeles 90033, USA.
- *E-mail: ziboli@usc.edu; Fax: +1 323-442-3252* ^b Brown Laboratories, Department of Chemistry and Biochemistry, University of Delaware, Newark, DE 19803, USA.
- *E-mail: jmfox@udel.edu; Fax: +1 302-831-6335*
- ^c School of Chemistry, University of Southampton, Highfield, Southampton, SO171BJ, UK. E-mail: rcb1@soton.ac.uk; Fax: +44 (0)238-059-6805
- † Electronic supplementary information (ESI) available: Experimental procedures, NMR spectra and Radio-HPLC data. See DOI: 10.1039/ c0cc03078c
- ‡ Equal contributions were made by these authors.

that 3,6-diaryl-s-tetrazines offer an excellent combination of fast reactivity and stability for both the conjugate and starting material.⁶ In particular, 3,6-di(2-pyridyl)-s-tetrazines were shown to display excellent characteristics.^{6,10} Thus, the reaction between *trans*-cyclooctene and **1a** proceeds with a rapid rate $(k_2 \approx 2000 \text{ M}^{-1} \text{ s}^{-1} \text{ in 9}: 1 \text{ MeOH}: \text{ water})$, and is successful in cell media and cell lysate.⁶ 3,6-Di(2-pyridyl)-s-tetrazines can easily be functionalized as their amido derivatives (**1b**),⁶ which display excellent stability toward water and biological nucleophiles.¹⁰

After we described the TTCO-ligation, the groups of Hilderbrand^{11a} and Pipkorn and Braun^{11b} described ligations between tetrazines and less reactive dienophiles. However, the use of *trans*-cyclooctene is the key to fast rates of reactivity, and other groups have subsequently developed applications for the TTCO-ligation.^{11c,d,12}

Because of the fast rate of reactivity, the TTCO-ligation offers opportunities for the rapid conjugation of radionuclides to biomolecules, both of which are often available only at low concentration. Very recently, Robillard and coworkers elegantly demonstrated that our system for TTCO-ligation could be utilized in In¹¹¹-imaging of tumors in live mice using a DOTA-derivative of **1b**.¹² Herein, we report the development of an extremely fast and reactive method for generating ¹⁸F labeled probes based on the TTCO-ligation.

While a number of indirect methods for ¹⁸F-incorporation into tetrazines or *trans*-cyclooctene derivatives could be envisioned (*e.g.* derivatization by *N*-succinimidyl-4-¹⁸F-fluorobenzoate),¹³ such approaches are complicated by the need to carry ¹⁸F through added manipulations. Thus, we focused on the development of direct methods for ¹⁸F-incorporation *via* reactions with fluoride ion, with an initial focus on the



Scheme 1 Synthesis of ¹⁸F-labeled tetrazine. Attempts to prepare 3 by nucleophilic substitution were low yielding. Fluorination of 4 (¹⁸F-fluoride, MeCN, 85 °C, 15 min) gives 5 in only 1% radiochemical yield.





Scheme 2 Synthesis of *trans*-cyclooctene nosylate 8. *Reagents and conditions*: (a) NaH, α -bromoacetic acid. (b) CH₂N₂, 69%, 2 steps. (c) DIBAL, 78%. (d) Methyl benzoate-sensitized photoisomerization with active removal of *trans*-isomers, AgNO₃, 55%. (e) NsCl, Et₃N, 87%.

synthesis of ¹⁸F-labeled tetrazines (Scheme 1). Our attempts to convert nitrotetrazine derivatives **2a** and **2b** into ¹⁸F-labeled substitution products **3** using ¹⁸F-fluoride/kryptofix or ¹⁸F-TBAF gave decomposition products and only traces of radiolabeled products. We also combined mesylate **4** with fluoride: in the most successful experiment (¹⁸F-TBAF at 85 °C for 15 min), ¹⁸F-labeled product **5** was obtained in ~1% labeling yield.

The instability of tetrazine precursors to F-labeling conditions prompted us to consider methods for preparing ¹⁸F-labeled *trans*-cyclooctenes. To this end, we synthesized the nosylate **8** as shown in Scheme 2. The key step in the synthesis was photoisomerization of **6** to **7** using a flow reactor that continuously removes the *trans*-isomers through selective metal complexation.⁸ The major diastereomer of **7** was carried forward in the synthesis of **8**.

Nosylate **8** reacted efficiently with TBAF to provide ${}^{19}F-9$ in high yield. With this standard in hand, conditions for the preparation of ${}^{18}F-9$ were optimized (Table 1). It was found that ${}^{18}F-9$ could be obtained in good yield by adding nosylate **8** to a mixture of [${}^{18}F$]-fluoride (100 mCi)/tetrabutyl-ammonium bicarbonate (TBAB) in acetonitrile (0.8 mL). We determined the effect of varying the concentration of **8** in reactions conducted for 15 min at 75 °C (Table 1, entries 1–4). The highest labeling yield (71%) was achieved with 7.0 mM **8**, but a useful radiochemical yield (18%) was still obtained with 0.35 mM **8**. Different reaction temperatures were studied, and 75 °C was found to be optimal (Table 1, entries 3, 5–7).

 Table 1
 Optimization of synthesis of ¹⁸F-labeled trans-cyclooctene

Finally, we investigated the efficiency of the ¹⁸F labeling as a function of time using 2 mg of **8** at 75 °C (Table 1, entries 8–10). While the labeling yield was optimal (71%) in an experiment conducted for 15 min, useful labeling yields were also obtained after 3 min (43%) and 7 min (68%). The conclusion of the experiments summarized in Table 1 is that the conditions of entry 3 (7.0 mM **8**, 75 °C, 15 min) are optimal for ¹⁸F labeling.

As derivatives of 3,6-di(2-pyridyl)-s-tetrazine (e.g. **1b**, Fig. 1) are readily accessible,^{6,10,12} 3,6-di(2-pyridyl)-s-tetrazine (**1a**) was used to test the efficiency of the ¹⁸F-labeled *trans*-cyclooctene **9** in the TTCO-ligation. A 'cold' study with ¹⁹F–**9** was initially conducted. As expected, the conjugate ¹⁹F–**10** formed immediately, and then slowly isomerized to 1,4-dihydropyrazine ¹⁹F–**11** as a mixture of isomers⁶ (Scheme 3a). These isotopically stable conjugates served as coinjection standards for analysis of reactions with ¹⁸F–**9**.

The conjugations between **1a** and 18 F-**9** were carried out in 1 : 1 acetonitrile/water, and were analyzed within 10 seconds of mixing. Prior to the conjugation, 18 F-**9** was purified by



Scheme 3 (a) Conjugation between 9 and 1a occurs rapidly to form conjugate 10, which slowly rearranges to 11 as a mixture of regio- and stereoisomers. (b) Radio-HPLC analysis of the crude product from Table 2, entry 3.

			H 9	
Entry	Amount of 8	$Temp/^{\circ}C$	Reaction time/min	Radiochemical yield (%)
1	100 μg (0.35 mM)	75	15	18
2	500 µg (1.8 mM)	75	15	25
3	2.0 mg (7.0 mM)	75	15	71
4	3.0 mg (11 mM)	75	15	71
5	2.0 mg (7.0 mM)	40	15	24
6	2.0 mg (7.0 mM)	55	15	34
7^a	2.0 mg (7.0 mM)	90	15	71
8	2.0 mg (7.0 mM)	75	3	43
9	2.0 mg (7.0 mM)	75	7	68
10	2.0 mg (7.0 mM)	75	30	71
^a Comparable	e results (70% RCY) were obtained	under similar conditions	using K ₂ CO ₃ /K ₂₂₂ instead of TBAE	3.

Entry	$^{18}\text{F-}9^{a}$	1a	Solvent	Reaction time/s	Radiochemical yield ($^{18}F-10 + ^{18}F-11$) (%)
1	1 mCi (2 μM)	210 µM	MeCN/H ₂ O	<10	> 98
2	$1 \text{ mCi} (2 \mu \text{M})$	21 µM	MeCN/H ₂ O	<10	>98
3	$0.1 \text{ mCi} (0.2 \mu \text{M})$	21 µM	MeCN/H ₂ O	<10	98
4	$0.1 \text{ mCi} (0.2 \mu \text{M})$	2.1 µM	MeCN/H ₂ O	<10	56
5	$0.1 \text{ mCi} (0.2 \mu \text{M})$	0.21 µM	MeCN/H ₂ O	<10	15
6	$1 \text{ mCi} (2 \mu M)$	21 µM	PBS buffer	<10	>98
7	$1 \text{ mCi} (2 \mu M)$	21 μM	Serum	<10	> 98

Table 2 The effect of concentration on the formation of 18 F-10 through the conjugation between 18 F-9 and 3,6-di(2-pyridyl)-s-tetrazine. All reactions were performed at room temperature

^{*a*} The concentration of 18 F–9 was estimated based on the specific activity of fluoride after bombardment (~4 Ci µmol⁻¹), taking into account a correction for the rate of radioactive decay.

HPLC and easily separated from unreacted precursor **8**. When ¹⁸F–**9** (1 mCi, 2 μ M) was combined with **1a** (concentrations of $\geq 21 \,\mu$ M), ¹⁸F–**9** was completely consumed within 10 s, and ¹⁸F–**10** had formed in 98% radiochemical yield, accompanied by ¹⁸F–**11** (1%) (Table 2, entries 1 and 2). When the concentration of ¹⁸F–**9** was decreased to 0.1 mCi (0.2 μ M), the conjugate ¹⁸F–**10/11** was still formed in excellent radiochemical yield (98%, entry 3). Useful radiochemical yields could also be obtained with even lower concentrations of **1a** (entries 4 and 5). We also investigated the efficiency of the conjugation between **1a** (21 μ M) and ¹⁸F–**9** (1 mCi, 2 μ M) in PBS buffer and serum media: ¹⁸F–**10** is formed in quantitative yield within 10 seconds (entries 6 and 7).

The Diels–Alder conjugate **10** was found to be stable in water, and the benign isomerization to **11** was the only side reaction. Thus, ¹⁹F–**10** was the only product detected by ¹H NMR analysis immediately after the conjugation. In CD₃CN/H₂O, the rearrangement of ¹⁹F–**10** to ¹⁹F–**11** proceeded to 11% conversion after 4 hours, and >95% conversion after 48 hours. The stability of the radiolabeled conjugation product was monitored in PBS buffer and serum media for 4 hours, and no degradation products of ¹⁸F–**10** were observed.

In conclusion, a new radiolabeling method for bioconjugation based on the TTCO-ligation has been described. The method utilizes a novel F-18 labeled *trans*-cyclooctene probe, and the efficiency of the reaction has been demonstrated with 3,6-di-(2-pyridyl)-*s*-tetrazine, which can be readily functionalized *via* amido derivatives. The reaction proceeds with exceptionally fast rates, making it an effective conjugation method at low concentration. We anticipate that the TTCO-ligation will provide the foundation for a new class of reliable methods for ¹⁸F-labeling of biomolecules in PET applications.

This work was supported by NIH Grant Number P20 RR017716 from the COBRE Program of the NCRR, and by the USC Department of Radiology. Spectra were obtained

with instrumentation supported by NSF CRIF:MU grants: CHE 0840401 and CHE-0541775.

Notes and references

- 1 D. J. Schlyer, Ann. Acad. Med., 2004, 33, 146-154.
- 2 R. Schirrmacher, C. Wängler and E. Schirrmacher, *Mini-Rev. Org. Chem.*, 2007, 4, 317–329.
- 3 P. H. Elsinga, *Methods*, 2002, **27**, 208–217; R. Schirrmacher, C. Wängler and E. Schirrmacher, *Eur. J. Org. Chem.*, 2008, 2853–2873.
- 4 J. Marik and J. L. Sutcliffe, *Tetrahedron Lett.*, 2006, 47, 6681–6684.
- 5 Z. B. Li, Z. Wu, K. Chen, F. T. Chin and X. Chen, *Bioconjugate Chem.*, 2007, **18**, 1987–1994.
- 6 M. L. Blackman, M. Royzen and J. M. Fox, J. Am. Chem. Soc., 2008, 130, 13518–13519.
- 7 For reviews of Cu-free bioconjugation chemistry: J. C. Jewett and C. R. Bertozzi, *Chem. Soc. Rev.*, 2010, **39**, 1272–1279; C. R. Becer, R. Hoogenboom and U. S. Schubert, *Angew. Chem., Int. Ed.*, 2009, **48**, 4900–4908; Reviews of bioorthogonal chemistry: R. K. V. Lim and Q. Lin, *Chem. Commun.*, 2010, **46**, 1589; J. Kalia and R. T. Raines, *Curr. Org. Chem.*, 2010, **14**, 138–147; E. M. Sletten and C. R. Bertozzi, *Angew. Chem., Int. Ed.*, 2009, **48**, 6974.
- 8 M. Royzen, G. P. A. Yap and J. M. Fox, J. Am. Chem. Soc., 2008, 130, 3760–3761.
- 9 (a) F. Thalhammer, U. Wallfahrer and J. Sauer, *Tetrahedron Lett.*, 1990, **31**, 6851; (b) D. L. Boger, *Chem. Rev.*, 1986, **86**, 781.
- 10 Acetamido-substituted dipyridyltetrazines (1b) display fast kinetics,¹² and are stable for prolonged periods (>24 h) in water, and toward amines and thiols in MeOH: R. Selvaraj, M. L. Blackman, J. M. Fox, unpublished results.
- (a) N. K. Devaraj, R. Weissleder and S. A. Hilderbrand, Bioconjugate Chem., 2008, 19, 2297-2299; (b) R. Pipkorn, W. Waldeck, B. Didinger, M. Koch, G. Mueller, M. Wiessler and K. Braun, J. Pept. Sci., 2010, 15, 235-241; (c) N. K. Devaraj, R. Upadhyay, J. B. Haun, S. A. Hilderbrand and R. Weissleder, Angew. Chem., Int. Ed., 2009, 48, 7013; (d) N. K. Devaraj, S. Hilderbrand, R. Upadhyay, R. Mazitschek and R. Weissleder, Angew. Chem., Int. Ed., 2010, 19, 2869-2872.
- 12 R. Rossin, P. R. Verkerk, S. M. v. d. Bosch, R. C. M. Vulders, I. Verel, J. Lub and M. S. Robillard, *Angew. Chem., Int. Ed.*, 2010, 49, 3375–3378.
- 13 G. Vaidyanathan and M. R. Zalutsky, Nat. Protoc., 2006, 1, 1655–1661.