

# **CHEMISTRY** A European Journal



# **Accepted Article** Title: Enantioselective Catalytic (4+1)-Cyclization of ortho-Hydroxypara-Quinone Methides with Allenoates Authors: Katharina Zielke, Ondřej Kováč, Michael Winter, Jiří Pospíšil, and Mario Waser This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article. To be cited as: Chem. Eur. J. 10.1002/chem.201901784 Link to VoR: http://dx.doi.org/10.1002/chem.201901784 **Supported by**

ACES

WILEY-VCH

## Enantioselective Catalytic (4+1)-Cyclization of ortho-Hydroxypara-Quinone Methides with Allenoates

Katharina Zielke,<sup>#[a]</sup> Ondřej Kováč,<sup>#[b]</sup> Michael Winter,<sup>[a]</sup> Jiří Pospíšil,<sup>[b,c]</sup> and Mario Waser<sup>\*[a]</sup>

**Abstract:** We herein report the first highly asymmetric catalytic synthesis of densely functionalized dihydrobenzofurans starting from ortho-hydroxy-containing para-quinone methides. The reaction relies on an unprecedented formal (4+1)-annulation of these quinone methides with allenoates in the presence of a commercially available chiral phosphine catalyst. The hereby accessed chiral dihydrobenzofurans were obtained as single diastereomers in yields up to 90% and with enantiomeric ratios up to 95:5.

## Introduction

The 2,3-dihydrobenzofuran scaffold is a prominent structural motif found in numerous biologically active (natural) compounds<sup>[1]</sup> and the development of novel synthesis strategies to access these targets has been a heavily investigated topic over the last years.<sup>[2-</sup> <sup>8]</sup> One especially appealing approach to access chiral 2,3dihydrobenzofurans is the formal (4+1)-cyclization<sup>[9]</sup> between a suitable C1 building block and a carefully chosen (maybe in situ generated) acceptor-donor containing C4 building block. The most versatile class of C4 building blocks used to obtain the dihydrobenzofuran skeleton 1 via a formal (4+1)-cyclization are ortho-quinone methides (o-QMs) 2.[10] These, usually in situ generated, reactive compounds have recently been very successfully used for racemic as well as highly stereoselective (4+1)-annulations with either sulfonium or ammonium ylides, <sup>[4]</sup>  $\alpha$ halocarbonyl compounds,<sup>[5]</sup> or with diazocompounds as C1 synthons (Scheme 1A).<sup>[6]</sup> Alternatively, the hydroxy-containing para-quinone methides 3 have very recently emerged as powerful building blocks for formal (4+n)-annulations as well.[7,11-13] Interestingly however, their applicability for asymmetric (4+1)cyclizations to access dihydrobenzofurans 1 has so far been rather limited, with highly asymmetric protocols still being rare (Scheme 1B).<sup>[7]</sup> Two years ago we reported the first highly enantioselective synthesis of compounds 1 by reacting preformed chiral ammonium ylides with in situ formed o-QMs 2.[4c] More recently we found that the highly functionalized allenoates 4 can undergo a very unique (and up to then unprecedented) formal

[a]	Dl <sup>in</sup> K. Zielke, DI M. Winter, Prof. M. Waser Johannes Kepler University Linz, Institute of Organic Chemistry
	Altenbergerstraße 69, 4040 Linz, Austria
	E-mail: mario.waser@jku.at
[b]	Mgr. O. Kováč, Dr. J. Pospíšil
	Department of Organic Chemistry,
	Faculty of Science, Palacky University,
	tř. 17. listopadu 1192/12, CZ-771 46 Olomouc, Czech Republic
[c]	Dr. J. Pospíšil
	Laboratory of Growth Regulators,
	The Czech Academy of Sciences, Institute of Experimental Botany
	& Palacký University
	Šlechtitelů 27, CZ-78371 Olomouc, Czech Republic
#	These authors contributed equally

Supporting information for this article is given via a link at the end of the document.((Please delete this text if not appropriate))

(4+1)-cyclization with acceptors 2 in the presence of a stoichiometric amount of PPh3 (Scheme 1C).<sup>[8]</sup> The (unexpected) outcome of this reaction was in sharp contrast to other previously described reactions between o-QMs 2 and (differently substituted) allenoates, which all resulted in formal (4+2)annulations.[14] Unfortunately however, we were only able to carry out this reaction in racemic manner, as even the use of a stoichiometric amount of different commonly used chiral phosphine catalysts gave low yields and poor enantioselectivities only.<sup>[8]</sup> In especially we found that the in situ formed o-QMs 2 decomposed rather rapidly under the previously developed reaction conditions, thus making a catalytic approach difficult. Given these limitations in catalyst turnover and asymmetric induction, we thought that maybe an alternative and slightly more stable acceptor would be beneficial to address these challenges. Thus, we decided to investigate if this methodology may also be extended to the formal (4+1)-cyclization of the o-hydroxycontaining p-QMs 3 with allenoates 4.[12] We reasoned that this preformed and, compared to 2, more stable acceptor molecule may be better suited to establish a truly catalytic as well as asymmetric protocol, which would then allow for the first enantioselective (4+1)-annulation of p-QMs 3 to access the highly functionalized chiral dihydrobenzofurans 5 (Scheme 1D).



B. Recently reported (4+1)-cyclizations of p-QMs 3:[7]



C. Our allenoate 4-based (4+1)-cyclization of o-QMs 2:[8]

4



catalytic asymmetric protocol

protocols

**Scheme 1.** Previous (4+1)-approaches for the syntheses of dihydrobenzofurans (A-C) and the herein reported novel catalytic asymmetric strategy (D).

## **Results and Discussion**

#### Initial Optimization of the Racemic Reaction.

We started our investigations by carrying out the racemic reaction between p-QM 3a and diethyl allenoate 4a in the presence of PPh<sub>3</sub> (Table 1 gives an overview of the most significant screening results). Our first reactions were carried out in analogy to the conditions developed for the annulation of o-QMs 2<sup>[8]</sup> (please note that we previously used a twofold excess of the guinone methide 2 to compensate for its competing decomposition under the reaction conditions). Gratifyingly the targeted dihydrobenzofuran 5a could be obtained as a single diastereomer in this initial attempt already (entry 1). The relative configuration of the product 5a was confirmed by NOESY experiments (as shown in Scheme 3) and we also observed the same correlations for other products 5 later (Scheme 2). As the reaction was found to be rather slow, with significant amounts of unreacted 3a being recovered (indicating its increased stability compared to o-QMs 2), we next increased the amount of base (entry 2), which however had a detrimental effect (complete decomposition of starting materials).

Table 1: Initial screening with PPh3.



Entry <sup>a</sup>	3a : 4a	PPh₃	Base (eq.)	Solvent	Yield (%) <sup>b</sup>
1	2 : 1	1 eq.	Cs <sub>2</sub> CO <sub>3</sub> (2 eq.)	CH <sub>2</sub> Cl <sub>2</sub>	16°
2	2 : 1	1 eq.	Cs <sub>2</sub> CO <sub>3</sub> (10 eq.)	CH <sub>2</sub> Cl <sub>2</sub>	-
3	1:2	1 eq.	Cs <sub>2</sub> CO <sub>3</sub> (2 eq.)	CH <sub>2</sub> Cl <sub>2</sub>	33°
4	1:2	1 eq.	Cs <sub>2</sub> CO <sub>3</sub> (2 eq.)	Toluene	45 <sup>d</sup>
5	1:2	1 eq.	Cs <sub>2</sub> CO <sub>3</sub> (2 eq.)	EtOAc	-
6	1:2	1 eq.	Cs <sub>2</sub> CO <sub>3</sub> (0.2 eq.)	CH <sub>2</sub> Cl <sub>2</sub>	63
7	1:2	1 eq.	K <sub>2</sub> CO <sub>3</sub> (2 eq.)	CH <sub>2</sub> Cl <sub>2</sub>	82
8	1:2	0.2 eq.	K <sub>2</sub> CO <sub>3</sub> (2 eq.)	CH <sub>2</sub> Cl <sub>2</sub>	81
9	1:2	0.1 eq.	K <sub>2</sub> CO <sub>3</sub> (2 eq.)	CH <sub>2</sub> Cl <sub>2</sub>	49
10	1:2	0.2 eq.	K <sub>2</sub> CO <sub>3</sub> (0.5 eq.)	CH <sub>2</sub> Cl <sub>2</sub>	89
11	1:2	0.2 eq.		CH <sub>2</sub> Cl <sub>2</sub>	90 (86) <sup>e</sup>

[a] All reactions were carried out on 0.1 mmol scale (based on the p-QM 3a):[b] Isolated yield; [c] Incomplete conversion of 3a and noticeable amounts of

side products; [d] complete conversion of **3a** and large amounts of unidentified side products; [e] 1 mmol scale reaction.

As decomposition of the acceptor **3a** was obviously not very fast in the first attempts with 2 equiv. of base we next used an excess of allenoate **4a**, which lead to a measurable increase in yield (entry 3). The screening of different solvents revealed that toluene allows for a slightly higher yield (entry 4), but also accompanied with a more pronounced formation of various not identified sideor decomposition products. Other solvents were not satisfying (see entry 5 for one example) and we thus carried out further optimizations with CH<sub>2</sub>Cl<sub>2</sub>. Very interestingly, lowering the amount of base (entry 6) significantly improved the yield and suppressed the side product formation. By testing other bases next, K<sub>2</sub>CO<sub>3</sub> turned out to be the most promising (entry 7). It should be noted that other simple trialkylphosphines were tested as well,<sup>[15]</sup> but in analogy to our previous observation<sup>[8]</sup> these did not allow for this (4+1)-annulation.

With these first high yielding conditions set, we next lowered the amount of PPh<sub>3</sub>. Gratifyingly, and in sharp contrast to the reaction with o-QMs **2**,<sup>[8]</sup> the use of 20 mol% PPh<sub>3</sub> allowed for the same yield as when using a stoichiometric amount (compare entries 7 and 8). Further lowering of the catalyst amount unfortunately slowed down the reaction measurably (entry 9). Considering the beneficial effect of using less base when using Cs<sub>2</sub>CO<sub>3</sub> (entry 6), we finally also lowered the amount of K<sub>2</sub>CO<sub>3</sub> (entries 10, 11), and much to our surprise the reaction proceeded well even without any base (entry 11; the reaction was reproduced several times on different scales and also on 1 mmol scale).

#### Development of an Asymmetric Catalytic Protocol.

Having established high yielding and robust catalytic procedures for the racemic synthesis of 5a we next focused on the use of chiral phosphine catalysts. As already mentioned before, we were not able to identify a suited asymmetric catalyst for our previous (4+1)-annulation of o-QMs 2. However, given the fact that p-QM 3a performed very well in the racemic reaction and also allowed for a catalytic approach, we were confident that the well-described bulky chiral phosphines A<sup>[16]</sup> or B<sup>[17]</sup> may allow for a truly catalytic enantioselective protocol (Table 2). We first used the binaphthylbased phosphines A1-3, but unfortunately neither of them allowed for any product formation (entries 1-3). Gratifyingly however, by switching to the commercially available chiral spiro phosphine B ((R)-SITCP)[17] we observed a very clean and reasonably enantioselective product formation when using 20 mol% of this catalyst under base-free conditions in CH<sub>2</sub>Cl<sub>2</sub> (entry 4). Lowering the reaction temperature to 0 °C unfortunately did not allow for product formation anymore (entry 5). When carrying out the reaction in the presence of two equiv. of K<sub>2</sub>CO<sub>3</sub>, the outcome was only slightly affected in this solvent (entry 6).

## WILEY-VCH

# **FULL PAPER**

Table 2: Development of an enantioselective catalytic protocol.



Entry <sup>a</sup>	PR₃ (mol%)	Solvent	Base (eq.)	Temp. (°C)	Yield (%) <sup>b</sup>	e.r.º
1	<b>A1</b> (20%)	CH <sub>2</sub> Cl <sub>2</sub>	-	25	-	-
2	<b>A2</b> (20%)	$CH_2CI_2$	-	25	-	-
3	<b>A3</b> (20%)	CH <sub>2</sub> Cl <sub>2</sub>	-	25	-	-
4	<b>B</b> (20%)	CH <sub>2</sub> Cl <sub>2</sub>	-	25	84	88:12
5	<b>B</b> (20%)	CH <sub>2</sub> Cl <sub>2</sub>	-	0	-	-
6	<b>B</b> (20%)	CH <sub>2</sub> Cl <sub>2</sub>	K₂CO₃ (2 eq.)	25	70	87:13
7	<b>B</b> (20%)	Toluene	-	25	87	94:6
8	<b>B</b> (20%)	Toluene	K <sub>2</sub> CO <sub>3</sub> (2 eq.)	25	90	92:8
9	<b>B</b> (20%)	Toluene	K₂CO₃ (2 eq.)	10	89	94:6
10	<b>B</b> (20%)	Toluene	K <sub>2</sub> CO <sub>3</sub> (2 eq.)	0	-	
11	<b>B</b> (10%)	Toluene	K₂CO₃ (2 eq.)	10	-	-

[a] All reactions were carried out on 0.1 mmol scale (based on **3a**): [b] Isolated yield; [c] Determined by HPLC using a chiral stationary phase with the (+)-enantiomer as the major product. The relative configuration was assigned by NOESY experiments (see Scheme 4) but the absolute configuration could not be determined yet.

Interestingly, when changing to toluene (other solvents like THF were found to be not suited), we were able to improve the enantioselectivity significantly (entries 7-9). At room temperature reactions in the presence of  $K_2CO_3$  as well as under base-free conditions performed very similarly, with a slightly higher e.r. in the absence of base (compare entries 7 and 8). However, when we investigated the application scope later on, we realized that the base-mediated conditions were more robust when using differently substituted starting materials **3**, while not all of those allowed for good conversions under base-free conditions. Other bases were found to be less satisfactory (with e.g.  $Cs_2CO_3$  giving

lower yields and K<sub>3</sub>PO<sub>4</sub> giving no product at all). We thus tested if any further improvement in the presence of 2 equiv. of K<sub>2</sub>CO<sub>3</sub> may be possible (entries 8-11). However, lowering of the reaction temperature was possible to some extent only (entries 9, 10), but reducing the catalyst loading to 10 mol% was unfortunately not possible anymore (entry 11). Accordingly, the best-suited and most robust catalytic enantioselective approach to access **5a** as a single diastereomer was to carry out the reaction in toluene at 10 °C in the presence of 2 equiv. K<sub>2</sub>CO<sub>3</sub> by using 20 mol% of the commercially available phosphine catalyst **B** (entry 9, the reaction was reproduced by different persons on 0.05 - 0.1 mmol **3a** scale giving identical results).

#### Application Scope.

CI

Having established a high yielding and robust catalytic procedure for the synthesis of dihydrobenzofuran **5a**, we next tested the use of differently substituted quinone methides **3** and allenoates **4** (Scheme 2).



 $\begin{array}{l} \label{eq:scheme 2. Application scope (Conditions: 0.05 - 0.2 mmol 3, 2 equiv. 4, 2 equiv. \\ \ensuremath{K_2CO_3}, 20 mol\% \ensuremath{B}, toluene (0.025 \ensuremath{M}), 10 \ensuremath{^\circ C}, min. 20 \ensuremath{h}; \ensuremath{All yields are isolated yields and enantiomeric ratios were determined by HPLC using a chiral stationary phase with the (+)-enantiomer being the major product in each case). \\ \end{array}$ 

First, we could show that replacement of one of the allenoate ethyl ester groups for a benzyl ester was tolerated very well (see product **5b**). Then it turned out that a dimethyl-based p-QM **3** can be used as well to obtain the enantioenriched product **5c** (albeit with a slightly lower selectivity than for the parent t-Bu-based **5a**). Interestingly, substituents in the 5 and 6-position of the

benzofuran backbone were tolerated very well (see compounds 5d, 5e, 5f, 5i, 5j, 5k, 5m). In contrast, substituents in positions 4 and 8 turned out to be more limiting and product 5g was only accessible with a rather low enantiomeric ratio of 79:21. Surprisingly, compound 5h was not formed at all under the asymmetric conditions (even with longer reaction times). We were however able to obtain racemic 5h in high yield when using PPh<sub>3</sub> as an achiral catalyst. Very interestingly, while we found initially that benzyl ester containing allenoates were tolerated similarly well as ethyl ester-based ones (see targets 5a and 5b), we found that t-butyl esters resulted in somewhat lower enantiomeric ratios compared to ethyl and benzyl esters (compare 5k and 5l as well as 5m, 5n, and 5o). All asymmetric reactions were initially carried out on 0.05 mmol scale of the limiting agent 3 and we also reproduced selected reactions on up to 0.2 mmol scale without affecting the outcome, thus indicating that the asymmetric procedure is of similar robustness as the racemic one (Table 1, entry 11).

All substrate combinations gave the (+)-enantiomer as the major product, but unfortunately we were so far not able to obtain suited crystals of any of the products **5** to determine the absolute configuration by means of single crystal X-ray analysis.

It has been described by others that the t-butyl groups of the phenol derivatives obtained by addition of nucleophiles to QMs can be cleaved off under (Lewis) acidic conditions.<sup>[7a,11c]</sup> We thus carried out a few (unoptimized) test reactions to see if a similar debutylation is also possible on the highly functionalized diestercontaining dihydrobenzofuranes 5 (Scheme 3). Carrying out the reaction at elevated temperature lead to guick decomposition only. In contrast, at room temperature the slow formation of the debutylated diester 6a was observed by MS. Interestingly however, the major reaction product was found to be the debutylated monoester 7a that was formed in around 30% after one day and around 50-60% after 3 days (accompanied with some decomposition products) and which could also be isolated after column chromatography (NMR clearly confirmed that the ester group on the stereogenic center was hydrolysed). It should be noted that no further attempts to optimize this reaction were undertaken, but this result clearly shows that the highly functionalized compounds 5a can be used for further transformations and that the two ester groups have different reactivities.



#### Mechanistic Considerations.

Mechanistically this is a rather complex reaction and it should be admitted that so far we only have some hints that may allow us to postulate the mechanistic scenario depicted in Scheme 4. This proposal is also based on our recent observations made for the racemic (4+1)-annulation of o-QMs **2** where we found that intramolecular rapid proton transfers are crucial to explain the outcome of this (4+1)-cyclization.<sup>[8]</sup>



Scheme 4. Postulated mechanism and NOESY results to determine the relative configuration.

Addition of the phosphine to the allenoate is supposed to give the required zwitterion I after proton transfer on the primary addition product. Following the reaction between PPh<sub>3</sub> and **4a** by <sup>31</sup>P NMR shows the appearance of two new signals around 27 ppm (the parent PPh<sub>3</sub> peak is at -5 ppm) substantiating the formation of alkylated phosphine species (these addition products decomposed very quickly in the absence of any electrophile). Upon addition of the guinone methide 3a immediately a strong red colour evolves, which can be rationalized by the 1.6-addition of I to 3a to give the phenolate II. Similar colour changes can also be observed when adding different nucleophiles to other p-QMs, substantiating the assumed initial 1,6-addition. With respect to the nature of the electrophile 3a one could however also postulate that a prototropic shift from the phenol to the para-QM moiety gives an ortho-QM in situ, which then reacts with I to give III directly).<sup>[18]</sup> However, we found compound **3a** being rather stable under basic conditions and we never observed any other species or got any experimental hint that supports this pathway, but it should not be ruled out completely with the current state of knowledge. The phenolate II then needs to undergo two proton transfer reactions towards the betaine IV, which then finally can react to the product 5a via an  $S_N2$ ' type cyclization. We have recently shown for the cyclizations of o-QMs 2 that these proton transfers are rather likely processes and we reason that the

presence of a base is beneficial for these reactions, which would be an explanation why the herein presented (4+1)-cyclization is more robust under basic conditions. This beneficial effect of base became especially pronounced in those cases where no electrondonating ring substituent para to the OH group is present (these reactions usually proceeded a bit slower as well). This observation supports a scenario where the final ring-closure may be the rate-determining step, which also rationalizes why slightly larger amounts of catalyst were necessary to obtain satisfying catalyst turnover.

With respect to the observed stereoselectivity it is likely that the catalyst controls the absolute configuration in the 1,6-addition step. An alternative may be a less selective 1,6-addition followed by base-mediated isomerization of the benzylic position on one of the chiral catalyst-bound intermediates II or III. However, as the observed enantioselectivity was more or less the same under basic and base-free conditions (compare with Table 2), this option seems less likely. The diastereoselectivity is then controlled in the final proton transfer - cyclization sequence. Given the fact that S<sub>N</sub>2' reactions usually proceed via cis-orientation of nucleophile and leaving group<sup>[19]</sup> the proton transfer towards **IV** is supposed to be highly selective, and may be steered by electrostatic attraction between the phenolate anion and the phosphonium cation in the nonpolar reaction solvent. However, it should clearly be pointed out that this is just a mechanistic hypothesis and although we were able to observe the presence of some alkylated phosphonium species by <sup>31</sup>P NMR during the reaction, none of these intermediates could be isolated or more carefully analysed.

## Conclusions

The first highly asymmetric catalytic formal (4+1)-annulation of ohydroxy-p-quinone methides **3** with allenoates **4** was developed. The outcome of this reaction is in sharp contrast to other recently reported reactions between quinone methides **3** and allenoates.<sup>[12]</sup> Key to success was the use of the commercially available chiral phosphine **B** as a catalyst under carefully optimized reaction conditions. This methodology allowed for the so far unprecedented synthesis of the chiral dihydrobenzofurans **5** as single diastereomers in yields up to 90% and with enantiomeric ratios up to 95:5.

## **Experimental Section**

General details can be found in the online supporting information. This document also contains detailed synthesis procedures and analytical data of novel compounds and reaction products as well as copies of NMR spectra and HPLC traces.

**General asymmetric (4+1)-cyclization procedure:** A mixture of the paraquinone methide **3** (0.05-0.2 mmol),  $K_2CO_3$  (2 equiv.), and chiral phosphine **B** (20 mol%) was cooled to 10°C and a solution of the allenoate **4** (2 equiv.) in dry toluene (20 ml per mmol **4**) was added. The resulting mixture was stirred at 10°C under an Ar atmosphere for approximately 20 h. The mixture was diluted by adding CH<sub>2</sub>Cl<sub>2</sub> (5 ml), filtrated over a pad of Na<sub>2</sub>SO<sub>4</sub> and the residue was rinsed with CH<sub>2</sub>Cl<sub>2</sub> (5 x 5 ml). The combined organic layers were evaporated to dryness (under reduce pressure) and the products were purified by silica gel column chromatography (gradient of heptanes and EtOAc) giving the corresponding dihydrobenzofurans **5** in the reported yields and enantiopurities (Syntheses of racemic samples were carried out in analogy using PPh<sub>3</sub> instead).

**Dihydrobenzofuran 5a.** Obtained as a yellow residue in 89% and e.r. = 94:6.  $[α]_D^{23} = 64.6$  (c = 0.15, CHCl<sub>3</sub>, e.r. = 94:6); 1H NMR (300 MHz, δ, CDCl<sub>3</sub>, 298 K): 0.82 (t, *J* = 7.2 Hz, 3H), 1.36 (t, *J* = 7.1 Hz, 3H), 1.36 (s, 18H), 1.89 (d, *J* = 7.2 Hz, 3H), 3.52-3.77 (m, 2H), 4.26-4.36 (m, 2H), 5.11 (s, 1H), 5.24 (s, 1H), 6.40 (q, *J* = 7.1 Hz, 1H), 6.83 (s, 2H), 6.94 (t, *J* = 7.3 Hz, 1H), 7.01-7.09 (m, 2H), 7.19-7.26 (m, 1H) ppm. <sup>13</sup>C NMR (75 MHz, δ, CDCl<sub>3</sub>, 298 K): 13.4, 14.2, 15.5, 30.2, 34.2, 55.8, 60.8, 61.1, 94.0, 110.2, 121.8, 125.7, 126.1, 128.9, 129.4, 130.0, 132.7, 133.2, 135.1, 153.0, 158.1, 167.2, 168.4 ppm; HRMS (ESI): m/z calcd for C<sub>31</sub>H<sub>40</sub>O<sub>6</sub>: 509.2898 [M+H]+; found: 509.2897. The enantioselectivity was determined by HPLC (YMC Chiral Art Cellulose-SB, eluent: hexane:*i*-PrOH = 95:5, 0.5 mL/min, 10 °C, retention times: t<sub>major</sub> = 9.4 min, t<sub>minor</sub> = 11.0 min).

## Acknowledgements

This work was supported by the Austrian Science Funds (FWF): Project No. P26387-N28. The NMR spectrometers used were acquired in collaboration with the University of South Bohemia (CZ) with financial support from the European Union through the EFRE INTERREG IV ETC-AT-CZ program (project M00146, "RERI-uasb"). O. Kováč was supported from Internal Grant Agency of Palacky University (IGA\_PrF\_2019\_027). J. Pospíšil was supported from European Regional Development Fund Project "Centre for Experimental Plant Biology" (No. CZ.02.1.01/0.0/0.0/16\_019/0000738).

**Keywords:** Allenoates • Enantioselectivity • Diastereoselectivity • Organocatalysis • Annulation

- For illustrative reviews and selected examples please see: (a) R. S. Ward, Nat. Prod. Rep., 1999, 16, 75-96; (b) S. Apers, D. Paper, J. Bürgermeister, S. Baronikova, S. Vam Dyck, G. Lemiere, A. Vlietinck, L. Pieters, J. Nat. Prod., 2002, 65, 718-720; (c) S- Apers, A. Vlietinck, L. Pieters, Phytochem. Rev. 2003, 2, 201-207; (d) T. She, X-N. Wang, H.-X. Lou, Nat. Prod. Rep. 2009, 26, 916-935; (e) A. Radadiya, A. Shah, Eur. J. Med. Chem., 2015, 97, 356-376; (f) H. K. Shamsuzzaman, Eur. J. Med. Chem., 2015, 97, 483-504; (g) R. J. Nevagi, S. N. Dighe, S. N. Dighe, Eur. J. Med. Chem. 2015, 97, 561-581.
- (a) F. Bertolini, M. Pineschi, Org. Prep. Proced. Int. 2009, 41, 385-418; (b) T.
   D. Sheppard, J. Chem. Res. 2011, 35, 377-385.
- For selected recent examples employing complementary synthesis strategies see: (a) E. D. Coy, L. Jovanovic, M. Sefkow, Org. Lett., 2010, 12, 1976-1979; (b) J. Mangas-Sanchez, E. Busto, V. Gotor-Fernandez, V. Gotor, Org. Lett., 2010, 12, 3498-3501; (c) F. Baragona, T. Lomberget, C. Duchamp, N. Henriques, E. L. Piccolo, P. Diana, A. Montalbano, R. Barret, Tetrahedron, 2011, 67, 8731-8739; (d) J. Fischer, P. G. Savage, M. J. Coster, Org. Lett. 2011, 13, 3376-3379; (e) A. Lu, K. Hu, Y. Wang, H. Song, Z. Zhou, J. Fang, C. Tang, J. Org. Chem., 2012, 77, 6208-6214; (f) N. Ortega, S. Urban, B. Beiring, F. Glorius, Angew. Chem. Int. Ed. 2012, 51, 1710-1713; (g) A. k. Shaikh, G. Varvounis, RSCAdv. 2015, 5, 14892-14896; (h) S. Sharma, S. K. R. Parumala, R. K. Peddinti, Synlett, 2016, 28, 239-244; (i) Y. Cheng, Z. Gang, W. Li, P. Li, Org. Chem. Front. 2018, 5, 2728-

## WILEY-VCH

# **FULL PAPER**

2733; (j) T. Zhou, T. Xia, Z. Liu, L. Liu, J. Zhang, Adv. Synth. Catal. 2018, 360, 4475-4479.

- For ylide-based (4+1) cyclizations of ortho-quinone methides: (a) M.-W. Chen,
  L.-L. Cao, Z.-S. Ye, G.-F. Jiang, Y.-G. Zhou, *Chem. Commun.*, 2013, 49, 1660-1662; (b) V. A. Osyanin, D. V. Osipov, Y. N. Klimochkin, *J. Org. Chem.*, 2013, 78, 5505-5520; (c) N. Meisinger, L. Roiser, U. Monkowius, M. Himmelsbach, R. Robiette, M. Waser, *Chem. Eur. J.* 2017, 23, 5137-5142; (d) Q.-Q. Yang, W.-J. Xiao, *Eur. J. Org. Chem.* 2017, 233-236; (e) Y.-Q. Liu, Q.-Z. Li, H.-P. Zhu, X. Feng, X. Peng, W. Huang, J.-L. Li, B. Han, *J. Org. Chem.* 2018, 83, 12753-12762;
- For a-halocarbonyl-based (4+1) cyclizations with ortho-quinone methides: (a)
   X.-L. Lian, A. Adili, B. Liu, Z.-L. Tao, Z.-Y. Han, Org. Biomol. Chem. 2017, 15, 3670-3673; (b)
   X.-L. Jiang; S.-J. Liu, Y.-Q. Gu, G.-J. Mei, F. Shi, Adv. Synth. Catal. 2017, 359, 3341-3346;
- 6 For a phosphoric acid-catalyzed (4+1) cyclization of ortho-quinone methides: A. Suneja, C. Schneider, *Org. Lett.* **2018**, *20*, 7576-7580.
- For para-quinone methide based (4+1) cyclizations: (a) L. Liu, Z. Yuan, R. Pan, Y. Zeng, A. Lin, H. Yao, Y. Huang, Org. Chem. Front. 2018, 5, 623-628; (b) Y. Zhi, K. Zhao, C. von Essen, K. Rissanen, D. Enders, Org. Chem. Front. 2018, 5, 1348-1351; (c) Y.-J. Xiong, S.-Q. Shi, W.-J. Hao, S.-J. Tu, B. Jiang, Org. Chem. Front. 2018, 5, 3483-3487; (d) J. Zhou, G. Liang, X. Hu, L. Zhou, H. Zhou, Tetrahedron 2018, 74, 1492-1496.
- 8 K. Zielke, M. Waser, Org. Lett. 2018, 20, 768-771.
- 9 For reviews on (4 + 1)-cyclizations: (a) C. Zhu, Y. Ding, L.-W. Ye, Org. Biomol. Chem., 2015, 13, 2530-2536; (b) J.-R. Chen, X.-Q. Hu, L.-A. Lu, W.-J. Xiao, Chem. Rev., 2015, 115, 5301-5365.
- For reviews on quinone methides: (a) T. P. Pathak, M. S. Sigman, J. Org. Chem.
   2011, 76, 9210-9215; (b) W.-J. Bai, J.-G. David, Z.-G. Feng, M. G. Weaver, K.-L. Wu, T. R. R. Pettus, Acc. Chem. Res. 2014, 47, 3655-3664; (c) N. J. Willis, C. D. Bray, Chem. Eur. J. 2012, 18, 9160-9173; (d) M. S. Singh, A. Nagaraju, N. Anand, S. Chowdhury, RSC Adv. 2014, 4, 55924-55959; (e) L. Caruana, M. Fochi, L. Bernardi, Molecules 2015, 20, 11733-11764; (f) A. Parra, M. Tortosa, ChemCatChem 2015, 7, 1524-1526; (g) D. V. Osipov, V. A. Osyanin, Y. N. Klimochkin, Russ. Chem. Rev. 2017, 86, 625-687; (h) C. D. T. Nielsen, H. Abas, A. C. Spivey, Synthesis, 2018, 50, 4008-4018.

- For recent (4+2)-cyclizations see: (a) K. Zhao, Y. Zhi, T. Shu, A. Valkonen, K. Rissanen, D. Enders, *Angew. Chem. Int. Ed.* 2016, *55*, 12104-12108; (b) C. Duan, L. Ye, W. Xu, X. Li, F. Chen, Z. Zhao, X. Li, *Chin. Chem. Lett.* 2018, *29*, 1273-1276; (c) G.-J. Mei, S.-L. Xu, W.-Q. Zheng, C.-Y. Bian, F. Shi, *J. Org. Chem.* 2018, *83*, 1414-1421.
- 12 For two very recent (4+2) cyclizations of allenoates with p-QMs 3: (a) F.-R. Yuan, F. Jiang, K.-W. Chen, G.-J. Mei, Q. Wu, F. Shi, Org. Biomol. Chem. 2019, 17, 2361-2369; (b) Y. Zhu, D. Wang, Y. Huang, Org. Lett. 2019, 21, 908-912.
- (4+3): (a) F. Jiang, F.-R. Yuan, L.-W. Jin, G.-J. Mei, F. Shi, ACS Catal. 2018, 8, 10234-10240; (b) W. Li, H. Yuan, Z. Liu, Z. Zhang, Y. Cheng, P. Li, Adv. Synth. Catal. 2018, 360, 2460-2464. (c) Q. Liu, S. Li, X.-Y. Chen, K. Rissanen, D. Enders, Org. Lett. 2018, 20, 3622–3626.
- (a) P. Chen, K. Wang, W. Guo, X. Liu, Y. Liu, C. Li, Angew. Chem. Int. Ed. 2017, 56, 3689-3693; (b) Z. Wang, T. Wang, W. Yao, Y. Lu, Org. Lett. 2017, 19, 4126-4129; (c) Y.-H. Deng, W.-D. Chu, X.-Z. Zhang, X. Yan, K.-Y. Yu, L.-L. Yang, H. Huang, C.-A. Fan, J. Org. Chem. 2017, 82, 5433-5440.
- 15 For a recent report using PBu<sub>3</sub> to activate allenoates 4 see: A. Eitzinger, K. Zielke, M. Widhalm, R. Robiette, M. Waser, Asian J. Org. Chem. 2018, 7, 1620-1625.
- 16 Prepared according to: S. Enthaler, G. Erre, K. Junge, J. Holz, A. Borner, E. Alberico, I. Nieddu, S. Gladiali, M. Beller; Org. Proc. Res. Dev. 2007, 11, 568-577.
- 17 For a seminal contribution on this now commercially available phosphine: S.-F. Zhu, Y. Yang, L.-X. Wang, B. Liu, Q.-L. Zhou, *Org. Lett.* **2005**, *7*, 2333-2335.
- 18 It was recently shown that this isomerization is energetically possible: Z.-P. Zhang, K.-X. Xie, C. Yang, M. Li, X. Li, *J. Org. Chem.* **2018**, *83*, 364-373.
- For pioneering studies: (a) G. Stork, W. N. White, J. Am. Chem. Soc. 1953, 75, 4119-4120; (b) G. Stork, W. N. White, J. Am. Chem. Soc. 1956, 78, 4609-4619.

## WILEY-VCH

## Entry for the Table of Contents (Please choose one layout)

Layout 1:

## FULL PAPER

**FULL PAPER** 





Katharina Zielke, Ondřej Kováč, Michael Winter, Jiří Pospíšil and Mario Waser\*

Page No. – Page No.

Enantioselective Catalytic (4+1)-Cyclization of ortho-Hydroxy-para-Quinone Methides with Allenoates

Layout 2:

# FULL PAPER

Text for Table of Contents

((Insert TOC Graphic here; max. width: 11.5 cm; max. height: 2.5 cm))

Author(s), Corresponding Author(s)\*

Page No. – Page No. Title uthor(s)\*