

## A Molecular Platform for Design and Synthesis of Targeted Dual-modality Imaging Probes

Amit Kumar, Shanrong Zhang, Guiyang Hao, Gedaa Hassan, Saleh Ramezani, Su-tang Lo, Koji Sagiya, Masaya Takahashi, A. Dean Sherry, Orhan K Oz, Zoltán Kovács, and Xiankai Sun

*Bioconjugate Chem.*, **Just Accepted Manuscript** • DOI: 10.1021/acs.bioconjchem.5b00028 • Publication Date (Web): 23 Jan 2015

Downloaded from <http://pubs.acs.org> on February 18, 2015

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



**ACS Publications**  
High quality. High impact.

Bioconjugate Chemistry is published by the American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036  
Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

# A Molecular Platform for Design and Synthesis of Targeted Dual-modality Imaging Probes

Amit Kumar<sup>1</sup>, Shanrong Zhang<sup>2</sup>, Guiyang Hao<sup>1</sup>, Gedaa Hassan<sup>1</sup>, Saleh Ramezani<sup>1</sup>, Koji Sagiya-  
ma<sup>2</sup>, Su-Tang Lo<sup>1</sup>, Masaya Takahashi<sup>2</sup>, A. Dean Sherry<sup>2</sup>, Orhan, K. Öz<sup>1</sup>, Zoltan Kovacs<sup>2</sup>, Xiankai  
Sun<sup>1,2\*</sup>

<sup>1</sup>Department of Radiology, <sup>2</sup>Advanced Imaging Research Center, University of Texas Southwestern Medical Center, Dallas, Texas 75390, USA

**ABSTRACT:** We report a versatile dendritic structure based platform for construction of targeted dual-modality imaging probes. The platform contains multiple copies of 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA) branching out from a 1,4,7-triazacyclononane-N,N',N''-triacetic acid (NOTA) core. The specific coordination chemistries of the NOTA and DOTA moieties offer specific loading of <sup>68/67</sup>Ga<sup>3+</sup> and Gd<sup>3+</sup>, respectively, into a common molecular scaffold. The platform also contains three amino groups which can potentiate targeted dual-modality imaging of PET/MRI or SPECT/MRI (PET: Positron Emission Tomography; SPECT: Single Photon Emission Computed Tomography; MRI: Magnetic Resonance Imaging) when further functionalized by targeting vectors of interest. To validate this design concept, a bimetallic complex was synthesized with six peripheral Gd-DOTA units and one Ga-NOTA core at the center, whose ion T<sub>1</sub> relaxivity per gadolinium atom was measured to be 15.99 mM<sup>-1</sup>s<sup>-1</sup> at 20 MHz. Further, the bimetallic agent demonstrated its anticipated *in vivo* stability, tissue distribution, and pharmacokinetic profile when labeled with <sup>67</sup>Ga. When conjugated with a model targeting peptide sequence, the trivalent construct was able to visualize tumors in a mouse xenograft model by both PET and MRI via a single dose injection.

## INTRODUCTION


Molecular imaging is gaining importance for noninvasive assessment of biological events in animal and human subjects. Through *in vivo* real-time visualization and imaging quantification of biological targets of interest, a better understanding of the biological processes underlining the initiation and progression of diseases can be obtained for more efficacious diagnosis and treatment.<sup>1-2</sup> However, a single imaging technique is often unable to deliver all the necessary information required for a definitive clinical decision. To date, successful attempts have been seen to combine two or more imaging modalities in order to overcome the shortcomings of single modality systems for optimal visualization and better quantitative delineation of targets of interest. Consequently, multimodal imaging techniques have become a norm in both clinical practice and preclinical research as evidenced by the implementation of PET/CT (PET: Positron Emission Tomography; CT: Computed Tomography) and PET/MRI (MRI: Magnetic Resonance Imaging) procedures to patient care and the increasing use of a variety of combinations of the currently available imaging techniques in translational or basic biomedical research, such as MRI/optical and PET/NIRF (NIRF: Near-infrared Fluorescence).<sup>3-8</sup> The fusion of PET and MRI is especially desirable because technically their physical features are complementary. While radionuclide-based imaging techniques, either PET or SPECT (Single Photon Emission Computed Tomography), are extremely sensitive and quantitative, which allows the studies of biological events or processes at the molecular and cellular level, their spatial resolution is limited (> 1 cm for current clinical scanners).<sup>9-11</sup> On the other hand, while MRI is able to provide high spatial resolution (< 0.1 cm) with exquisite soft tissue contrast even without exogenous contrast agents, contrast agents are often required to highlight specific biological or physiological processes in an

amount a few orders of magnitude higher than PET or SPECT.<sup>12-14</sup>

Given that more and more PET/MRI systems will likely be used in the practice of diagnostic radiology, it becomes important to develop dual modality imaging probes that can realize the full potential of both modalities.

To take advantage of the strengths of PET and MRI, our approach is to design and synthesize a dual-modality agent scaffold offering a “single pharmacological behavior” for both imaging acquisitions. The merging of PET and MRI probe moieties to a single molecular platform would facilitate co-location and cross-validation of each modality in targeted regions of interest (two measures of one event). While MRI can provide the exact location of the probe, motion artifact correction, and PET partial volume correction, PET can afford better imaging quantification for higher detection sensitivity. Further, given that the proton MRI contrast actually reflects the map of the proton density, a co-localized PET signal distinct from the proton background could make the MRI contrast more identifiable, which further improves the MRI sensitivity.

Indeed several nanoparticle-based PET/MRI agents have been reported with enhanced magnetic relaxivities and considerable promise as dual-modality imaging agents.<sup>15-17</sup> However, a nanoparticle platform for PET/MRI probe development presents challenges. First, the nanoparticle-based PET/MRI probes are not single molecular entities and often have questionable *in vivo* stability or molecular integrity. Second, to serve the purpose of diagnostic imaging, contrast agents are desired to have a reasonable blood circulation half-life for efficient accumulation in targets and minimal nonspecific deposition in the mononuclear phagocyte system (MPS) organs after systemic administration. However, it is well known that most nanoparticles if not all possess suboptimal *in vivo* distribution as shown by rapid sequestration into MPS organs and thereafter slow



**Figure 1.** The dual-modality molecular probe design

clearance. These properties present potential health hazards<sup>18-20</sup> which could impede their translation toward clinical applications. Third, the majority of nanoparticle-based PET/MRI agents provide  $T_2$  contrast, only a handful are  $T_1$ -based agents, which are preferred in the cases of low proton density in the target tissues. To date, a wide range of macromolecules, such as perfluorocarbon emulsions,<sup>15</sup> silica nanoparticles,<sup>16, 21-22</sup> liposomal vesicles,<sup>17, 23-25</sup> dendrimers,<sup>26-32</sup> and polymers,<sup>33</sup> have been explored for the development of gadolinium-based  $T_1$  contrast agents. Of the macromolecular systems, dendrimers, which possess definitive molecular structures and formula, have shown a promising role. For instance, the dendrimer systems of poly(amido amine) (PAMAM) have been used as blood pool,<sup>34</sup> liver,<sup>35</sup> renal,<sup>36</sup> lymphatic,<sup>37</sup> and tumor-specific<sup>38</sup> contrast agents. However, the Gd-PAMAM complexes rarely afford an ionic relaxivity greater than 11 mM<sup>-1</sup>s<sup>-1</sup> per gadolinium ion.<sup>39-40</sup> When such dendrimer platforms are used to develop PET/MRI or SPECT/MRI agents, it is challenging to achieve precise control of radioisotope loading into specific chelating moieties. Herein, we present a dual-modality molecular probe design (Figure 1) carrying six 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA) moieties branched out from a single 1,4,7-triazacyclononane-1,4,7-triacetic acid (NOTA) core structure. In this work, we demonstrate an exclusive loading of Gd<sup>3+</sup> into the DOTA units affording a  $T_1$  ion relaxivity of 15.99 mM<sup>-1</sup>s<sup>-1</sup> at 20 MHz, while the NOTA core remains available to form a chelate with <sup>67/68</sup>Ga<sup>3+</sup> for nuclear imaging. If the targeted accumulation event is rapid, <sup>68</sup>Ga ( $t_{1/2}$  = 68 min;  $\beta^+$  = 89%,  $E_{\beta^+ \text{ max}}$  = 1.92 MeV; EC = 11%) will be used for PET imaging; otherwise, <sup>67</sup>Ga ( $t_{1/2}$  = 3.26 d;  $\gamma$  = 184 keV) will be used for SPECT imaging.


## RESULTS AND DISCUSSION

The dual-modality molecular probe design also possesses three amino groups for potential conjugation with multiple copies of targeting molecules, thereby providing multivalent effect to the construct for better specific binding to the biological target of interest. Of note, the strong interaction between the targeting molecule tethered probe and the corresponding target, triggered by multivalent effect, is expected to further enhance the relaxivity of the probe. For proof of concept we conjugated each amine with a model integrin  $\alpha_v\beta_3$  targeting peptide, c(RGDyK). The main features of this molecular dual-

modality imaging probe design include the apparent *in vivo* stability (due to the inertness of the Gd-DOTA and Ga-NOTA complexes) and the defined molecular formula with selective loading of two metal ions.

**Synthesis.** The desired ligand H<sub>21</sub>L was synthesized in 7 steps with an overall yield of approximately 15%. The structure of each intermediate was verified by <sup>1</sup>H NMR and <sup>13</sup>C NMR as well as mass spectroscopy. The synthesis route to compound H<sub>21</sub>L is outlined in Scheme 1 and its functionalization is depicted in Scheme 2. Compound **2** was obtained from bis(3-aminopropyl)amine and carbobenzyloxy (Cbz) protected lysine in three steps (see Supporting Information). The free primary amine of **2** was coupled to the carboxylic acid bearing side arms of NOTA-(OH)<sub>3</sub> (Scheme 1) via carbodiimide chemistry to form **3**. The N-phthaloyl protected amino groups of **3** were freed by deprotection via hydrazine to afford **4**, which contains six amino groups serving as points of attachment for the DOTA units.

**Scheme 1.** Synthetic route to ligand H<sub>21</sub>L




The N-hydroxy succinamide activated DOTA ester, **5**, was reacted to the six amino groups forming **6**. Attempts to deprotect the Boc groups of compound **6** led to partial deprotection of the Cbz groups. To overcome this problem, the ligand was fully deprotected by 30% HBr in an acetic acid solution. The synthesized ligand, H<sub>21</sub>L, was fully characterized. The ligand carries six DOTA units intended for chelating Gd<sup>3+</sup> and one NOTA unit intended for chelating Ga<sup>3+</sup>. Since Gd<sup>3+</sup> and Ga<sup>3+</sup> can be specifically loaded into the DOTA and NOTA moieties, respectively, based on their ion-size preference, the bimetallic molecular complex is expected to carry the two metal ions in a specified ratio so as to facilitate PET imaging based quantification.

Functionalization of ligand H<sub>21</sub>L was performed by utilizing the previously mentioned amino groups. Maleimido groups were tethered to amines using the commercially available reagent of N-succinimidyl 3-maleimidopropionate (Scheme 2). The model targeting peptide, c(RGDyK), was tethered to the maleimido carrying ligand H<sub>21</sub>L via thiol-maleimide coupling. Given that targeting molecules attached to the bimetallic complex may not be able to stretch out for specific binding because of the steric hindrance from the construct, additionally we intro-

duced a PEG<sub>12</sub> linker between the complex and targeting vector.

**Scheme 2.** Synthesis of integrin- $\alpha_v\beta_3$  targeted ligand scaffolds




The thiol group without a spacer (zero) or with a PEG<sub>12</sub> linker was introduced to c(RGDyK) using the commercially available reagent: 2-pyridyldithiol-tetraoxaoctatriacontane-N-hydroxysuccinimide (SPDP). The thiol terminated c(RGDyK) peptides with zero and PEG<sub>12</sub> linker were subsequently added to H<sub>21</sub>L(Mal)<sub>3</sub> to afford Gd<sub>6</sub>H<sub>3</sub>L[c(RGDyK)]<sub>3</sub> and Gd<sub>6</sub>H<sub>3</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub>, respectively.

**Complexation with Gd<sup>3+</sup> and Ga<sup>3+</sup> to form Gd<sub>6</sub>LGa.** The thermodynamic stability constants of Gd-DOTA (logK<sub>Gd-DOTA</sub> = 24.7), Gd-NOTA (logK<sub>Gd-NOTA</sub> = 14.3), and Ga-NOTA (logK<sub>Ga-NOTA</sub> = 31.0)<sup>41-42</sup> suggest that when ligand H<sub>21</sub>L is treated with six equivalents of Gd<sup>3+</sup>, its complex can be formed exclusively with the DOTA units. The bimetallic complex of H<sub>21</sub>L, Gd<sub>6</sub>LGa, can then be formed by incorporation of Ga<sup>3+</sup> into the NOTA core. However, considering that the reaction of L with the stoichiometric amount of Gd<sup>3+</sup> may be slow and could lead to partial complexation of the DOTA units, the first complexation was performed with excess of Gd<sup>3+</sup> to give complete chelation of both the DOTA and NOTA units forming Gd<sub>7</sub>L. This was confirmed with mass spectroscopy (see Supporting Information) as well as <sup>1</sup>H NMR spectra using Eu<sup>3+</sup> instead of Gd<sup>3+</sup>. Later, Gd<sup>3+</sup> was efficiently removed from the NOTA core by treating the complex with excess diethylene triamine pentaacetic acid (DTPA) (the log K of Gd-DTPA<sup>42</sup> is approximately eight orders of magnitude higher than that of Gd-NOTA). The incorporation of either cold Ga<sup>3+</sup>

or <sup>68</sup>Ga<sup>3+</sup> into the NOTA core of complex of Gd<sub>6</sub>H<sub>3</sub>L was then carried out at pH 3-5 in an HEPES buffer. The cold Gd<sub>6</sub>LGa complex was purified and characterized by HPLC and mass spectrometry. Various concentration of Gd<sub>6</sub>LGa were injected to inductively coupled plasma mass spectrometry (ICP-mass) and the ratio of Gd:Ga was calculated to be 6:1 (see Supporting Information). The Gd<sub>6</sub>L<sup>68</sup>Ga was purified and characterized by radio-HPLC using the cold complex as reference.

**Coordination chemistry of the NOTA core of H<sub>21</sub>L.** To prove the efficient removal of Gd<sup>3+</sup> from the NOTA core during the complexation procedure, the same complexation procedure was followed using Eu<sup>3+</sup> instead of Gd<sup>3+</sup>. Chemically, Eu<sup>3+</sup> and Gd<sup>3+</sup> have similar reactivities towards NOTA and DOTA. Since Eu<sup>3+</sup> is only weakly paramagnetic, the <sup>1</sup>H-NMR signals of its complexes are relatively sharp and also paramagnetically shifted. This facilitates relatively straightforward structural characterization by NMR spectroscopy. To accomplish this, the Eu<sup>3+</sup> complex of H<sub>21</sub>L, were synthesized via two procedures, *i.e.*, with and without a challenge by excess DTPA, followed by HPLC purification. Both complexes were characterized by <sup>1</sup>H NMR spectroscopy (400 MHz). Figures 2a and 2b show the <sup>1</sup>H NMR spectra of Eu<sup>3+</sup> complexes of H<sub>21</sub>L with and without the DTPA challenge, respectively. The DTPA treated Eu<sup>3+</sup> complex (Figure 2a) showed sharp peaks in the NOTA region (around 3 – 4 ppm) suggesting a metal free NOTA core as opposed to the untreated complex (Figure 2b), which showed broad peaks indicating the interaction of metal with the NOTA core. This characterization along with the mass spectroscopy results of Gd<sub>7</sub>L and Gd<sub>6</sub>H<sub>3</sub>L indicates the exclusive Gd<sup>3+</sup> coordination with the six DOTA moieties of H<sub>21</sub>L. Further, the formation of Gd<sub>6</sub>LGa was confirmed by the molar ratio of Gd:Ga determined by ICP-mass (supporting information). These results demonstrate that Gd<sup>3+</sup> and Ga<sup>3+</sup> have been exclusively loaded to the DOTA and NOTA moieties as designed, respectively, for the preparation of the desired dual-modality molecular imaging probe.




**Figure 2.** 400 MHz <sup>1</sup>H NMR spectra of a) Eu<sup>3+</sup> complex of H<sub>21</sub>L treated with excess DTPA b) untreated Eu<sup>3+</sup> complex of H<sub>21</sub>L, and c) Eu-DOTA complex.

**Relaxometric measurements.** The T<sub>1</sub> relaxivity of Gd<sub>6</sub>LGa was measured at 20 MHz, 25 °C, to be 95.98 mM<sup>-1</sup>s<sup>-1</sup> with R<sup>2</sup> value greater than 0.99 (Each Gd<sup>3+</sup> in the complex accounted for 15.99 mM<sup>-1</sup>s<sup>-1</sup> (ion relaxivity)). Under the same conditions, the relaxivity of Magnavist was found to be 4.1 mM<sup>-1</sup>s<sup>-1</sup>. Impressively, the T<sub>1</sub> relaxivity of Gd<sub>6</sub>H<sub>3</sub>L was approximately 4 times higher than that of Gd-DOTA complex and comparable



to the Gd-PAMAM complexes, which have been reported with the ion relaxivity up to  $11 \text{ mM}^{-1}\text{s}^{-1}$ .<sup>39–40</sup> It is noteworthy that a similar relaxivity was also reported for a polylysine-based contrast agent (MW 17,453) with 24 Gd-DOTA units.<sup>43–44</sup> At the physiological temperature of  $37^\circ\text{C}$ , the relaxivity of Gd<sub>6</sub>LGa was unchanged while its relaxivity in rat serum at  $25^\circ\text{C}$  increased to  $153.1 \text{ mM}^{-1}\text{s}^{-1}$ . These observations suggest two things; first, the lack of temperature dependence between  $25^\circ\text{C}$  and  $37^\circ\text{C}$  suggests that the water exchange rate in the appended Gd-DO3A-monoamide complexes limit the relaxivity somewhat at  $25^\circ\text{C}$ . Given that water exchange is faster at  $37^\circ\text{C}$ , the expected decline in relaxivity with increasing temperatures is offset by faster water exchange and a more favorable relaxivity at  $37^\circ\text{C}$ . This was an unanticipated yet welcomed added advantage for this probe platform design. The higher relaxivity in serum indicates that Gd<sub>6</sub>LGa does interact with one or more serum proteins.



**Figure 3.**  $T_1$ -weighted MR images of samples recorded at 1.0 T at  $25^\circ\text{C}$ . Samples of Magnevist (a  $\rightarrow$  e) and Gd<sub>6</sub>H<sub>3</sub>L (v  $\rightarrow$  z) containing a series of decreasing concentrations from 0.4 mM, 0.2 mM, 0.1 mM, 0.04 mM, to 0.02 mM.

**MR imaging and relaxivity measurements of Gd<sub>6</sub>LGa at 1.0 T.** The MR imaging potential of Gd<sub>6</sub>LGa was evaluated at magnetic field of 1.0 T. Figure 3 shows the  $T_1$  weighted imaging of Gd<sub>6</sub>LGa in comparison with Magnevist at a series of decreasing concentrations (0.4 mM – 0.02 mM determined by ICP-mass). The images of Gd<sub>6</sub>LGa were obviously brighter than those of Magnevist ( $4.1 \text{ mM}^{-1}\text{s}^{-1}$ ) at each equivalent concentration. The  $T_1$  relaxivity of Gd<sub>6</sub>LGa was measured at  $113.36 \text{ mM}^{-1}\text{s}^{-1}$  ( $T_1$  ion relaxivity:  $18.89 \text{ mM}^{-1}\text{s}^{-1}$ ) at 1.0 T ( $25^\circ\text{C}$ ). Since the agent is intended for conjugation to a targeting moiety for receptor-based imaging, such interactions could further enhance the relaxivity of the agent *in vivo*.

***In vivo* tissue distribution, pharmacokinetics, and stability of Gd<sub>6</sub>LGa.** As compared to other nanoparticle-based dual modality imaging platforms, our molecular probe design is expected to have a reasonable blood circulation half-life and an efficient clearance profile from the organs. To evaluate the *in vivo* behavior of this probe design, Gd<sub>6</sub>LH<sub>3</sub> was radio-labeled with <sup>67</sup>Ga ( $t_{1/2} = 3.26 \text{ d}$ ) so that the tissue biodistribution could be evaluated over an extended period. As shown in Table 1, the bimetallic complex, Gd<sub>6</sub>L<sup>67</sup>Ga, indeed displayed a rapid and efficient clearance profile from all major organs except kidneys in normal mice. The high renal uptake can be attributed to the net positive charge on the probe due to the presence of amine groups.<sup>45</sup> However, an efficient renal clearance is obvious after the initial high uptake as seen from the decrease of the kidney uptake level from  $101.0 \pm 4.3 \text{ \%ID/g}$  at 1 h p.i. to  $43.0 \pm 5.1 \text{ \%ID/g}$  at 24 h p.i. One would anticipate

that kidney clearance would be even faster once the positively charged amino groups are removed from the agent through conjugation with a targeting vector. Indeed when the amine groups of the agent were capped with c(RGDyK), the kidney uptake was impressively reduced to *ca.* 3 %ID/g at 2 h p.i.

The pharmacokinetics of Gd<sub>6</sub>L<sup>67</sup>Ga showed a two-compartment profile with  $t_{1/2\alpha} = 11.3 \text{ min}$  and  $t_{1/2\beta} = 7.3 \text{ h}$ , quite different compared to nanoparticle-based imaging agents which can be described by one compartment models.<sup>46–47</sup> The  $t_{1/2\alpha}$  reveals that the tissue distribution of Gd<sub>6</sub>L<sup>67</sup>Ga was relatively rapid while the reasonably long  $t_{1/2\beta}$  enables the probe to accumulate in its targeted tissues without incurring high background uptake. This indicates that the probe behaves similarly to other clinically available small molecule contrast agents.<sup>48</sup> The presence of amine groups on the ligand provides the ease of incorporation of polyethylene glycol (PEG) chains into the probe for optimization of the *in vivo* kinetics if necessary.

The *in vitro* and *in vivo* stabilities of Gd<sub>6</sub>L<sup>67</sup>Ga were evaluated through incubation in rat serum and by mouse urine metabolite analysis, respectively. The bimetallic complex remained intact during the entire course of study, which further validates this molecular probe platform design.


**Table 1.** Tissue distribution data and pharmacokinetic parameters of Gd<sub>6</sub>L<sup>67</sup>Ga in normal Balb/c mice (n = 4). Data are presented as mean  $\pm$  standard deviation.

Organ	%ID/g		
	1h	4h	24h
Blood	$3.51 \pm 0.22$	$2.63 \pm 0.84$	$0.21 \pm 0.06$
Heart	$1.21 \pm 0.13$	$0.94 \pm 0.40$	$0.21 \pm 0.02$
Lung	$2.59 \pm 0.5$	$1.78 \pm 0.87$	$0.38 \pm 0.09$
Liver	$1.39 \pm 0.11$	$2.13 \pm 0.81$	$1.39 \pm 0.24$
Kidney	$101.32 \pm 4.32$	$57.42 \pm 17.29$	$43.90 \pm 5.12$
Spleen	$0.73 \pm 0.19$	$0.69 \pm 0.07$	$0.42 \pm 0.12$
Muscle	$0.27 \pm 0.09$	$0.38 \pm 0.13$	$0.08 \pm 0.02$
Fat	$0.27 \pm 0.10$	$0.27 \pm 0.10$	$0.06 \pm 0.02$
Brain	$0.08 \pm 0.01$	$0.07 \pm 0.02$	$0.02 \pm 0.00$
Two-compartment clearance profile (min)			
$t_{1/2\alpha}$	$11.33 \pm 1.31$	$t_{1/2\beta}$	$438.37 \pm 15.11$

#### Receptor-binding assay of targeted Gd<sub>6</sub>LGa complexes.

Two integrin  $\alpha_v\beta_3$  targeted Gd<sub>6</sub>LGa complexes Gd<sub>6</sub>H<sub>3</sub>L[c(RGDyK)]<sub>3</sub> and Gd<sub>6</sub>H<sub>3</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub> were prepared to validate that our molecular construct design can be utilized for specific imaging of cancer biomarkers. The  $\alpha_v\beta_3$ -binding affinities were determined by a competitive cell-binding assay using <sup>125</sup>I-echistatin (PerkinElmer) as the  $\alpha_v\beta_3$ -specific radioligand. The best-fit IC<sub>50</sub> values (inhibitory concentration where 50% of the <sup>125</sup>I-echistatin bound on U87MG cells are displaced) of c(RGDyK), Gd<sub>6</sub>H<sub>3</sub>L[c(RGDyK)]<sub>3</sub> and Gd<sub>6</sub>H<sub>3</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub> were measured to be 199 nM, 489 nM, and 37 nM (Supporting Information), respectively. The significantly improved  $\alpha_v\beta$  binding affinity of Gd<sub>6</sub>H<sub>3</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub> clearly indicates the critical role

of the PEG<sub>12</sub> linker in the desired targeting property of our molecular scaffold design.



**Figure 4.** Representative MRI (upper panel), PET (lower panel – left), and fused PET/MR (lower panel – right) images of U87MG tumor xenograft in SCID mice at 1 h post injection of Gd<sub>6</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub><sup>68</sup>Ga. The white arrow indicates the tumor site.

**PET/MR Imaging using Gd<sub>6</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub><sup>68</sup>Ga.** To overcome the sensitivity difference between PET and MR, the radioactivity of <sup>68</sup>Ga was controlled in the labeling of Gd<sub>6</sub>H<sub>3</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub>. The final dose injected into each animal carried about 6.0 mg of the final compound with an activity of 100–150 μCi (Gd: *ca.* 0.027 mmol/kg; c(RGDyK): 2.0 μmol). Figure 4 shows *in vivo* dual PET/MR images of integrin α<sub>v</sub>β<sub>3</sub> positive U87MG tumor enabled by a single dose injection of Gd<sub>6</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub><sup>68</sup>Ga. The average tumor uptake measured by PET was 0.88 %ID/g. It should be noted that the tumor uptake value was reduced by roughly 3 times when the injected amount of the agent increased from 0.1 mg to 6.0 mg. This decrease also indicates the integrin α<sub>v</sub>β<sub>3</sub> imaging specificity. The observed MR contrast enhancement in the tumor was 3.06 (n = 3) when compared to the contrast before injection.

To date, few PET/MRI or SPECT/MRI agents with a defined structure have been reported. *In vivo* pH monitoring PET/MRI agents carrying one Gd-DOTA unit along with <sup>18</sup>F or <sup>68</sup>Ga as PET contrast have been reported.<sup>49</sup> Bimetallic complexes carrying one Gd-DOTA conjugated to <sup>68</sup>Ga-NOTA and one Gd-DOTA conjugated to cold Cu or In-DOTA do exist.<sup>50–51</sup> However, they were not designed to serve the goal of providing targeted contrast to both imaging techniques. To provide proof of concept, the probe design in this work consists of only six gadolinium chelatable DOTA units. While the anticipated T<sub>1</sub> relaxivity enhancement was achieved, we recognize that further amplification of MRI sensitivity is necessary in order to realize the practical application of the molecular probe design as dual-modality imaging agents. Given the versatility of the

design concept, we believe that it can be readily modified or adapted to reach the desired MRI sensitivity.

In conclusion, we have successfully demonstrated a molecular platform intended for targeted dual-modality imaging of PET/MRI or SPECT/MRI, which carries quantifiable numbers of <sup>67/68</sup>Ga<sup>3+</sup> and Gd<sup>3+</sup> ions in their specific chelation moieties. This probe design displays a very favorable high T<sub>1</sub> relaxivity for MRI contrast enhancement. With the current rapid growth of hybrid PET/MRI systems in diagnostic radiology, our molecular design of dual-modality agents possessing a “single pharmacological behavior” offers a versatile platform for the development of multimodality imaging agents with potential applications for noninvasive molecular profiling of various diseases.

## EXPERIMENTAL PROCEDURES

**General Materials and Methods** All reactions were carried out under N<sub>2</sub> atmosphere in degassed dried solvents. Commercially available starting materials were purchased from commercial vendors and used directly without further purification unless otherwise stated. All aqueous solutions were prepared with Milli-Q water. Bulk solvents were removed using rotary evaporator under reduced pressure at 40 °C. Trace solvents were removed under high vacuum. Matrix-assisted laser desorption/ionization (MALDI) mass spectra were acquired on an Applied Biosystems Voyager-6115 mass spectrometer. Radiolabeled conjugates were purified by Light C-18 Sep-Pak cartridges (Waters, Milford, MA). High performance liquid chromatography (HPLC) was performed on a Waters Xterra Shield RP C-18 semiprep column (250 × 10 mm, 10 μm) and read by a Waters 2996 photodiode array detector and an inline Shell Jr. 2000 radio-detector. The mobile phase was H<sub>2</sub>O with 0.1% trifluoroacetic acid (TFA) (solvent A) and acetonitrile with 0.1% TFA (solvent B). The gradient consisted of 0% B to 80% B in 0–40 min at 4.0 mL/min flow rate. The radioactivity of excised tissue samples and radioactive standards were counted by a Wizard2 300 automatic γ-counter (PerkinElmer).

## Synthesis Procedures

**Compound 3.** To a solution of the protected acid **1** (0.10 g, 0.15 mmol) in CH<sub>3</sub>CN (1.0 mL) were added the deprotected amine **2** (0.40 g, 0.60 mmol), dicyclohexylcarbodiimide (0.15 g, 0.83 mmol) and triethylamine (0.30 g, 0.27 mmol). The resultant solution was stirred for 12 h, filtered and the solvent evaporated. The crude product was purified by flash chromatography (ethyl acetate) to give **3**, a NOTA derivative, as a white solid (0.26 g, 0.10 mmol, 67%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.73 (bs, 6H), 7.71 (m, 12H), 7.61 (m, 12H), 7.23 (m, 15H), 5.56 (bs, 2H), 4.99 (m, 5H), 4.74 (m, 3H), 3.79–3.27 (m, 24H), 3.25–2.77 (m, 13H), 2.60–2.17 (m, 4H), 2.15–1.71 (m, 12H), 1.60 (m, 4H), 1.38 (m, 43H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 172.5, 168.3, 156.8, 136.5, 134.1, 133.9, 131.8, 128.4, 127.9, 123.3, 123.2, 79.0, 69.8, 66.5, 64.8, 49.3, 45.9, 44.3, 40.5, 35.5, 35.3, 32.5, 32.2, 29.3, 28.3, 27.9, 26.8, 22.6. MS (MALDI) *m/z* calcd for C<sub>141</sub>H<sub>168</sub>N<sub>18</sub>O<sub>30</sub>: 2594.2; found: 2595.9 ([M + H]<sup>+</sup>).

**Compound 4.** To a solution of **3** (0.26 g, 0.11 mmol) in ethanol (1.0 mL) was added hydrazine monohydrate (0.1 mL, 2.0 mM) and the mixture was stirred for 12 h at room temperature. After the reaction, the precipitate was removed by filtration. The filtrate was evaporated and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 10 mL). The combined organic layers were evaporated to give **4** as light yellow oil (0.15 g, 0.08 mmol, 80%). This compound was used for the next step without further purification. <sup>1</sup>H

NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  7.31 (s, 15H), 4.99 (m, 6H), 4.66 (m, 3H), 3.68-3.39 (m, 10H), 3.38-3.31 (m, 6H), 3.20-2.94 (m, 17H), 2.93-2.72 (m, 9H), 2.65-2.30 (m, 7H), 1.95-1.81 (m, 6H), 1.78-1.57 (m, 6H), 1.55-1.20 (m, 44H). <sup>13</sup>C NMR (100 MHz, CD<sub>3</sub>OD):  $\delta$  173.7, 173.1, 157.6, 136.9, 128.9, 127.3, 126.8, 126.5, 82.6, 77.3, 65.9, 64.5, 50.2, 48.8, 45.9, 44.5, 42.3, 38.5, 38.1, 36.7, 30.3, 29.1, 27.8, 26.6, 25.2 MS (MALDI)  $m/z$  calcd for C<sub>93</sub>H<sub>156</sub>N<sub>18</sub>O<sub>18</sub>: 1814.1; found: 1816.3 ([M + H]<sup>+</sup>).

**Compound 6.** To a solution of **4** (0.15 g, 0.08 mmol) in DMF (1 mL) were added **5** (0.53 g, 0.80 mM), an N-hydroxysuccinimide ester of DOTA, and the mixture was stirred for 24 h at room temperature. The solvent was evaporated and the product purified by reverse phase HPLC using water and acetonitrile solvent mixture to give **6** as a white solid (0.29 g, 0.06 mmol, 70%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>):  $\delta$  7.40-7.18 (s, 15H), 5.15-4.99 (m, 6H), 4.71 (m, 6H), 4.38-4.83 (m, 26H), 3.82-3.36 (m, 61H), 3.30-2.62 (m, 17H), 2.60-2.30 (m, 12H), 2.23-1.98 (m, 10H), 1.99-1.80 (m, 10H), 1.79-0.96 (m, 174H). <sup>13</sup>C NMR (100 MHz, CD<sub>3</sub>OD):  $\delta$  172.8, 172.6, 170.8, 170.6, 157.3, 137.2, 128.2, 127.6, 127.2, 84.5, 81.5, 65.8, 64.7, 53.6, 51.6, 49.2, 48.5, 45.3, 43.6, 42.0, 40.1, 36.5, 31.9, 27.2, 28.5, 27.2, 27.1, 22.7. MS (MALDI)  $m/z$  calcd for C<sub>261</sub>H<sub>456</sub>N<sub>42</sub>O<sub>60</sub>: 5139.4; found: 5138.7 ([M + H]<sup>+</sup>).

**Ligand H<sub>21</sub>L.** To a solution of **6** (0.10 g, 0.02 mmol) was added 30% HBr in AcOH (2 mL) and the solution was allowed to stir for 4 h. The solvent was evaporated, the product neutralized and purified by reverse phase HPLC using water and acetonitrile solvent mixture to give H<sub>21</sub>L as a white solid (0.06 g, 0.02 mmol, 75%). <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD):  $\delta$  4.42-3.62 (m, 89H), 3.59-3.32 (m, 67H), 3.24-2.85 (m, 48H), 2.80-2.30 (m, 22H), 2.34-2.03 (m, 17H), 2.06-1.85 (m, 14H), 1.88-1.65 (m, 20H), 1.65-1.39 (m, 11H), 1.31 (m, 6H)  $\delta$  MS (MALDI)  $m/z$  calcd for C<sub>153</sub>H<sub>270</sub>N<sub>42</sub>O<sub>54</sub>: 3561.9; found: 3563.0 ([M + H]<sup>+</sup>).

**Compound H<sub>21</sub>L(Mal)<sub>3</sub>.** To the free ligand H<sub>21</sub>L (0.200 g, 0.056 mmol) dissolved in DMF (1 mL) was added triethyl amine (0.022 g, 0.224 mmol) and N-( $\gamma$ -maleimidobutyryloxy)succinimide and the solution was allowed to stir for 24 h. The solvent was evaporated; the product neutralized and purified by reverse phase HPLC using water and acetonitrile solvent mixture to give H<sub>21</sub>L(Mal)<sub>3</sub> as a white solid (0.102 g, 0.052 mmol, 45.1%). MS (MALDI)  $m/z$  calcd for C<sub>174</sub>H<sub>285</sub>N<sub>45</sub>O<sub>63</sub>: 4015.0; found: 4016.3 ([M + H]<sup>+</sup>).

**Compound c(RGDyK)-SH.** To the DMF solution (1 mL) of the commercially available  $\alpha_v\beta_3$  integrin targeting peptide c(RGDyK) (0.030 g, 0.048 mmol) (Peptides International Inc, Kentucky) was added N-succinimidyl 3-(2-pyridyldithio)propionate (0.020 g, 0.064 mmol) (Thermo Scientific, IL) and the solution was allowed to stir for 6 h. The solvent was evaporated; the product neutralized and purified by reverse phase HPLC using water and acetonitrile solvent mixture and lyophilized. The resultant white solid was dissolved in DMF (1.0 mL) and dithiothreitol (0.010 g, 0.065 mmol) was then added. The solution was allowed to stir for 3 h. The solvent was evaporated and the product neutralized and purified by reverse phase HPLC using water and acetonitrile solvent mixture to give c(RGDyK)-SH as a white solid (0.103 g, 0.014 mmol, 29.5%). MS (MALDI)  $m/z$  calcd for C<sub>30</sub>H<sub>45</sub>N<sub>9</sub>O<sub>9</sub>S: 707.3; found: 708.3 ([M + H]<sup>+</sup>).

**Compound H<sub>21</sub>L[c(RGDyK)]<sub>3</sub>.** To the maleimide carrying ligand, H<sub>21</sub>L(Mal)<sub>3</sub>, (0.020 g, 0.005 mmol) dissolved in PBS

(1 x) was added the thiol carrying c(RGDyK)-SH (0.014 g, 0.029 mmol) and the solution was allowed to stir for 18 h. The solution was purified by reverse phase HPLC using water and acetonitrile solvent mixture to give H<sub>21</sub>L[c(RGDyK)]<sub>3</sub> as a white solid (0.010 g, 0.002 mmol, 33.3%). MS (MALDI)  $m/z$  calcd for C<sub>264</sub>H<sub>420</sub>N<sub>72</sub>O<sub>90</sub>S<sub>3</sub>: 6136.9; found: 6139.1 ([M + H]<sup>+</sup>).

**Compound c(RGDyK)PEG<sub>12</sub>SH.** To the DMF solution (1 mL) of the  $\alpha_v\beta_3$  integrin targeting peptide c(RGDyK) (0.040 g, 0.064 mmol) (Peptides International Inc, Kentucky) dissolved in DMF (1.0 mL) was added N-2-pyridyldithiol-tetraoxacontane-N-hydroxysuccinimide (0.060 g, 0.065 mmol) (Thermo Scientific, IL) and the solution was allowed to stir for 6 h. The solvent was evaporated and the product neutralized and purified by reverse phase HPLC using water and acetonitrile solvent mixture and lyophilized. The resultant white solid was dissolved in DMF (1 mL) and dithiothreitol (0.010 g, 0.065 mmol) was then added and the solution was allowed to stir for 3 h. The solvent was evaporated and the product neutralized and purified by reverse phase HPLC using water and acetonitrile solvent mixture to give c(RGDyK)PEG<sub>12</sub>SH as a colorless viscous liquid (0.025 g, 0.019 mmol, 30.2%). MS (MALDI)  $m/z$  calcd for C<sub>57</sub>H<sub>98</sub>N<sub>10</sub>O<sub>22</sub>S: 1306.6; found: 1307.5 ([M + H]<sup>+</sup>).

**Compound H<sub>21</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub>.** To the maleimide carrying ligand, H<sub>21</sub>L(Mal)<sub>3</sub> (0.030 g, 0.007 mmol), dissolved in PBS (1 x) was added the thiol carrying peptide c(RGDyK)-SH (0.038 g, 0.028 mmol) and the solution was allowed to stir for 18 h. The solution was purified by reverse phase HPLC using water and acetonitrile solvent mixture to give H<sub>21</sub>L[PEG<sub>12</sub>c(RGDyK)]<sub>3</sub> as a white solid (0.019 g, 0.002 mmol, 35.3%). MS (MALDI)  $m/z$  calcd for C<sub>345</sub>H<sub>579</sub>N<sub>75</sub>O<sub>129</sub>S<sub>3</sub>: 7936.0; found: 7936.7 ([M + H]<sup>+</sup>).

**Compound Gd<sub>7</sub>L.** The free ligand H<sub>21</sub>L (0.060 g, 0.013 mmol) was dissolved in water (1 mL) and the pH was adjusted to 7.0 with NaOH (0.1 M). To this solution was added an excess amount of GdCl<sub>3</sub>·6H<sub>2</sub>O and the pH was readjusted to 6.5 and allowed to stir at room temperature overnight. The pH of the resultant solution was raised above 8 using 1 M aqueous NaOH, causing the excess Gd<sup>3+</sup> to precipitate as Gd(OH)<sub>3</sub>. The solution was filtered and the pH was readjusted to 7.0 using 1 N HCl. the solution was purified using HPLC and the fractions pooled together and lyophilized to give a white solid (0.051 g, 0.011 mmol, 84.6%). MS (MALDI)  $m/z$  calcd for C<sub>153</sub>H<sub>261</sub>Gd<sub>7</sub>N<sub>42</sub>O<sub>60</sub>: 4644.2; found: 4645.1 ([M + H]<sup>+</sup>).

**Compound Gd<sub>6</sub>H<sub>3</sub>L.** To the solution of Gd<sub>7</sub>L (0.05 g, 0.011 mmol), DTPA (0.1 mM, 1mL) was added. The solution was stirred at room temperature for 2 h and then purified using HPLC. The desired fractions were pooled together and lyophilized to give Gd<sub>6</sub>H<sub>3</sub>L as a white solid (0.04 g, 0.01 mmol, 81%). MS (MALDI)  $m/z$  calcd for C<sub>153</sub>H<sub>264</sub>Gd<sub>6</sub>N<sub>42</sub>O<sub>60</sub>: 4489.3; found: 4634.9 ([M + K + 6H<sub>2</sub>O]<sup>+</sup>).

**Compound Eu<sub>6</sub>H<sub>3</sub>L.** The free ligand H<sub>21</sub>L (0.030 g, 0.008 mmol) was dissolved in water (1 mL) and the pH was adjusted to 7.0 with NaOH (0.1 M). To this solution was added an excess amount of EuCl<sub>3</sub>·6H<sub>2</sub>O and the pH was readjusted to 6.5 and allowed to stir at room temperature overnight. The pH was raised above 8.0 using 1.0 N aqueous NaOH, which caused the excess Eu<sup>3+</sup> to precipitate as Eu(OH)<sub>3</sub>. The solution was filtered and the pH was readjusted to 7.0 using 1.0 N HCl. To the resulting solution, DTPA (0.1 mM, 1mL), was added and the solution was purified using HPLC to give the desired complex. The desired fractions were pooled together and lyophilized to

give  $\text{Eu}_6\text{LH}_3$  as a white solid. (0.042 g, 0.009 mmol, 54%). MS (MALDI)  $m/z$  calcd for  $\text{C}_{153}\text{H}_{252}\text{Eu}_6\text{N}_{42}\text{O}_{54}$  : 4453.4; found: 4454.5 ( $[\text{M} + \text{H}]^+$ ).

**Compound Cold  $\text{Gd}_6\text{LGa}$ .** The Gadolinium complex,  $\text{Gd}_6\text{H}_3\text{L}$  (0.001 g, 0.225  $\mu\text{mol}$ ) was dissolved in a solution of 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES, pH = 6.5, 1.0 M, 1.0 mL). To the resulting solution was added a solution of  $\text{GaCl}_3$  (0.0001 g, 0.567  $\mu\text{mol}$ ) in 0.6 N HCl (0.3 mL) and the resulting solution was stirred for 1 h. To the mixture was added 500  $\mu\text{L}$  of 5.0 mM ethylenediaminetetraacetic acid (EDTA) and the mixture was stirred for another 5 min at room temperature (EDTA was used to remove nonspecifically bound or free  $\text{GaCl}_3$  from the complex). The purification of the bimetallic complex was carried out by passing the mixture through a preconditioned Sep-Pak C-18 heavy cartridge. After thorough rinsing ( $3 \times 5$  mL, water) of the cartridge, the bimetallic complex,  $\text{Gd}_6\text{LGa}$ , was eluted by an ethanol-water mixture (70:30). The product was purified with HPLC and characterized by mass spec. MS (MALDI)  $m/z$  calcd for  $\text{C}_{153}\text{H}_{264}\text{GaGd}_6\text{N}_{42}\text{O}_{60}$  : 4555.28; found: 4556.9 ( $[\text{M} + \text{H}]^+$ ).

**Compound  $\text{Gd}_6\text{L}^{68}\text{Ga}$ .** In a 1.5 mL eppendorf tube containing 30  $\mu\text{g}$   $\text{Gd}_6\text{H}_3\text{L}$  complex in 1.0 mL of HEPES (pH = 6.5) solution, was added a solution of 8.0 mCi of  $^{68}\text{GaCl}_3$  in 0.6 N HCl. The reaction mixture was incubated at 75  $^\circ\text{C}$  for 0.5 h on a shaker. To this solution was added DTPA (5.0 mM, 5.0  $\mu\text{L}$ ) and the reaction mixture was stirred for 5 min at room temperature (DTPA was used to remove nonspecifically bound or free  $^{68}\text{Ga}$  from  $\text{Gd}_6\text{L}^{68}\text{Ga}$ ). The  $\text{Gd}_6\text{L}^{68}\text{Ga}$  complex was purified by passing the mixture through a preconditioned Sep-Pak C-18 light cartridge. After thorough rinsing ( $3 \times 3$  mL, water) of the cartridge,  $\text{Gd}_6\text{L}^{68}\text{Ga}$  was eluted by an ethanol-water mixture (70:30) to give 3.6 mCi of labeled compound. The product was first analyzed by a Rita Star Radioisotope TLC Analyzer (Straubenhardt, Germany) on instant thin-layer chromatography (ITLC-SG) plates (Pall Life Sciences, East Hills, NY) and then by radio-HPLC to determine the radiochemical purity of the product. The compound was determined to have more than 95% purity.

**Compound  $\text{Gd}_6\text{L}^{67}\text{Ga}$ .** In a 1.5 mL eppendorf tube containing 30  $\mu\text{g}$   $\text{Gd}_6\text{H}_3\text{L}$  complex in 1.0 mL of HEPES (pH = 6.5) solution, was added a solution of 4.0 mCi of  $^{67}\text{GaCl}_3$  in 0.6 N HCl. The reaction mixture was shaken and incubated at 75  $^\circ\text{C}$  for 0.5 h. To this solution, was added DTPA (5.0 mM, 5.0  $\mu\text{L}$ ) and the reaction mixture was incubated for 5 min at room temperature. The  $\text{Gd}_6\text{L}^{67}\text{Ga}$  complex was purified by passing the mixture through a preconditioned Sep-Pak C-18 light cartridge. After thorough rinsing ( $3 \times 3$  mL, water) of the cartridge,  $\text{Gd}_6\text{L}^{67}\text{Ga}$  was eluted by an ethanol-water mixture (70:30) to give 2.1 mCi of the labeled compound. The product was first analyzed by a TLC Analyzer on ITLC-SG plates and then by radio-HPLC to determine the radiochemical purity of the product. The compound was determined to have more than 95% purity.

**Compound  $\text{Gd}_6\text{H}_3\text{L}[\text{c}(\text{RGDyK})]_3$**  The c(RGDyK) modified ligand  $\text{H}_{21}\text{L}[\text{c}(\text{RGDyK})]_3$  (0.010 g, 0.002 mmol) was dissolved in water (1 mL) and the pH was adjusted to 7 with NaOH (0.1 M). To this solution was added an excess of  $\text{GdCl}_3 \cdot 6\text{H}_2\text{O}$  and the pH was again adjusted to 6.5 and allowed to stir at room temperature overnight. The pH was raised above 8.0 using 1.0 N aqueous NaOH, which caused the excess  $\text{Gd}^{3+}$  to precipitate as  $\text{Gd}(\text{OH})_3$ . The solution was filtered and the pH was readjusted to 7.0 using 1.0 N HCl. To the re-

sulting solution, DTPA (0.1 mM, 1 mL), was added and the solution was purified using HPLC to give the desired complex. The desired fractions were pooled together and lyophilized to give  $\text{Gd}_6\text{H}_3\text{L}[\text{c}(\text{RGDyK})]_3$  as a white solid. (0.005 g, 0.001 mmol, 48%). MS (MALDI)  $m/z$  calcd for  $\text{C}_{264}\text{H}_{402}\text{Gd}_6\text{N}_{72}\text{O}_{90}\text{S}_3$  : 7064.4; found: 7085.0 ( $[\text{M} + \text{Na}]^+$ ).

**Compound  $\text{Gd}_6\text{L}[\text{c}(\text{RGDyK})]_3^{68}\text{Ga}$ .** In a 1.5 mL eppendorf tube containing 80  $\mu\text{g}$   $\text{Gd}_6\text{H}_3\text{L}[\text{c}(\text{RGDyK})]_3$  complex in 1 mL of HEPES (pH = 6.5) solution, was added a solution of 15.0 mCi of  $^{68}\text{GaCl}_3$  in 0.6 N HCl. The reaction mixture was incubated at 75  $^\circ\text{C}$  for 0.5 h on a shaker. To this solution was added DTPA (5.0 mM, 5.0  $\mu\text{L}$ ) and the reaction mixture was incubated for 5 min at room temperature. The  $^{68}\text{Ga}$ -labeled conjugate,  $\text{Gd}_6\text{L}[\text{c}(\text{RGDyK})]_3^{68}\text{Ga}$ , was purified by passing the mixture through a preconditioned Sep-Pak C-18 light cartridge. After thorough rinsing ( $3 \times 5$  mL, water) of the cartridge,  $\text{Gd}_6\text{L}[\text{c}(\text{RGDyK})]_3^{68}\text{Ga}$  was eluted by an ethanol-water mixture (70:30) to give 9.0 mCi of the labeled compound. The product was analyzed by radio-HPLC to determine the radiochemical purity of the product. The radiochemical purity of the compound was determined to be higher than 95%.

**Compound  $\text{Gd}_6\text{H}_3\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3$**  The c(RGDyK) modified ligand  $\text{H}_{21}\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3$  (0.019 g, 0.002 mmol) was dissolved in water (1.0 mL) and the pH was adjusted to 7.0 with NaOH (0.1 M). To this solution was added an excess of  $\text{GdCl}_3 \cdot 6\text{H}_2\text{O}$  and the pH was again adjusted to 6.5 and allowed to stir at room temperature overnight. The pH was raised above 8.0 using 1.0 N aqueous NaOH, which caused the excess  $\text{Gd}^{3+}$  to precipitate as  $\text{Gd}(\text{OH})_3$ . The solution was filtered and the pH was readjusted to 7.0 using 1.0 N HCl. To the resulting solution, DTPA (0.1 mM, 1.0 mL), was added and the solution was purified using HPLC to give the desired complex. The desired fractions were pooled together and lyophilized to give  $\text{Gd}_6\text{H}_3\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3$  as a white solid. (0.006 g, 0.001 mmol, 51%). MS (MALDI)  $m/z$  calcd for  $\text{C}_{345}\text{H}_{561}\text{Gd}_6\text{N}_{75}\text{O}_{129}\text{S}_3$  : 8864.3; found: 8864.7 ( $[\text{M} + \text{H}]^+$ ).

**Compound  $\text{Gd}_6\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3\text{Ga}$ .** The gadolinium complex,  $\text{Gd}_6\text{H}_3\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3$  (0.001 g, 0.112  $\mu\text{mol}$ ) was dissolved in an HEPES solution (pH = 6.5, 1.0 M, 1.0 mL). To the resulting solution was added a solution of  $\text{GaCl}_3$  (0.001 g, 0.567  $\mu\text{mol}$ ) in 0.6 N HCl (0.3 mL) and the resulting solution was stirred for 1 h. To the mixture was added 500  $\mu\text{L}$  of 5.0 mM of EDTA and the mixture, was allowed to incubate for another 5 min at room temperature (EDTA was used to remove nonspecifically bound or free  $\text{GaCl}_3$ ). The purification of  $\text{Gd}_6\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3\text{Ga}$  was carried out by passing the mixture through a preconditioned Sep-Pak C-18 heavy cartridge. After thorough rinsing ( $3 \times 5$  mL, water) of the cartridge,  $\text{Gd}_6\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3\text{Ga}$  was eluted by an ethanol-water mixture (70:30). The product was characterized by mass spec. MS (MALDI)  $m/z$  calcd for  $\text{C}_{345}\text{H}_{558}\text{GaGd}_6\text{N}_{75}\text{O}_{129}\text{S}_3$  : 8930.9; found: 8954.8 ( $[\text{M} + \text{Na}]^+$ ).

**Compound  $\text{Gd}_6\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3^{68}\text{Ga}$ .** For PET/MR imaging, a slightly different labeling procedure was followed. To a 1.5 mL eppendorf tube containing 6.0 mg of  $\text{Gd}_6\text{H}_3\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3$  complex in 1.0 mL of HEPES (pH = 6.5) solution was added a solution of 300  $\mu\text{Ci}$  of  $^{68}\text{GaCl}_3$  in 0.6 N HCl. The reaction mixture was incubated at 75  $^\circ\text{C}$  for 10 min on a shaker. To this solution was added DTPA (5.0 mM, 5.0  $\mu\text{L}$ ) and the reaction mixture was incubated for 1 min at room temperature. The  $^{68}\text{Ga}$ -labeled conjugate was purified by passing the mixture through a preconditioned



tioned Sep-Pak C-18 plus cartridge. After thorough rinsing (3 × 5 mL, water) of the cartridge,  $\text{Gd}_6\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3^{67}\text{Ga}$  was eluted by an ethanol-water mixture (70:30) to give 180–200  $\mu\text{Ci}$  of the labeled compound. The product was analyzed by radio-HPLC to determine the radiochemical purity of the product. The radiochemical purity of the compound was determined to be higher than 95%.

**Biodistribution and pharmacokinetics studies of  $\text{Gd}_6\text{L}^{67}\text{Ga}$ .** Male BALB/C mice were injected with 300  $\mu\text{Ci}$  of  $\text{Gd}_6\text{L}^{67}\text{Ga}$  complex to evaluate the tissue distribution of the tracer in mice. Mice were sacrificed 1 h, 4 h, and 24 h post injection (p.i.). The organs of interest (blood, heart, lung, liver, spleen, kidney, stomach, muscle, fat, small intestine, large intestine, and brain) were harvested, weighed, and radioactivity was quantified using a  $\gamma$ -counter. Standards were prepared and counted along with the tissue samples to calculate the percentage-injected dose per gram (%ID/g). To determine the pharmacokinetic parameters, mice injected with the tracer were blood sampled from the retro-orbital sinus at 2 min, 5 min, 10 min, 30 min, 1 h, 2 h, 24 h, and 48 h p.i. and quantified using a  $\gamma$ -counter. The pharmacokinetic parameters were calculated based on a two-compartment open model.

***In vitro* and *ex vivo* stability of  $\text{Gd}_6\text{L}^{67}\text{Ga}$ .** The *in vitro* stability test was performed in rat serum. Briefly,  $\text{Gd}_6\text{L}^{67}\text{Ga}$  complex (100  $\mu\text{Ci}$ , 5  $\mu\text{L}$ ) was added into 100  $\mu\text{L}$  of rat serum ( $n = 3$ ). The solution was incubated for 1 h, 4 h 24 h and 48 h at 37 °C, respectively. The solution was vortexed and centrifuged for 5 min at the speed of 21,000 g. The supernatant was then analyzed by HPLC. For the *in vivo* stability evaluation, each male mouse was injected with 600  $\mu\text{Ci}$  of  $\text{Gd}_6\text{L}^{67}\text{Ga}$  complex in 100  $\mu\text{L}$  of saline via the tail vein. The urine samples were collected at 1h, 4 h 24 h and 48 h p.i. and then analyzed by HPLC.

**Integrin  $\alpha_v\beta_3$  receptor-binding assay.** The  $\alpha_v\beta_3$  integrin-binding affinities of  $\text{c}(\text{RGDyK})$ ,  $\text{Gd}_6\text{H}_3\text{L}[\text{c}(\text{RGDyK})]_3$  and  $\text{Gd}_6\text{H}_3\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3$  were determined by a competitive cell-binding assay using  $^{125}\text{I}$ -echistatin (PerkinElmer) as the  $\alpha_v\beta_3$ -specific radioligand. The experiments were performed on U87MG human glioblastoma cells following a previously reported method.<sup>52</sup> Briefly, U87MG cells were grown in RPMI 1640 medium supplemented with penicillin, streptomycin, and 10% (v/v) fetal bovine serum (FBS) at 37 °C under 5%  $\text{CO}_2$ . Suspended U87MG cells in the binding buffer (20 mM Tris, pH 7.4, 150 mM NaCl, 2.0 mM  $\text{CaCl}_2$ , 1.0 mM  $\text{MgCl}_2$ , 1.0 mM  $\text{MnCl}_2$ , 0.1% bovine serum albumin) were seeded on multiwell DV plates (Millipore) with  $5 \times 10^4$  cells per well and then incubated with  $^{125}\text{I}$ -echistatin (10,000 cpm/well) in the presence of increasing concentrations (0–5000 nM) of each  $\text{c}(\text{RGDyK})$  peptide conjugate for 2 h. The final volume in each well was maintained at 200  $\mu\text{L}$ . At the end of incubation, unbound  $^{125}\text{I}$ -echistatin was removed by filtration followed by three rinses with cold binding buffer. The retentate was collected and the radioactivity was measured using a  $\gamma$ -counter. The best-fit  $\text{IC}_{50}$  values (inhibitory concentration where 50% of the  $^{125}\text{I}$ -echistatin bound on U87MG cells are displaced) of  $\text{c}(\text{RGDyK})$ ,  $\text{Gd}_6\text{H}_3\text{L}[\text{c}(\text{RGDyK})]_3$  and  $\text{Gd}_6\text{H}_3\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3$  were calculated by fitting the data with nonlinear regression using GraphPad Prism (GraphPad-Software, Inc.). Experiments were repeated with quintuplicate samples.

**Relaxivity measurements at 0.5 Tesla (T).** The  $T_1$  values were recorded at 23 MHz (0.5 T) at 25 °C by using a Maran

Ultra relaxometer (Oxford Instruments, UK). Longitudinal relaxation times were measured by using the inversion-recovery pulse sequence ( $180^\circ\text{-t-}90^\circ$ ). The  $T_1$  relaxivities were determined by the linear regression analysis of the water proton relaxation rates in solutions ranging in concentration from 0.005 to 12.0 mM, in Millipore water in triplicates.

**MRI imaging and relaxivity measurements at 1.0 T.** The  $T_1$ -weighted MR images of samples in 0.5 mL microfuge tubes were collected using Aspect Imaging M2TM 1.0 T (43.5–45 MHz, 60 mm diameter volume coil) Gradient Echo Spoiled External Averaging (GRE-EXT) sequence. For imaging, the following parameters were used: TR = 14.2 ms; effective echo time (TE) = 2.8 ms; FOV  $64 \times 100 \text{ mm}^2$ , data matrix =  $256 \times 256$ , averaging = 4, slice = 1 mm. The  $T_1$ -maps of the samples at 1.0 T were determined from a series of multi- $T_1$  (0.01 ms to 5 s) Inversion Recovery Spin Echo sequence (IR-SE with minimum TE = 7.0 ms); TR = 10 s; FOV =  $64 \times 100 \text{ mm}^2$ ; matrix =  $128 \times 128$ ; averaging = 1, steady state scans = 10, fitted and calculated using Matlab code. All the fits for  $T_1$  values used to calculate the longitudinal relaxivity,  $r_1$ , had fitting coefficients,  $R^2 \geq 0.99$ . Three trials were performed. The Gd and Ga metal concentrations of samples were determined by inductively coupled plasma mass spectrometry (ICP-MS).

**PET/MR Imaging using  $\text{Gd}_6\text{L}[\text{PEG}_{12}\text{c}(\text{RGDyK})]_3^{68}\text{Ga}$ .** The MR and PET imaging studies were performed on a 1.0 T MR scanner (Aspect Imaging, Shoham, Israel) using a 35 mm mouse body coil, and a Siemens Inveon Multimodality PET/CT system (Siemens Medical Solutions Inc., Knoxville, TN, USA), respectively. Ten minutes prior to imaging, the animals were anesthetized using 3% isoflurane at room temperature until stable vitals were established. Once the animals were sedated, they were placed onto a custom made interchangeable bed between the MR and PET scanners under 2% isoflurane anesthesia for the duration of the imaging. A  $T_1$ -weighted gradient echo, spoiled sequence (echo time/repetition time [TE/TR] 3/15 msec; field of view [FOV]  $80 \times 100 \text{ mm}$ ; slice thickness 0.8 mm; matrix  $256 \times 256$ ; number of excitations [NEX] = 3; flip angle =  $25^\circ$ ) was performed for each mouse. Each mouse was injected with 100 – 150  $\mu\text{Ci}$  (Gd: ca. 0.027 mmol/kg) via the tail-vein. MR scans were obtained pre-contrast and 1 h after contrast medium injection.

PET imaging was acquired with the mouse mounted on the interchangeable bed directly following the MR acquisition. A 15 min PET static scan was performed and reconstructed using Fourier Rebinning and Ordered Subsets Expectation Maximization 3D (OSEM3D) algorithm. Reconstructed MR and PET images were fused and analyzed using the VivoQuant software (Invivo LLC, Boston, USA). For quantification, regions of interest were placed on the tumor and muscle. The latter was used for normalization. The resulting quantitative data were expressed in intensity and %ID/g for MR and PET images.

## ASSOCIATED CONTENT

**Supporting Information.** Supporting schemes, synthesis of compounds, ICP quantification are available free of charge via the Internet at <http://pubs.acs.org>.

## AUTHOR INFORMATION

### Corresponding Author

\*Correspondence to Xiankai Sun, Ph.D., Department of Radiology and Advanced Imaging Research Center, University of Texas Southwestern Medical Center at Dallas, 2201 Inwood Road, Texas 75390-8542, USA;  
E-mail: Xiankai.Sun@UTSouthwestern.edu.

### Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

### Funding Sources

This work was partially supported by the Prostate Cancer Research Program of the United States Army Medical Research and Materiel Command (W81XWH-12-1-0336) and the Dr. Jack Krohmer Professorship Funds.

### REFERENCES

- (1) Weissleder, R. (2006) Molecular Imaging in Cancer. *Science*, 312, 1168-1171.
- (2) Cheon, J.; Lee, J.-H. (2008) Synergistically Integrated Nanoparticles as Multimodal Probes for Nanobiotechnology. *Acc. Chem. Res.* 41, 1630-1640.
- (3) Jennings, L. E.; Long, N. J. (2009) 'Two is better than one'-probes for dual-modality molecular imaging. *Chem. Commun.* 28, 3511-3524.
- (4) Ntziachristos, V.; Yodh, A. G.; Schnall, M.; Chance, B. (2000) Concurrent MRI and diffuse optical tomography of breast after indocyanine green enhancement. *Proc. Natl. Acad. Sci. U.S.A.* 97, 2767-2772.
- (5) Beyer, T.; Townsend, D. W.; Brun, T.; Kinahan, P. E.; Charron, M.; Roddy, R.; Jerin, J.; Young, J.; Byars, L.; Nutt, R. (2000) A Combined PET/CT Scanner for Clinical Oncology. *J. Nucl. Med.* 41, 1369-1379.
- (6) Murray, C. B.; Norris, D. J.; Bawendi, M. G. (1993) Synthesis and characterization of nearly monodisperse CdE (E = sulfur, selenium, tellurium) semiconductor nanocrystallites. *J. Am. Chem. Soc.* 115, 8706-8715.
- (7) Link, S.; El-Sayed, M. A. (1999) Spectral Properties and Relaxation Dynamics of Surface Plasmon Electronic Oscillations in Gold and Silver Nanodots and Nanorods. *J. Phys. Chem. B.* 103, 8410-8426.
- (8) Alivisatos, A. P. (1996) Semiconductor Clusters, Nanocrystals, and Quantum Dots. *Science*, 271, 933-937.
- (9) Catana, C.; Wu, Y.; Judenhofer, M. S.; Qi, J.; Pichler, B. J.; Cherry, S. R. (2006) Simultaneous Acquisition of Multislice PET and MR Images: Initial Results with a MR-Compatible PET Scanner. *J. Nucl. Med.* 47, 1968-1976.
- (10) Cherry, S. R. (2006) The 2006 Henry N. Wagner Lecture: Of Mice and Men (and Positrons)--Advances in PET Imaging Technology. *J. Nucl. Med.* 47, 1735-1745.
- (11) Cherry, S. (2001) Fundamentals of positron emission tomography and applications in preclinical drug development. *J. Clin. Pharmacol.* 41, 482-491.
- (12) Caravan, P.; Ellison, J. J.; McMurry, T. J.; Lauffer, R. B. (1999) Gadolinium(III) Chelates as MRI Contrast Agents: Structure, Dynamics, and Applications. *Chem. Rev.* 99, 2293-2352.
- (13) Raymond, K. N.; Pierre, V. r. C. (2004) Next Generation, High Relaxivity Gadolinium MRI Agents. *Bioconjugate Chem.* 16, 3-8.
- (14) Seo, W. S.; Lee, J. H.; Sun, X.; Suzuki, Y.; Mann, D.; Liu, Z.; Terashima, M.; Yang, P. C.; McConnell, M. V.; Nishimura, D. G., et al. (2006) FeCo/graphitic-shell nanocrystals as advanced magnetic-resonance-imaging and near-infrared agents. *Nat. Mater.* 5, 971-976.
- (15) Morawski, A. M.; Winter, P. M.; Crowder, K. C.; Caruthers, S. D.; Fuhrhop, R. W.; Scott, M. J.; Robertson, J. D.; Abendschein, D. R.; Lanza, G. M.; Wickline, S. A. (2004) Targeted nanoparticles for quantitative imaging of sparse molecular epitopes with MRI. *Magn. Reson. Med.* 51, 480-486.
- (16) Santra, S.; Bagwe, R. P.; Dutta, D.; Stanley, J. T.; Walter, G. A.; Tan, W.; Moudgil, B. M.; Mericle, R. A. (2005) Synthesis and Characterization of Fluorescent, Radio-Opaque, and Paramagnetic Silica Nanoparticles for Multimodal Bioimaging Applications. *Adv. Mater.* 17, 2165-2169.
- (17) Cheng, Z.; Tsourkas, A. (2008) Paramagnetic Porous Polymersomes. *Langmuir*, 24, 8169-8173.
- (18) Cheng, L.; Yang, K.; Li, Y.; Chen, J.; Wang, C.; Shao, M.; Lee, S.-T.; Liu, Z. (2011) Facile Preparation of Multifunctional Upconversion Nanoprobes for Multimodal Imaging and Dual-Targeted Photothermal Therapy. *Angew. Chem. Int. Ed.* 123, 7523-7528.
- (19) Chen, J.; Wiley, B.; Li, Z. Y.; Campbell, D.; Saeki, F.; Cang, H.; Au, L.; Lee, J.; Li, X.; Xia, Y. (2005) Gold Nanocages: Engineering Their Structure for Biomedical Applications. *Adv. Mater.* 17, 2255-2261.
- (20) Gao, X.; Cui, Y.; Levenson, R. M.; Chung, L. W. K.; Nie, S. (2004) In vivo cancer targeting and imaging with semiconductor quantum dots. *Nat. Biotech.* 22, 969-976.
- (21) Lin, Y.-S.; Hung, Y.; Su, J.-K.; Lee, R.; Chang, C.; Lin, M.-L.; Mou, C.-Y. (2004) Gadolinium(III)-Incorporated Nanosized Mesoporous Silica as Potential Magnetic Resonance Imaging Contrast Agents. *J. Phys. Chem. B.* 108, 15608-15611.
- (22) Rieter, W. J.; Kim, J. S.; Taylor, K. M. L.; An, H.; Lin, W.; Tarrant, T.; Lin, W. (2007) Hybrid Silica Nanoparticles for Multimodal Imaging. *Angew. Chem. Int. Ed.* 119, 3754-3756.
- (23) Hak, S.; Sanders, H. M. H. F.; Agrawal, P.; Langereis, S.; Gröll, H.; Keizer, H. M.; Arena, F.; Terreno, E.; Strijkers, G. J.; Nicolay, K. (2009) A high relaxivity Gd(III)DOTA-DSPE-based liposomal contrast agent for magnetic resonance imaging. *Eur. J. Pharm. Biopharm.* 72, 397-404.
- (24) Terreno, E.; Delli Castelli, D.; Cabella, C.; Dastrù, W.; Sanino, A.; Stancanella, J.; Tei, L.; Aime, S. (2008) Paramagnetic Liposomes as Innovative Contrast Agents for Magnetic Resonance (MR) Molecular Imaging Applications. *Chem. Biodivers.* 5, 1901-1912.
- (25) Unger, E. C.; MacDougall, P.; Cullis, P.; Tilcock, C. (1989) Liposomal Gd-DTPA: Effect of encapsulation on enhancement of hepatoma model by MRI. *Magn. Reson. Imaging*, 7, 417-423.

- (26) Boswell, C. A.; Eck, P. K.; Regino, C. A. S.; Bernardo, M.; Wong, K. J.; Milenic, D. E.; Choyke, P. L.; Brechbiel, M. W. (2008) Synthesis, Characterization, and Biological Evaluation of Integrin  $\alpha v \beta 3$ -Targeted PAMAM Dendrimers. *Mol. Pharmaceutics*, **5**, 527-539.
- (27) Kobayashi, H.; Brechbiel, M. W. (2005) Nano-sized MRI contrast agents with dendrimer cores. *Adv. Drug Delivery Rev.* **57**, 2271-2286.
- (28) Langereis, S.; de Lussanet, Q. G.; van Genderen, M. H. P.; Meijer, E. W.; Beets-Tan, R. G. H.; Griffioen, A. W.; van Engelshoven, J. M. A.; Backes, W. H. (2006) Evaluation of Gd(III)DTPA-terminated poly(propylene imine) dendrimers as contrast agents for MR imaging. *NMR Biomed.* **19**, 133-141.
- (29) Langereis, S.; Dirksen, A.; Hackeng, T. M.; van Genderen, M. H. P.; Meijer, E. W. (2007) Dendrimers and magnetic resonance imaging. *New J. Chem.* **31**, 1152-1160.
- (30) Rudovský, J.; Botta, M.; Hermann, P.; Hardcastle, K. I.; Lukeš, I.; Aime, S. (2006) PAMAM Dendrimeric Conjugates with a Gd-DOTA Phosphinate Derivative and Their Adducts with Polyaminoacids: The Interplay of Global Motion, Internal Rotation, and Fast Water Exchange. *Bioconjugate Chem.* **17**, 975-987.
- (31) Bolskar, R. D. (2008) Gadofullerene MRI contrast agents. *Nanomedicine*, **3**, 201-213.
- (32) Zhu, W.; Okollie, B.; Bhujwalla, Z. M.; Artemov, D. (2008) PAMAM dendrimer-based contrast agents for MR imaging of Her-2/neu receptors by a three-step pretargeting approach. *Magn. Reson. Med.* **59**, 679-685.
- (33) Duarte, M. G.; Gil, M. H.; Peters, J. A.; Colet, J. M.; Elst, L. V.; Muller, R. N.; Geraldes, C. F. G. C. (2001) Synthesis, Characterization, and Relaxivity of Two Linear Gd(DTPA)-Polymer Conjugates. *Bioconjugate Chem.* **12**, 170-177.
- (34) Brasch, R. C. (1991) Rationale and applications for macromolecular Gd-based contrast agents. *Magn. Reson. Med.* **22**, 282-287.
- (35) Kobayashi, H.; Kawamoto, S.; Saga, T.; Sato, N.; Hiraga, A.; Ishimori, T.; Akita, Y.; Mamede, M. H.; Konishi, J.; Togashi, K., et al. (2001) Novel liver macromolecular MR contrast agent with a polypropylenimine diaminobutyl dendrimer core: Comparison to the vascular MR contrast agent with the polyamidoamine dendrimer core. *Magn. Reson. Med.* **46**, 795-802.
- (36) Kobayashi, H.; Kawamoto, S.; Jo, S. K.; Sato, N.; Saga, T.; Hiraga, A.; Konishi, J.; Hu, S.; Togashi, K.; Brechbiel, M. W., et al. (2002) Renal tubular damage detected by dynamic micro-MRI with a dendrimer-based magnetic resonance contrast agent. *Kidney Int.* **61**, 1980-1985.
- (37) Sugaya, M.; Watanabe, T.; Yang, A.; Starost, M. F.; Kobayashi, H.; Atkins, A. M.; Borris, D. L.; Hanan, E. A.; Schimel, D.; Bryant, M. A., et al. (2005) Lymphatic dysfunction in transgenic mice expressing KSHV k-cyclin under the control of the VEGFR-3 promoter. *Blood*, **105**, 2356-2363.
- (38) Wiener, E. C.; Konda, S.; Shadron, A.; Brechbiel, M.; Gansow, O. (1997) Targeting dendrimer-chelates to tumors and tumor cells expressing the high-affinity folate receptor. *Invest. Radiol.* **32**, 748-754.
- (39) Kobayashi, H.; Brechbiel, M. W. (2003) Dendrimer-based Macromolecular MRI Contrast Agents: Characteristics and Application. *Mol. Imaging*, **2**, 1-10.
- (40) Venditto, V. J.; Regino, C. A. S.; Brechbiel, M. W. (2005) PAMAM Dendrimer Based Macromolecules as Improved Contrast Agents. *Mol. Pharm.* **2**, 302-311.
- (41) Kumar, K.; Chang, C. A.; Francesconi, L. C.; Dischino, D. D.; Malley, M. F.; Gougoutas, J. Z.; Tweedle, M. F. (1994) Synthesis, Stability, and Structure of Gadolinium(III) and Yttrium(III) Macrocyclic Poly(amino carboxylates). *Inorg. Chem.* **33**, 3567-3575.
- (42) Reichert, D. E.; Hancock, R. D.; Welch, M. J. (1996) Molecular Mechanics Investigation of Gadolinium(III) Complexes. *Inorg. Chem.* **35**, 7013-7020.
- (43) Dong, Q.; Hurst, D. R.; Weinmann, H. J.; Chenevert, T. L.; Londy, F. J.; Prince, M. R. (1998) Magnetic Resonance Angiography With Gadomer-17: An Animal Study. *Invest Radiol.* **33**, 699-708.
- (44) Nicolle, G. M.; Tóth, É.; Schmitt-Willich, H.; Radüchel, B.; Merbach, A. E. (2002) The Impact of Rigidity and Water Exchange on the Relaxivity of a Dendritic MRI Contrast Agent. *Chem. Eur. J.* **8**, 1040-1048.
- (45) Behr, T. M.; Goldenberg, D. M.; Becker, W. (1998) Reducing the renal uptake of radiolabeled antibody fragments and peptides for diagnosis and therapy: present status, future prospects and limitations. *Eur J Nucl Med.* **25**, 201-212.
- (46) Ma, H. L.; Xu, Y. F.; Qi, X. R.; Maitani, Y.; Nagai, T. (2008) Superparamagnetic iron oxide nanoparticles stabilized by alginate: Pharmacokinetics, tissue distribution, and applications in detecting liver cancers. *Int. J. Pharm.* **354**, 217-226.
- (47) Lux, F.; Mignot, A.; Mowat, P.; Louis, C.; Dufort, S.; Bernhard, C.; Denat, F.; Boschetti, F.; Brunet, C.; Antoine, R., et al. (2011) Ultrasmall Rigid Particles as Multimodal Probes for Medical Applications. *Angew. Chem. Int. Ed.* **123**, 12507-12511.
- (48) Iznaga-Escobar, N.; Arocha, L. A. T.; Morales, A. M.; Suzarte, M. R.; Mesa, N. R.; Rodríguez, R. P. (1998) Technetium-99m-Antiepidermal Growth Factor-Receptor Antibody in Patients with Tumors of Epithelial Origin: Part II. Pharmacokinetics and Clearances. *J. Nucl. Med.* **39**, 1918-1927.
- (49) Vologdin, N.; Rolla, G. A.; Botta, M.; Tei, L. (2013) Orthogonal synthesis of a heterodimeric ligand for the development of the Gd-III-Ga-III ditopic complex as a potential pH-sensitive MRI/PET probe. *Org. Biomol. Chem.* **11**, 1683-1690.
- (50) Notni, J.; Hermann, P.; Dregely, I.; Wester, H.-J. (2013) Simultaneous in vivo PET/MRI imaging with a bi-metal bimodal tracer: Towards Ga-68-labeled responsive gadolinium contrast agents. *J. Labelled Comp. Radiopharm.* **56**, S62-S62.
- (51) Suchy, M.; Bartha, R.; Hudson, R. H. E. (2013) "Click" chemistry toward bis(DOTA-derived) heterometallic complexes: potential bimodal MRI/PET(SPECT) molecular imaging probes. *R. Soc. Chem. Adv.* **3**, 3249-3259.
- (52) Liu, W.; Hao, G. Y.; Long, M. A.; Anthony, T.; Hsieh, J. T.; Sun, X. K. (2009) Imparting Multivalency to a Bifunctional Chelator: A Scaffold Design for Targeted PET Imaging Probes. *Angew. Chem. Int. Ed.* **48**, 7346-7349.

Insert Table of Contents artwork here

