

Featured Article

## Cu-catalyzed O-Difluoromethylation of Functionalized Aliphatic Alcohols: An Access to Complex Organic Molecules with OCF<sub>2</sub>H group.

Kostiantyn Levchenko, Olexandr P. Datsenko, Oleh Serhiichuk,  
Andrei Tolmachev, Viktor O. Iaroshenko, and Pavel K. Mykhailiuk

*J. Org. Chem.*, **Just Accepted Manuscript** • DOI: 10.1021/acs.joc.6b00628 • Publication Date (Web): 06 May 2016

Downloaded from <http://pubs.acs.org> on May 6, 2016

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



# Cu-catalyzed O-Difluoromethylation of Functionalized Aliphatic Alcohols: An Access to Complex Organic Molecules with OCF<sub>2</sub>H group.


Kostiantyn Levchenko,<sup>†</sup> Olexandr P. Datsenko,<sup>†</sup> Oleh Serhiichuk,<sup>§</sup> Andrei Tolmachev,<sup>†</sup> Viktor O. Iaroshenko,<sup>⊥\*</sup> Pavel K. Mykhailiuk<sup>†‡\*</sup>

<sup>†</sup>Enamine Ltd., 78 Chervonotkatska str., 02094, Kyiv, Ukraine, [www.enamine.net](http://www.enamine.net)

<sup>‡</sup>Department of Chemistry, Taras Shevchenko National University of Kyiv, Volodymyrska Street, 64, Kyiv 01601, Ukraine.

<sup>§</sup>Department of Organic Chemistry, Kiev Polytechnic Institute, Pr. Pobedy 37, 03056 Kiev, Ukraine.

<sup>⊥</sup> Homogeneous Catalysis and Molecular Design Research Group at Center of Molecular and Macromolecular Studies, Polish Academy of Sciences, Sienkiewicza 112, PL-90-363 Łódź, Poland.




**ABSTRACT:** Two-step synthetic strategy towards the difluoromethyl ethers *via* a CuI-catalyzed reaction of the alcohols, bearing additional protected functionalities, with FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>H has been developed. The high potential of the developed protocol has been shown by preparing novel OCF<sub>2</sub>H-analogues of GABA and *L*-Proline. The described transformation has good functional group compatibility and can serve as a powerful synthetic tool for late-stage preparation of complex OCF<sub>2</sub>H-containing organic compounds as well as building blocks for drug discovery.

## INTRODUCTION

The isosteric substitution of hydrogen atoms with fluorine or introduction of fluorine-bearing functionalities into complex organic molecules are well-established strategies in medicine and life science for the design and optimization of biologically active molecules, as well as in material science for devising new functional materials.<sup>1,2</sup> The fluorine atom is similar in size to hydrogen. This

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

feature is accepted to be the main reason why the biological objects, like microorganisms and enzymes, under particular circumstances, do not recognize the difference between a natural substrate and its fluorine-modified analogue, obtained *via* the replacement of the particular C-H bond by C-F bond. This phenomenon is often considered as the “mimic effect” of fluorine for hydrogen.<sup>1b</sup> However, the C-F bond comparing with C-H has much stronger bonding energy which results in the increased stability of the C-F bond. These structural alterations are the main reasons for the changes in the metabolic pathways of fluorinated compounds and in particular augmentation of the *in vivo* stability towards various enzymes. For instance, introduction of a fluorine into aromatic core is a well-known strategy to improve the stability of the lead compounds, for instance towards the Cytochrome C enzymes of liver cells, which dramatically changes the clearance and longevity of the drug candidate *in vivo*.<sup>3</sup> Nonetheless, twenty seven among worldwide 150-best-selling drugs in the year 2013 are fluorinated compounds.<sup>1b,d</sup>



**Figure 1.** Pharmacologically relevant aromatic OCF<sub>2</sub>H compounds.


31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41

The effects associated with stereo-electronic interplays of fluorine atom with other fluorines or fluorine-containing groups, as well as with various polar or charged functional groups including, oxygen and nitrogen, are commonly used in the medicinal chemistry<sup>1c,d,4</sup> and in the catalytic science for the development of pharmacologically significant substances, catalysts<sup>5,6</sup> and/or catalytic moieties with enhanced properties, edited efficacy and performance. In medicinal chemistry, however, these particular tools deliver a practical handle for “fine-tune” of biological and physicochemical profiles of lead structures.<sup>1c,d</sup> In this context, several fluorine modifications of naturally occurring and pharmacologically relevant structures were undertaken.<sup>1b</sup>

42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

In a continuation of our ongoing medicinal chemistry program dedicated to engineering of fluorinated pharmacologically relevant scaffolds,<sup>7</sup> we have undertaken the current study aiming the synthesis of OCF<sub>2</sub>H-containing complex organic molecules. Our choice of this group is rather not at random, but was stipulated by a pivotal role this functionality plays as a pharmacophore in the marketed drugs;<sup>3</sup> for instance, such well-known medicines as Roflumilast **I**<sup>1b,c,3</sup> and Pantoprazole **II**<sup>1b,c,3</sup> contain this group (Figure 1). Besides, the OCF<sub>2</sub>H group is a commonly occurring structural motif found in numerous pharmaceuticals and biologically active compounds<sup>8</sup> including enzyme inhibitors, anti-HIV agents and antimicrobial agents. In the life science this group is presented by several OCF<sub>2</sub>H containing nucleosides,<sup>1c,d</sup> having visible impact on the conformational preferences of the nucleosides themselves and the DNA double strand in particular. In addition to that, the OCF<sub>2</sub>R functionalities are commonly used for the stereocontrol of

rotational profile<sup>9</sup> in organic molecules of different types. The elevated interest to the difluoromethyl group can be also explained by the fact that it acts as bioisostere of alcohols and thiols.<sup>1b,4</sup> This makes the OCF<sub>2</sub>R groups and OCF<sub>2</sub>H in particular as desirable pharmacophores and valuable tool to tune the properties of organic molecules and their conformational bias.



**Table 1.** List of used starting materials.

Analyzing the OCF<sub>2</sub>H drugs presently available on the market as well as the drug candidates in clinical and preclinical studies, we witnessed the lack of aliphatic OCF<sub>2</sub>H molecules. In a view of recent trends in the field of medicinal chemistry related to the design of architecturally more complex molecules with greater three-dimensionality (3D-shape), *via* the formal saturation of aromatic precursors,<sup>10</sup> the functionalized OCF<sub>2</sub>H aliphatic building blocks with high degree of saturation are highly desirable scaffolds for drug discovery. The deficit of OCF<sub>2</sub>H aliphatic structural motives is a result of underdeveloped synthetic methodologies enabling the preparation of such scaffolds.


Concerning the synthetic routes known to date for the furnishing of organic compounds with the OCF<sub>2</sub>H group, there are four main strategies. Namely, (i) the O-difluoromethylation<sup>11</sup> of phenols and alcohols using various sources of difluorocarbene; (ii) the C-O coupling reaction using the chemical equivalents<sup>12</sup> of the [OCF<sub>2</sub>H]<sup>•</sup> synthon; (iii) the transformation of CO and CS groups;<sup>13</sup> (iv) the nucleophilic exchange<sup>14</sup> of other halogens to fluorine in the OCHal<sub>2</sub>H groups. Concerning the strategy (i), there are a number of sources amenable for generation of difluorocarbene; for instance, such chemical synthons as HalCF<sub>2</sub>CO<sub>2</sub>Na; FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>TMS; FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>H; TMSCF<sub>2</sub>R (R = F, Cl, Br); (Et<sub>2</sub>O)<sub>2</sub>POCF<sub>2</sub>Br; CF<sub>3</sub>SO<sub>3</sub>CF<sub>2</sub>H etc. are commonly used for this purpose.<sup>11a</sup> However,

the FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>H due to its high reaction efficiency, low price and synthetic availability, is often considered to be the optimal reagent of choice for O-difluoromethylation, in particular for large-scale synthesis. Despite the diversity of the strategies known to date, the most commonly used synthetic route towards difluoromethyl ethers from alcohols requires the use of ozone-depleting HCF<sub>2</sub>Cl, the so-called Freon 22.<sup>15</sup> The environment friendly preparative methods, however, often require elevated temperatures, are not compatible with many functional groups and are only efficient on the structurally simple substrates.<sup>11</sup> Due to the aforementioned limitations of presently used methodologies, the development of new, simple, straightforward and concise methods, to convert functional alcohols to difluoromethyl ethers, are of considerable interests.

Herein, we report a practical entry towards structurally diverse difluoromethyl ethers, using the set of readily available alcohols, bearing an additional functional group, and FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>H<sup>11</sup> as a source CF<sub>2</sub> structural motif. Our methodology commenced with the preparation of the protected substrates **1a-s**. For the current study we selected O-protected aliphatic diols **1a,b**; N-protected amino alcohols **1c-n**; carboxyl-protected hydroxyl acids **1m-q**, as well as N- and carboxyl- protected amino acids bearing free hydroxyl group **1r,s** (Table 1). The method we are communicating here is an extension of the methodology for non-functionalized alcohols developed by Chen at al.<sup>16</sup> that in our case covers the functionalized aliphatic alcohols and results in improved reaction efficiency and overall yields. Within the current research program we have developed the diverse libraries of complex OCF<sub>2</sub>H-containing organic compounds, however here as a quintessential we are communicating only 18 examples (Table 3).

## RESULTS AND DISCUSSION

With the set of the starting compounds **1a-s** in hand (Table 1), we focused on the setting up the standard reaction condition applicable for the title transformation (Scheme 1, Table 2). We pointed out the reaction conditions that were used in the early works of Chen<sup>17</sup> as a starting point of our research. Noteworthy, these conditions require from 2- to 3-fold excess of alcohol along with elevated temperatures, and in our opinion cannot be used for the multi gram preparation of structurally diverse OCF<sub>2</sub>H compounds. At the same time, many synthetic methods known to date utilise strong bases for the difluorocarbene generation; these methods usually demonstrate low functional group tolerance and cannot be employed for the late-stage functionalization of complex organic molecules.



**Scheme 1.** Synthetic scenario for the preparation of OCF<sub>2</sub>H substituted intermediates and final PG-free products.

Recent studies showcased that the transition metals enable the transfer of the carbene moieties by formation of corresponding carbene complexes. Additionally, the existence of such adducts and their stability were supported by spectroscopic means along with the chemical transformations of the aforementioned metallic species.<sup>18</sup> The difluorocarbene complexes of transition metals are studied, however their existence as intermediates was proven in several catalytic processes.<sup>19</sup> This in particular turned our attention towards transition-metal-mediated protocols and prompted us to start the reaction trials applying Cu(I) salts, which are known to be capable to trap carbenes forming suitable carbene complexes.<sup>18a,d</sup> On the other hand, the recent progress on the application of CuCF<sub>3</sub> as an intermediate of choice for the catalytic trifluoromethylation as well as difluoromethylation<sup>20</sup> brought us on the idea that the O-difluoromethylation can also follow the second possible mechanistic route, namely *via* the formation of a CF<sub>3</sub> metal species. Since we were not sure about the mechanism the reaction undergoes within the designed and tested reaction conditions, we decided to consider both possible routes mentioned above. Accordingly, we have selected a set of transition metals capable for formation of metal carbene complexes and CF<sub>3</sub> metal species.

**Table 2.** Optimization of reaction conditions for the synthesis of **2a**.<sup>a</sup>



entry	catalyst/solvent	T	time	yield (%) <sup>b</sup>
1	CuBr (0.2 eq)/C <sub>6</sub> H <sub>6</sub>	60°C	45min	traces
2	CuBr (0.2 eq)/C <sub>6</sub> H <sub>6</sub>	60°C	3h	traces
3	CuBr (0.2 eq)/C <sub>6</sub> H <sub>6</sub>	100°C	3h	traces
4	CuBr (0.2 eq)/MeCN	60°C	2h	5 <sup>c</sup>
5	CuI (0.2 eq)/C <sub>6</sub> H <sub>6</sub>	50°C	1h	17 <sup>c</sup>
6	CuI (0.2 eq)/C <sub>6</sub> H <sub>6</sub>	90°C	2h	12 <sup>c</sup>
7	CuI (0.2 eq)/MeCN	50°C	2h	67 (75 <sup>c</sup> )
8	CuI (0.2 eq)/MeCN	80°C	2h	60 (69 <sup>c</sup> )

9	CuI (0.2 eq)/MeCN	50°C	45min	73 (82 <sup>c</sup> )
10	CuI (0.1 eq)/MeCN	50°C	45min	58 <sup>c</sup>
11	CuI (0.3 eq)/MeCN	50°C	45min	81 <sup>c</sup>
12	NiBr <sub>2</sub> (0.2 eq)/MeCN	60°C	1h	15 <sup>c</sup>
13	NiI <sub>2</sub> (0.2 eq)/MeCN	60°C	1h	19 <sup>c</sup>
14	Cu(OAc) <sub>2</sub> (0.2 eq)/MeCN	60°C	1h	-
15	Cu(OTf) <sub>2</sub> (0.2 eq)/MeCN	60°C	1h	25 <sup>c</sup>
16	Rh <sub>2</sub> (OAc) <sub>4</sub> (0.1 eq)/C <sub>6</sub> H <sub>6</sub>	60°C	3h	45 <sup>c</sup>
17	Rh <sub>2</sub> (O <sub>2</sub> CCF <sub>3</sub> ) <sub>4</sub> (0.1 eq)/C <sub>6</sub> H <sub>6</sub>	60°C	3h	38 <sup>c</sup>


<sup>a</sup> Optimization of the reaction condition was performed on the 10 mmol scale taking 1.5 equivalents of FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>H.

<sup>b</sup> Yields.

<sup>c</sup> GC/MS detected yields.

The substance **1a** was chosen as a model compound. After reaction condition optimization, consisted in variation of solvents, amount and type of transition metals, temperature and time of reaction, the optimal reaction condition was developed (Table 2 entry 9). We mainly focused on the exploration of CuBr and CuI as catalysts enabling introduction of CF<sub>2</sub>H group. First, the CuBr was checked (Table 2, entries 1-4); accordingly, to our great disappointment we did observe the formation of desired product but in trace amounts. The best result showcased by utilizing the combination of CuBr with MeCN as a solvent. Under these conditions the product **2a** was detected by GC/MS in 5% yield. Furthermore, the CuI/C<sub>6</sub>H<sub>6</sub> system improved the situation dramatically, delivering the desired product in 17% GC/MS-yield (Table 2, entry 5). Nonetheless, these results were not sufficient enough to deploy a chemistry using these reaction conditions. The crucial improvement in terms of yields appeared to be the usage of MeCN as a solvent, which increased the GC/MS-yield up to 75% (Table 2, entry 7). The following crucial parameters during the reaction condition optimization were the screening of time and the temperature. We encountered with the problem of the thermal instability of the product **2a** under the Cu-catalysis (Table 2, entry 8). This observation prompted us to decrease the reaction time up to 45 minutes and the reaction temperature to 50 °C. Finally, we reached the synthesis of the corresponding difluoromethyl ether **2a** in 82% GC/MS-yield and 73% isolated yield (Scheme 1, Table 2). The best yield was obtained using 1.5 equivalents of difluoromethylation agent (FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>H) and 0.2 equivalents of CuI, in MeCN as reaction media at 50 °C (Table 2, entry 9). Further, increasing

the amount of CuI to 0.3 equivalents did not improve visibly the efficiency of the title reaction. However, when we took 0.1 equivalent of CuI, the GC/MS-yield dropped significantly (entry 10). At the same time, other transition metals, such as Ni(II) (Table 2, entries 12, 13) and Cu(II) (Table 2, entries 14, 15) salts, were not prone to catalyze the title reaction in a sufficient manner. In contrast, the Rh(II) salts, known previously to promote the formation of carbene metalocomplexes,<sup>18c</sup> demonstrated promising results when applied to the title reaction (Table 2, entries 16, 17). However, owing the availability and overall costs of Rh-based catalysts, we excluded this option for the further optimization of reaction condition basing on the goal of the gram scale production of desired OCF<sub>2</sub>H substituted substances.




**Table 3.** The scope of the OCF<sub>2</sub>H intermediates as well as final protecting-group-free products.

Prompted by these results and having optimized reaction conditions, we turned our attention towards evaluation of the generality of the developed synthetic protocol. We found that the O-protected diol **1b** under the aforementioned reaction conditions can be swift-

ly converted to the corresponding protected intermediate compound **2b** (Table 3). The same results were observed for a variety of N-protected amino alcohols **1c-n** and carboxyl-protected hydroxyl acids **1o-q** (Table 1 and 3). To our delight, the developed reaction conditions appeared to be general and allowed the synthesis of corresponding difluoromethyl ethers **2a-q** bearing various functionalities. From the results summarized in Table 3, it is evident that the described CuI-catalyzed difluoromethylation reaction turned out to be rather general. The obtained compounds were either high viscous oils or liquids; thus the purification was accomplished by distillation in vacuum or preparative column chromatography on the silica gel. In the case of compounds **2e**, **2j** and **3c** we observed the drop in the yields. For **2e** and **2j** most probably this can be addressed to the sterical bulkiness of the corresponding substrates **1e**, **1j**. The rather low yield of the amine **3c** can be explained by its high volatility. Upon scale-up production of compounds **2d**, **2l**, **2m**, **2r** and **2s** the amount of FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>H could be decreased up to 1.2-1.3 equivalents without visible loss in yields.

The removal of the concrete protecting group was achieved following the standard reaction condition commonly used for the given case (Scheme 1). For instance, the Bn and CO<sub>2</sub>Bn moieties, present in the compounds **2a-2q**, were easily removed from corresponding ethers and amides by the high pressure catalytic hydrogenation. The reaction was conducted in methanol, using 5%-Pd/C as catalyst under 30 atm of H<sub>2</sub> at room temperature. The title reaction can be easily scaled-up on gram- and decagram-scales without further optimization. Of note, the mildness of the developed methodology enabled the synthesis of enantiomerically pure compounds **3l** and **3m** from the corresponding (*S*) and (*R*) precursors without any notable racemization.



**Scheme 2.** Follow-up chemistry: Synthesis of (*S*)-3-(difluoromethoxy)pyrrolidin-2-one **5** by base-promoted cyclization of (*S*)-methyl 4-amino-2-(difluoromethoxy)butanoate **4**.

In the light of these results, we anticipated that the title reaction could be also applicable for the late-stage diversification of pharmacologically relevant compounds, for instance, hydroxy amino acids. In a view of our current interests in the fluorine modified natural and semi-natural products<sup>7</sup> like amino acids and nucleosides, for this study two model amino acids were selected; namely, the (*4R*)-4-hydroxy-L-proline and (*2S*)-2-hydroxy GABA ((*2S*)-4-amino-2-hydroxybutanoic acid). Following the known synthetic methods,<sup>21</sup> these compounds were converted into the protected derivatives **1r** and **1s**, which were suitable for the subsequent CF<sub>2</sub>-functionalization (Table 1). Using the optimal conditions, the corresponding compounds were successfully converted into intermediate structures **2r** and **2s** in good yields (80% and 84% respectively). The further saponification of these amino acid esters by solu-

tion of LiOH in THF afforded N-Boc-protected (4*R*)-4-(difluoromethoxy)-L-proline **3r** and N-t-Boc-protected (2*S*)-4-amino-2-(difluoromethoxy)butanoic acid **3s** in 61% and 58% yield respectively. The Boc and t-Boc protecting groups on the amino function were not removed, since according to our actual research plans those building blocks will be used further for the peptide synthesis. The attempts to incorporate the synthesized OCF<sub>2</sub>H-analogue of *L*-Proline into the oligo- and polypeptide strings<sup>7a-c, 22</sup> are currently underway in our laboratories.

In order to illustrate the follow-up chemistry, the synthetic transformation of the **2r** was undertaken. The cyclization of the protected (2*S*)-2-hydroxy GABA was achieved by the treatment of the title compound with TFA followed by the basic neutralization with K<sub>2</sub>CO<sub>3</sub>, which afforded to the pharmacologically relevant (3*S*)-3-(difluoromethoxy)pyrrolidin-2-one **5** in 52% yield as a pure (*S*)-enantiomer (Scheme 2). The racemization in the course of the reaction was not detected. At the same time, in order to cut off two extra steps related with protection and deprotection, we had a notion to set under investigation the unprotected hydroxy amino substrates including hydroxy amino acids. Unfortunately, these attempts experienced a failure, the corresponding reactions led to the formation of a mixture of unidentified products accompanied by the tar formation.

To confirm the constitution of the obtained structural scaffolds **2**, **3** and **5** we mainly used the NMR means. The characteristic features of the OCF<sub>2</sub>H products in <sup>1</sup>H and <sup>13</sup>C NMR spectra are accompanied by the equal volume of corresponding *J*<sub>C-F</sub>, *J*<sub>F-H</sub> coupling constants in proton and <sup>19</sup>F NMR spectra. For the structures with the chiral carbon in the close proximity to the OCF<sub>2</sub>H group we observed the diastereotopism of two geminal fluorine atoms with the characteristic spin pictures and coupling patterns.<sup>23</sup> The chemical shifts of OCF<sub>2</sub>H function is in accordance to the previously communicated data.<sup>23</sup>

## SUMMARY

In summary, we developed operationally simple protocol for O-difluoromethylation of various functional alcohols with 1.2-1.5 excess amount of FSO<sub>2</sub>CF<sub>2</sub>CO<sub>2</sub>H which is so far an optimal difluoromethylation reagent. The CuI-catalyzed reaction of the polyfunctional alcohols provides corresponding difluoromethyl ethers from moderate to high yields. Furthermore, this protocol is distinguished by a wide substrate scope, including aliphatic alcohols bearing additional protected OH, NH<sub>2</sub>, or CO<sub>2</sub>H groups, without compromising its efficiency and scalability. As the following task to be solved in the frames of this research program, we see the extension of the developed protocol on the synthesis of other OCF<sub>2</sub>H-modified amino acids and natural products as well as subsequent assembly of OCF<sub>2</sub>H-analogues of several pharmacologically relevant peptides. At the same time, we are intended to disclose the mechanism of the title CuI-catalyzed reaction by spectroscopic (1D, 2D NMR, IR) and rational means. These attempts are currently underway in our laboratories.

**EXPERIMENTAL SECTION**

Solvents were purified according to standard procedures. Column chromatography was performed using Kieselgel Merck 60 (230-400 mesh) as the stationary phase.  $^1\text{H}$ -,  $^{19}\text{F}$ -,  $^{13}\text{C}$  NMR spectra were recorded on at 499.9, 470.3 and 124.9 MHz. Chemical shifts are reported in ppm downfield from TMS ( $^1\text{H}$ ,  $^{13}\text{C}$ ) or  $\text{CFCl}_3$  ( $^{19}\text{F}$ ) as internal standards. The information associated with preparation of the compounds **1a-s** one can find in the SI.

**((4-(Difluoromethoxy)butoxy)methyl)benzene (2a).**

Compound **1a** (15 g, 0.083 mol, 1 eq) was dissolved in 200 mL of acetonitrile and copper iodide (3.16 g, 0.017 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (22.3 g, 0.125 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the solid phase was filtered out. The EtOAc solution was concentrated under vacuum to give the product **2a**. Yield = 73 % (14 g).

The product is a transparent colorless liquid.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 1.71-1.74 (m, 4H), 3.20, 3.49 (s, 2H), 3.85, 4.18 (t, 2H,  $^3J$  = 5.4 Hz), 4.49 (s, 2H), 6.16 (t, 1H,  $^2J_{\text{H-F}}$  = 75.3 Hz,  $\text{CF}_2\text{H}$ ), 7.27-7.32 (m, 5H).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta$  = -842 (d, 2F,  $^2J$  = 76.0 Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 25.9, 26.0, 63.4 (t,  $J$  = 5.9 Hz), 69.5, 72.8, 116.1 (t,  $^1J_{\text{C-F}}$  = 259.0 Hz,  $\text{CF}_2\text{H}$ ), 127.4, 127.5, 128.3, 138.5.

Anal. calcd for  $\text{C}_{12}\text{H}_{16}\text{F}_2\text{O}_2$ : C, 62.60; H, 7.00. Found: C, 62.74; H, 7.07.

**4-(Difluoromethoxy)butan-1-ol (3a).**

Compound **2a** (14 g, 0.061 mol, 1 eq) was dissolved in 100 mL of methanol and 1.5g of 5% -Pd/C was added thereto. The mixture was stirred under 30 atm of  $\text{H}_2$  for 96 h at room temperature. Then Pd/C was filtered out and the resulting mixture was concentrated under vacuum to give compound **3a**. Yield = 74% (6.3 g).

The product is a pale liquid.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 1.60-1.73 (m, 4H), 2.78 (br. s, 1H), 3.63 (t, 2H,  $^3J$  = 6.8 Hz), 3.85 (t, 2H,  $^3J$  = 6.7 Hz), 6.17 (t, 1H,  $^2J_{\text{H-F}}$  = 73.3 Hz,  $\text{CF}_2\text{H}$ ).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta$  = -84.5 (d, 2F,  $^2J_{\text{H-F}}$  = 75.1 Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 25.1, 29.3, 62.1, 63.0 (t,  $J$  = 5.5 Hz), 115.7 (t,  $^1J_{\text{C-F}}$  = 256.3 Hz,  $\text{CF}_2\text{H}$ ).

Anal. calcd for  $\text{C}_5\text{H}_{10}\text{F}_2\text{O}_2$ : C, 42.86; H, 7.19. Found: C, 42.72; H, 7.22.

**(((4-(Difluoromethoxy)cyclohexyl)oxy)methyl)benzene (2b) (Mixture of diastereomers).**

Compound **1b** (28.8 g, 0.14 mol, 1 eq) was dissolved in 300 mL of acetonitrile and copper iodide (5.3 g, 0.028 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (37.3 g, 0.21 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solid was filtered out. The resulting solution of EtOAc was evaporated and the crude of **2b** was purified by column chromatography (EtOAc : Hexane - 1 : 6,  $R_f = 0.65$ ). Yield = 73 % (27.6g). We encountered with the problems related to the prefabrication of the compound **2b**, however the impurities of the batch did not effected the purity of the final product **3b**.

The product is a transparent colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta = 1.15\text{-}1.19$  (m, 2H),  $1.69\text{-}1.72$  (m, 2H),  $1.95\text{-}1.99$  (m, 2H),  $2.07\text{-}2.14$  (m, 2H),  $3.50\text{-}3.53$  (m, 1H),  $4.20\text{-}4.25$  (m, 1H),  $4.58$  (s, 2H),  $6.27$  (dt, 1H,  $^2J_{\text{H-F}} = 75.6$  Hz,  $J = 9.0$  Hz,  $\text{CF}_2\text{H}$ ),  $7.33\text{-}7.34$  (m, 1H),  $7.39\text{-}7.40$  (m, 4H).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -81.1$  (t, 2F,  $^2J_{\text{H-F}} = 77.0$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{DMSO-}d_6$ ) (mixture of diastereomers):  $\delta = 27.4, 28.2, 28.4, 29.4, 69.3, 69.7, 74.3, 75.0, 117.7$  (dt,  $^1J_{\text{C-F}} = 258.5$  Hz,  $J = 12.2$  Hz,  $\text{CF}_2\text{H}$ ),  $127.6, 127.7, 128.6, 139.6, 139.6$  (d,  $J = 14.0$  Hz).

Anal. calcd for  $\text{C}_{14}\text{H}_{18}\text{F}_2\text{O}_2$ : C, 65.61; H, 7.08. Found: C, 65.43; H, 7.11.

#### 4-(Difluoromethoxy)cyclohexanol (**3b**) (Mixture of diastereomers).

Compound **2b** (25.5 g, 0.1 mol) was dissolved in 200 mL of methanol and 3.5 g of 5%-Pd/C was added thereto. The mixture was stirred under 30 atm of  $\text{H}_2$  for 96 h at room temperature. Then Pd/C was filtered out and the resulting mixture was concentrated under vacuum. The crude product was distilled under vacuum (1 Torr, 54 °C) to give compound **3b**. Yield = 64.7% (10.7 g).

The product is a pale liquid.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta = 1.25\text{-}1.71$  (m, 4H),  $1.75$  (s, 1H),  $1.90\text{-}2.02$  (m, 4H),  $3.73\text{-}3.75$  (m, 1H),  $4.18$  (m, 1H),  $6.22$  (dt, 1H,  $^2J_{\text{H-F}} = 75.9$  Hz,  $J = 7.8$  Hz,  $\text{CF}_2\text{H}$ ).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta = 28.2, 29.3, 29.8, 31.5, 35.1, 68.2, 72.6, 115.9$  (t,  $^1J_{\text{C-F}} = 259.6$  Hz,  $\text{CF}_2\text{H}$ ).

Anal. calcd for  $\text{C}_7\text{H}_{12}\text{F}_2\text{O}_2$ : C, 50.60; H, 7.28. Found: C, 50.39; H, 7.25.

#### Benzyl (2-(difluoromethoxy)ethyl)carbamate (**2c**).

Compound **1c** (6 g, 0.0307 mol, 1 eq) was dissolved in 60 mL of acetonitrile and copper iodide (1.15 g, 0.006 mol, 0.2 eq) was added thereto. The reaction mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (8.2 g, 0.046 mol, 1.5 eq) in 10 mL of acetonitrile was added dropwise over a period of 30 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solids were filtered

1 out. The resulting solution of EtOAc was evaporated and the crude of **2c** was purified by column chromatography (Hexane : EtOAc  
2 - 4 : 1,  $R_f = 0.65$ ). Yield = 86% (6.35 g).

3  
4 The product is a transparent colorless thick oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 3.40$  (br. s, 2H), 3.86 (m, 2H), 5.07 (s, 2H), 5.32  
5 (br. s, 1H), 6.15 (t, 1H,  $^2J_{\text{H-F}} = 71.4$  Hz,  $\text{CF}_2\text{H}$ ), 7.31 (s, 5H).

6  
7  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -84.6$  (d, 2F,  $^2J_{\text{H-F}} = 79.6$  Hz,  $\text{CF}_2\text{H}$ ).

8  
9  $^{13}\text{C}\{^1\text{H}\}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta = 40.2, 62.5, 66.9, 115.8$  (t,  $^1J_{\text{C-F}} = 260.0$  Hz,  $\text{CF}_2\text{H}$ ), 127.9, 128.0, 128.4, 136.3, 156.3.

10  
11 Anal. calcd for  $\text{C}_{11}\text{H}_{13}\text{F}_2\text{NO}_3$ : C, 53.88; H, 5.34; N, 5.71. Found: C, 54.00; H, 5.40; N, 5.65.  
12  
13

#### 14 15 **2-(Difluoromethoxy)ethanamine (3c).**

16  
17 Compound **2c** (6.35g, 0.026 mol, 1 eq) was dissolved in 65 mL of methanol and 0.6 g of 5% -Pd/C was added thereto. The mixture  
18 was stirred under 30 atm of  $\text{H}_2$  for 12 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated  
19 under atmospheric pressure. Product **3c** was purified by distilled under vacuum (18 Torr, 35 °C). Yield = 10% (0.29 g); the main  
20 reason of low yield is the high volatility of the product.  
21  
22

23  
24 The product is a transparent colorless liquid.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.22$  (s, 2H), 2.85 (t, 2H,  $^3J = 5.0$  Hz), 3.78 (t, 2H,  $^3J$   
25 = 5.0 Hz), 6.17 (t, 1H,  $^2J_{\text{H-F}} = 76.3$  Hz,  $\text{CF}_2\text{H}$ ).

26  
27  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -74.1$  (d, 2F,  $^2J_{\text{H-F}} = 74.7$  Hz).

28  
29  $^{13}\text{C}\{^1\text{H}\}$  NMR (125 MHz,  $\text{CDCl}_3$ ):  $\delta = 40.8, 65.5, 115.7$  (t,  $^1J_{\text{C-F}} = 258.8$  Hz,  $\text{CF}_2\text{H}$ ).

30  
31 Anal. calcd for  $\text{C}_3\text{H}_7\text{F}_2\text{NO}$ : C, 32.43; H, 6.35; N, 12.61. Found: C, 32.22; H, 6.28; N, 12.50.  
32  
33  
34  
35

#### 36 **Benzyl (3-(difluoromethoxy)propyl)carbamate (2d).**

37  
38 Compound **1d** (60 g, 0.287 mol, 1 eq) was dissolved in 600 mL of acetonitrile and copper iodide (10.95 g, 0.058 mol, 0.2 eq) was  
39 added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (61.3 g, 0.344 mol, 1.2  
40 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30  
41 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solids were filtered out. The  
42 resulting solution of EtOAc was evaporated and the crude of **2d** was purified by column chromatography (Hexane : EtOAc - 4 : 1,  
43  $R_f = 0.65$ ). Yield = 86% (63 g).  
44  
45

46  
47 The product is a transparent colorless thick oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.85$ -1.87 (m, 2H), 3.31 (br. s, 2H), 3.91 (m, 2H),  
48 4.99 (br. s, 1H), 5.10 (s, 2H), 6.20 (t, 1H,  $^2J_{\text{H-F}} = 72.4$  Hz,  $\text{CF}_2\text{H}$ ), 7.36 (s, 5H, Ph).

49  
50  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -81.8$  (dd, F,  $^2J_{\text{H-F}} = 76.8$  Hz), -81.0 (dd, F,  $^2J_{\text{H-F}} = 76.8$  Hz).

51  
52  $^{13}\text{C}\{^1\text{H}\}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta = 29.4, 38.00, 61.4, 66.8, 116.2$  (t,  $^1J_{\text{C-F}} = 261.3$  Hz,  $\text{CF}_2\text{H}$ ), 128.2, 128.6, 136.7, 156.7.

53  
54 Anal. calcd for  $\text{C}_{12}\text{H}_{15}\text{F}_2\text{NO}_3$ : C, 55.59; H, 5.83; N, 5.40. Found: C, 55.75; H, 5.87; N, 5.31.  
55  
56  
57  
58  
59  
60

**3-(Difluoromethoxy)propan-1-amine (3d).**

Compound **2d** (63 g, 0.243 mol, 1 eq) was dissolved in 300 mL of methanol and 5 g of 5% -Pd/C was added thereto. The mixture was stirred under 30 atm of H<sub>2</sub> for 36 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated under atmospheric pressure. Product **3d** was purified by distilled under vacuum (18 Torr, 43 °C). Yield = 72.6% (22.1 g).

The product is a transparent colorless liquid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 1.42 (s, 2H), 1.77 (quint, 2H, <sup>3</sup>J = 6.5 Hz), 2.81 (t, 2H, <sup>3</sup>J = 7.0 Hz), 3.92 (t, 2H, <sup>3</sup>J = 6.0 Hz), 6.18 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 75.2 Hz, CF<sub>2</sub>H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -84.6 (d, 2F, <sup>2</sup>J<sub>H-F</sub> = 76.1 Hz).

<sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ = 32.6, 38.6, 61.4, 116.0 (t, <sup>1</sup>J<sub>C-F</sub> = 272.0 Hz, CF<sub>2</sub>H).

Anal. calcd for C<sub>4</sub>H<sub>9</sub>F<sub>2</sub>NO: C, 38.40; H, 7.25; N, 11.19. Found: C, 38.55; H, 7.19; N, 11.13.

**Benzyl (2-hydroxy-2-methylpropyl)carbamate (1e).**

Benzyl chloroformate (CbzCl) (94.95 g, 0.557 mol, 1 eq) was added dropwise to the stirred solution of aminoalcohol (49.6 g, 0.557 mol, 1 eq) and potassium carbonate (153.8 g, 1.115 mol, 2 eq) in mixture of THF: water (800:400) at 0°C. Then the reaction mixture was stirred 18 hours at room temperature. After completion of the reaction, THF was rotary evaporated and the rest was extracted 3 times with DCM. DCM was dried with sodium sulfate and after filtration evaporated. Crude product was purified by flash-chromatography (EtOAc – petroleum ether 1:3). Yield = 71.9% (89.3 g).

The product is a transparent yellowish oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ = 1.19 (s, 6H), 2.66 (s, 1H), 3.17 (s, 2H), 5.09 (br. s, 2H), 5.45 (s, 1H), 7.33 (br. s, 5H).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 27.1, 51.8, 67.0, 71.0, 128.2, 128.3, 128.6, 136.6, 157.5.

Anal. calcd for C<sub>12</sub>H<sub>17</sub>NO<sub>3</sub>: C, 64.55; H, 7.67; N, 6.27; O, 21.50. Found: C, 64.34; H, 7.79; N, 6.18.

**Benzyl (2-(difluoromethoxy)-2-methylpropyl)carbamate (2e).**

Compound **1e** (89.3 g, 0.4 mol, 1 eq) was dissolved in 750 mL of acetonitrile and copper iodide (15.25 g, 0.08 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (106.85g, 0.6 mol, 1.5 eq) in 200 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solid was filtered out. The resulting solution of EtOAc was evaporated and the crude product **2e** was purified by column chromatography (Hexane : EtOAc - 3 : 1, R<sub>f</sub> = 0.6). Yield = 6.8% (7.4 g).

The product is a transparent colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 1.34 (s, 6H), 3.32 (d, 2H, <sup>3</sup>J = 6.7 Hz), 5.10 (br. s, 3H), 6.25 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 76.4 Hz, CF<sub>2</sub>H), 7.34 (br. s, 5H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -79.3 (d, 2F, <sup>2</sup>J<sub>H-F</sub> = 76.5 Hz).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 24.0, 50.3, 67.0, 80.3, 115.6 (t, <sup>1</sup>J<sub>C-F</sub> = 250.8 Hz, CF<sub>2</sub>H), 128.2, 128.3, 128.7, 136.6, 157.0.

Anal. calcd for C<sub>13</sub>H<sub>17</sub>F<sub>2</sub>NO<sub>3</sub>: C, 57.14; H, 6.27; N, 5.13. Found: C, 56.98; H, 6.30; N, 5.20.

### 2-(Difluoromethoxy)-2-methylpropan-1-amine (3e).

Compound **2e** (7.4 g, 0.027 mol, 1 eq) was dissolved in 100 mL of methanol and 1 g of 5%-Pd/C was added thereto. The mixture was stirred under 30 atm of H<sub>2</sub> for 48 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated under atmospheric pressure. The product **3e** was purified by distillation under vacuum (18 Torr, 32 °C). Yield = 25.2% (0.95 g).

The product is a transparent colorless liquid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ = 1.33 (s, 6H), 1.37 (s, 2H), 2.73 (s, 2H), 6.35 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 75.9 Hz, CF<sub>2</sub>H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -76.9 (d, 2F, <sup>2</sup>J<sub>H-F</sub> = 77.1 Hz).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 23.6, 51.6, 80.3, 115.1 (t, <sup>1</sup>J<sub>C-F</sub> = 250.0 Hz, CF<sub>2</sub>H).

Anal. calcd for C<sub>5</sub>H<sub>11</sub>F<sub>2</sub>NO: C, 43.16; H, 7.97; N, 10.07. Found: C, 43.08; H, 8.02; N, 10.01.

### Benzyl *trans*-2-(Difluoromethoxy)cyclopentylcarbamate (2f).

Compound **1f** (41.6 g, 0.177 mol, 1 eq) was dissolved in 400 mL of acetonitrile and copper iodide (6.7 g, 0.035 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (47.2 g, 0.265 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solid was filtered out. The resulting solution of EtOAc was evaporated and the crude of **2f** was purified by column chromatography (Hexane : EtOAc - 2 : 1, R<sub>f</sub> = 0.6). Yield = 67% (33.6g).

The product is a transparent colorless thick oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ = 1.49-2.03 (m, 6H), 3.97 (br. s, 1H), 4.39 (br. s, 1H), 5.10 (br. s, 3H), 6.32 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 75.2 Hz, CF<sub>2</sub>H), 7.34 (br. s, 5H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -81.7 (m, 2F).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 29.3, 30.1, 53.1, 57.3, 66.3, 80.1, 116.0 (t, <sup>1</sup>J<sub>C-F</sub> = 255.9 Hz, CF<sub>2</sub>H), 127.7, 127.8, 128.1, 136.0, 155.6.

Anal. calcd for C<sub>14</sub>H<sub>17</sub>F<sub>2</sub>NO<sub>3</sub>: C, 58.94; H, 6.01; N, 4.91. Found: C, 58.81; H, 6.05; N, 4.82.

### *trans*-2-(Difluoromethoxy)cyclopentanamine (3f).

1 Compound **2f** (33.6 g, 0.1177 mol, 1 eq) was dissolved in 300 mL of methanol and 4 g of 5% -Pd/C was added thereto. The mixture  
2 was stirred under 30 atm of H<sub>2</sub> for 36 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated  
3 under vacuum. The product **3f** was purified by distillation under vacuum (18 Torr, 67 °C). Yield = 71% (12.6 g).

4 The product is a transparent colorless liquid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ = 1.21-1.27 (m, 3H), 1.60-1.66 (m, 3H), 1.88-1.98 (m,  
5 2H), 3.16-3.20 (m, 1H), 4.01-4.03 (m, 1H), 6.16 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 76.3 Hz, CF<sub>2</sub>H).

6 <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -81.5 (dd, F, <sup>2</sup>J<sub>F-F</sub> = 164.6 Hz, <sup>2</sup>J<sub>H-F</sub> = 75.4 Hz), -80.9 (dd, F, <sup>2</sup>J<sub>F-F</sub> = 160.4 Hz, <sup>2</sup>J<sub>H-F</sub> = 76.0 Hz).

7 <sup>13</sup>C{<sup>1</sup>H}NMR (125 MHz, CDCl<sub>3</sub>): δ = 19.8, 30.0, 31.3, 57.5, 83.5, 115.8 (t, <sup>1</sup>J<sub>C-F</sub> = 257.7 Hz, CF<sub>2</sub>H).

8 Anal. calcd for C<sub>6</sub>H<sub>11</sub>F<sub>2</sub>NO: C, 47.68; H, 7.34; N, 9.27. Found: C, 47.80; H, 7.38; N, 9.30.

### 16 **Benzyl (3-(difluoromethoxy)cyclopentyl)carbamate (2g) (Mixture of diastereomers).**

17 Compound **1g** (35.7 g, 0.152 mol, 1 eq) was dissolved in 350 mL of acetonitrile and copper iodide (5.8 g, 0.03 mol, 0.2 eq) was  
18 added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (40.5 g, 0.227 mol, 1.5  
19 eq) in 100 mL of acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30  
20 minutes at 50 °C. Next, acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solid was filtered out. The  
21 resulting solution of EtOAc was evaporated and the crude of **2g** was purified by column chromatography (Hexane : EtOAc - 2 : 1,  
22 R<sub>f</sub> = 0.6). Yield = 83% (36 g).

23 The product is a transparent colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) (mixture of diastereomers): δ = 1.66-1.93 (m, 2H), 1.96-2.21  
24 (m, 4H), 4.20 (br. s, 1H), 4.68 (m, 1H), 5.05 (br. s, 1H), 5.10 (s, 2H), 6.18 (dt, 1H, <sup>2</sup>J<sub>H-F</sub> = 74.6 Hz, J = 6.7 Hz, CF<sub>2</sub>H), 7.33-7.36 (m,  
25 5H).

26 <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -82.4 - -81.9 (m, 2F).

27 <sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>) (mixture of diastereomers): δ = 31.1, 31.5, 40.1, 50.5, 53.0, 60.0, 66.3, 115.6 (t, <sup>1</sup>J<sub>C-F</sub> = 259.6 Hz,  
28 CF<sub>2</sub>H), 127.8, 128.2, 136.1, 144.5, 155.3.

29 Anal. calcd for C<sub>14</sub>H<sub>17</sub>F<sub>2</sub>NO<sub>3</sub>: C, 58.94; H, 6.01; N, 4.91. Found: C, 58.81; H, 6.06; N, 4.87.

### 34 **3-(Difluoromethoxy)cyclopentanamine (3g) (Mixture of diastereomers).**

35 Compound **2g** (36 g, 0.1261 mol, 1 eq) was dissolved in 300 mL of methanol and 4 g of 5% -Pd/C was added thereto. The mixture  
36 was stirred under 30 atm of H<sub>2</sub> for 36 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated  
37 under vacuum. Finally, the product **3g** was purified by distillation under vacuum (18 Torr, 65 °C). Yield = 66.6% (12.7 g).

38 The product is a transparent colorless oil. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) (mixture of diastereomers): δ = 1.24 (s, 2H), 1.42-1.69 (m,  
39 2H), 1.75-1.87 (m, 2H), 1.91-2.15 (m, 2H), 3.24, 3.48 (quintet, 1H, <sup>3</sup>J = 6.0 Hz), 4.50-64 (m, 1H), 6.10 (dt, 1H, <sup>2</sup>J<sub>H-F</sub> = 75.7 Hz, J =  
40 13.7 Hz, CF<sub>2</sub>H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -82.2- -81.7 (m, 2F).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>) (mixture of diastereomers): δ = 31.2, 31.5, 33.2, 33.8, 42.7, 42.8, 50.5, 51.1, 76.1, 76.2, 115.8 (t,

<sup>1</sup>J<sub>C-F</sub> = 258.1 Hz, J = 12.8 Hz).

Anal. calcd for C<sub>6</sub>H<sub>11</sub>F<sub>2</sub>NO: C, 47.68; H, 7.34; N, 9.27. Found: C, 47.81; H, 7.40; N, 9.23.

### Benzyl (*trans*-2-(difluoromethoxy)cyclohexyl)carbamate (**2h**).

Compound **1h** (39 g, 0.156 mol, 1 eq) was dissolved in 350 mL of acetonitrile and copper iodide (6 g, 0.032 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (41.8 g, 0.235 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solid was filtered out. The resulting solution of EtOAc was evaporated and the crude **2h** was purified by column chromatography (Hexane : EtOAc - 2 : 1, R<sub>f</sub> = 0.6). Yield = 78.6% (36.8 g).

The product is a transparent colorless thick oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ = 1.28-1.75 (m, 6H), 2.04-2.08 (m, 2H), 3.56 (br. s, 1H), 3.92 (m, 1H), 4.97 (br. s, 1H), 5.11 (s, 2H), 6.22 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 74.6 Hz, CF<sub>2</sub>H), 7.30-7.34 (m, 5H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -81.0 (m, 2F).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 23.5, 31.2, 31.9, 53.3, 60.0, 66.2, 75.6, 115.6 (t, <sup>1</sup>J<sub>C-F</sub> = 257.3 Hz, CF<sub>2</sub>H), 127.6, 128.1, 136.3, 155.7.

Anal. calcd for C<sub>15</sub>H<sub>19</sub>F<sub>2</sub>NO<sub>3</sub>: C, 60.19; H, 6.40; N, 4.68. Found: C, 60.23; H, 6.35; N, 4.60.

### *trans*-2-(Difluoromethoxy)cyclohexanamine (**3h**).

Compound **2h** (30 g, 0.1 mol, 1 eq) was dissolved in 300 mL of methanol and 4 g of 5% -Pd/C was added thereto. The mixture was stirred under 30 atm of H<sub>2</sub> for 36 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated under vacuum. The product **3h** was purified by destination under vacuum (18 Torr, 75 °C). Yield = 72.5% (12 g).

The product is a transparent colorless liquid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ = 0.96-1.14 (m, 4H), 1.19-1.25 (m, 4H), 1.49-1.57 (m, 2H), 1.73-1.87 (m, 2H), 2.47-2.53 (m, 1H), 3.43-3.48 (m, 1H), 6.14 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 75.0 Hz, CF<sub>2</sub>H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -80.7 (dd, F, <sup>2</sup>J<sub>F-F</sub> = 161.0 Hz, <sup>2</sup>J<sub>H-F</sub> = 70.0 Hz), -79.0 (dd, 2F, <sup>2</sup>J<sub>F-F</sub> = 155.0 Hz, <sup>2</sup>J<sub>H-F</sub> = 75.0 Hz).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 23.7, 23.8, 31.3, 33.0, 53.6, 81.5, 116.1 (t, <sup>1</sup>J<sub>C-F</sub> = 258.9 Hz, CF<sub>2</sub>H).

Anal. calcd for C<sub>7</sub>H<sub>13</sub>F<sub>2</sub>NO: C, 50.90; H, 7.93; N, 8.48. Found: C, 50.79; H, 7.96; N, 8.53.

### Benzyl (3-(difluoromethoxy)cyclohexyl)carbamate (**2i**) (Mixture of diastereomers).

Compound **1i** (49.8 g, 0.2 mol, 1 eq) was dissolved in 450 mL of acetonitrile and copper iodide (7.6 g, 0.04 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (53.4g, 0.3 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solids were filtered out. The resulting solution of EtOAc was evaporated and the crude of **2i** was purified by column chromatography (Hexane : EtOAc - 2 : 1,  $R_f$  = 0.6). Yield = 90.3% (54 g).

The product is a transparent colorless oil.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta$  = 1.16-2.26 (m, 6H), 3.62, 3.90 (br. s, 1H), 4.11, 4.48 (m, 1H), 5.01-5.18 (m, 3H), 5.26 (s, 2H), 6.20 (dt, 1H,  $^2J_{\text{H-F}} = 75.3$  Hz,  $J = 5.4$  Hz,  $\text{CF}_2\text{H}$ ), 7.31-7.34 (m, 5H).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta$  = -81.8 (d, F,  $^2J = 74.0$  Hz), -81.0 (d, F,  $^2J = 74.0$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta$  = 18.9, 20.2, 30.3, 31.5, 37.3, 38.7, 45.5, 47.8, 53.1, 66.1, 70.5, 72.3, 115.9 (dt,  $^1J_{\text{C-F}} = 258.1$  Hz,  $J = 10.6$  Hz,  $\text{CF}_2\text{H}$ ), 127.7, 128.1, 136.3, 155.2.

Anal. calcd for  $\text{C}_{15}\text{H}_{19}\text{F}_2\text{NO}_3$ : C, 60.09; H, 6.40; N, 4.68. Found: C, 60.22; H, 6.42; N, 4.76.

### **3-(Difluoromethoxy)cyclohexanamine (3i) (Mixture of diastereomers).**

Compound **2i** (30 g, 0.1 mol, 1 eq) was dissolved in 300 mL of methanol and 4 g of 5%-Pd/C was added thereto. The mixture was stirred under 30 atm of  $\text{H}_2$  for 36 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated under vacuum. The product **3i** was purified by destination under vacuum (18 Torr, 77 °C). Yield = 74.3% (12.3 g).

The product is a transparent colorless liquid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta$  = 0.97-1.46 (m, 4H), 1.52 (s, 2H), 1.49-1.57 (m, 2H), 1.62-2.18 (m, 2H), 2.68-3.11 (m, 1H), 4.02-4.49 (m, 1H), 6.19 (dt, 1H,  $^2J_{\text{H-F}} = 76.0$  Hz,  $J = 4.6$  Hz,  $\text{CF}_2\text{H}$ ).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta$  = -81.4 - -81.0 (m, 2F).

$^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta$  = 19.2, 21.7, 30.8, 32.2, 35.1, 35.3, 40.8, 43.2, 45.4, 49.0, 71.6, 73.0, 116.1, 116.4 (t,  $^1J_{\text{H-F}} = 253.4$  Hz,  $\text{CF}_2\text{H}$ ).

Anal. calcd for  $\text{C}_{15}\text{H}_{19}\text{F}_2\text{NO}_3$ : C, 60.19; H, 6.40; F, 12.69; N, 4.68. Found: C, 60.15; H, 6.42; F, 12.65; N, 4.65.

### **Benzyl ((4-hydroxytetrahydro-2H-pyran-4-yl)methyl)carbamate (1j).**

Benzyl chloroformate (CbzCl) (43 g, 0.252 mol, 1 eq) was added dropwise to the stirring solution of aminoalcohol (31.5 g, 0.516 mol, 2 eq) and potassium carbonate (66.4 g, 0.481 mol, 1.9 eq) in the mixture THF/water (400/200 mL) at 0 °C. Afterwards, the reaction mixture was stirred for additional 18 h at room temperature. Upon completion the THF was evaporated and the rest was

1  
2 extracted 3 times with DCM. The organic layer was dried with sodium sulphate, filtered and evaporated to dryness. The crude  
3 product **1j** was purified by flash-chromatography (EtOAc : Petroleum ether - 1 : 3). Yield = 75.1% (47.85 g).

4 The product is a white solid, mp = 143-146 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ = 1.48-1.51 (m, 2H), 1.59-1.64 (m, 2H), 2.72 (s,  
5 1H), 3.21 (br. s, 2H), 3.72 (br. s, 4H), 5.09 (s, 2H), 5.41 (br. s, 1H), 7.34 (s, 5H).

6  
7 <sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 35.5, 51.3, 63.7, 67.2, 69.2, 128.2, 128.4, 128.7, 136.4, 157.7.

8  
9 Anal. calcd for C<sub>14</sub>H<sub>19</sub>NO<sub>4</sub>: C, 63.38; H, 7.22; N, 5.28. Found: C, 63.22; H, 7.27; N, 5.25.

#### 10 11 12 13 **Benzyl ((4-(difluoromethoxy)tetrahydro-2H-pyran-4-yl)methyl)carbamate (2j).**

14  
15 Compound **1j** (47.85 g, 0.18 mol, 1 eq) was dissolved in 450 mL of acetonitrile and copper iodide (6.9 g, 0.036 mol, 0.2 eq) was  
16 added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (48.3 g, 0.271 mol, 1.5  
17 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30  
18 minutes at 50 °C. Afterwards, acetonitrile was evaporated, the rest was dissolved in EtOAc and the solid phase was filtered out. The  
19 resulting solution of EtOAc was evaporated and the crude of **2j** was purified by column chromatography (EtOAc : Hexane - 3 : 1,  
20 R<sub>f</sub> = 0.65). Yield = 12.8% (7.3 g).

21  
22 The product is a transparent colorless thick oil. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ = 1.70-1.83 (m, 4H), 3.44-3.46 (m, 2H), 3.74 (br. s,  
23 4H), 5.11 (s, 2H), 5.23 (br. s, 1H), 6.36 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 76.0 Hz, CF<sub>2</sub>H), 7.35 (s, 5H).

24  
25 <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -77.6 (d, 2F, <sup>2</sup>J<sub>H-F</sub> = 77.7 Hz).

26  
27 <sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 33.3, 47.6, 63.3, 67.3, 78.8, 114.7 (t, <sup>1</sup>J<sub>C-F</sub> = 264.6 Hz, CF<sub>2</sub>H), 127.7, 127.8, 128.2, 135.8,  
28 156.4.

29  
30 Anal. calcd for C<sub>15</sub>H<sub>19</sub>F<sub>2</sub>NO<sub>4</sub>: C, 57.14; H, 6.07; N, 4.44. Found: C, 57.01; H, 6.13; N, 4.41.

#### 31 32 33 34 **(4-(Difluoromethoxy)tetrahydro-2H-pyran-4-yl)methanamine (3j).**

35  
36 Compound **2j** (7.3 g, 0.232 mol, 1 eq) was dissolved in 100 mL of methanol and 1 g of 5%-Pd/C was added thereto. The mixture  
37 was stirred under 30 atm of H<sub>2</sub> for 48 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated  
38 under vacuum. The product **3j** was purified by distillation under vacuum (1 Torr, 44 °C). Yield = 42.9% (1.8 g).

39  
40 The product is a transparent colorless liquid. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 1.24 (m, 2H), 1.64-1.86 (m, 4H), 2.87 (s, 2H), 3.74-  
41 3.76(m, 4H), 6.54 (t, 1H, <sup>2</sup>J = 77.8 Hz, CF<sub>2</sub>H).

42  
43 <sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -77.2 (d, 2F, <sup>2</sup>J<sub>H-F</sub> = 84.7 Hz).

44  
45 <sup>13</sup>C{<sup>1</sup>H}NMR (100 MHz, CDCl<sub>3</sub>): δ = 33.2, 49.3, 63.1, 79.4, 115.2 (t, <sup>1</sup>J<sub>C-F</sub> = 231.2 Hz, CF<sub>2</sub>H).

46  
47 Anal. calcd for C<sub>7</sub>H<sub>13</sub>F<sub>2</sub>NO<sub>2</sub>: C, 46.40; H, 7.23; N, 7.73. Found: C, 46.53; H, 7.27; N, 7.68.

**Benzyl 3-((difluoromethoxy)methyl)pyrrolidine-1-carboxylate (2k).**

Compound **1k** (40g, 0.17 mol, 1 eq) was dissolved in 400 mL of acetonitrile and copper iodide (6.5 g, 0.034 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (45.4g, 0.2549 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Next, acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solid was filtered out. The resulting solution of EtOAc was evaporated and the crude of **2k** was purified by column chromatography (Hexane : EtOAc - 7 : 1,  $R_f = 0.6$ ). Yield = 80% (39 g).

The product is a transparent thick oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.67$  (m, 1H), 1.99 (br. s, 1H), 2.48 (m, 1H,  $\text{CH}_{\text{pyrrolidine}}$ ), 3.14-3.37 (m, 2H), 3.49-3.81 (m, 4H), 5.08 (s, 2H), 6.16 (t, 1H,  $^2J_{\text{H-F}} = 74.7$  Hz,  $\text{CF}_2\text{H}$ ), 7.28-7.33 (m, 5H).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -85.0$  (d, F,  $^2J_{\text{H-F}} = 74.2$  Hz),  $-84.9$  (d, F,  $^2J_{\text{H-F}} = 74.2$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta = 27.5, 28.3, 37.6, 38.5, 45.2, 45.6, 48.5, 48.9, 63.3, 66.9, 115.9$  (t,  $^1J_{\text{C-F}} = 262.3$  Hz,  $\text{CF}_2\text{H}$ ), 128.0, 128.1, 128.6, 137.1, 154.9.

Anal. calcd for  $\text{C}_{14}\text{H}_{17}\text{F}_2\text{NO}_3$ : C, 58.99; H, 6.01; N, 4.91. Found: C, 58.93; H, 6.07; N, 4.83.

**3-((Difluoromethoxy)methyl)pyrrolidine (3k).**

Compound **2k** (30 g, 0.105 mol, 1 eq) was dissolved in 150 mL of methanol and 3 g of 5% -Pd/C was added thereto. The mixture was stirred under 30 atm of  $\text{H}_2$  for 36 h at room temperature. After completion of the reaction Pd/C was filtered out and the reaction mixture was concentrated under vacuum. The product **3k** was purified by distillation under vacuum (18 Torr, 57 °C). Yield = 68.6% (10.9 g).

The product is a transparent colorless liquid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.15$ -1.20 (m, 1H), 1.61-1.70 (m, 1H), 1.75 (s, 1H), 2.11-2.18 (m, 1H), 2.41-2.45 (m, 1H), 2.59-2.80 (m, 3H), 3.46-3.56 (m, 2H), 5.95 (t, 1H,  $^2J_{\text{H-F}} = 75.1$  Hz,  $\text{CF}_2\text{H}$ ).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -84.5$  (d, 2F,  $^2J_{\text{H-F}} = 77.0$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125 MHz,  $\text{CDCl}_3$ ):  $\delta = 28.4, 37.8, 46.2, 49.5, 65.5$  (t,  $J = 4.7$  Hz), 115.4 (t,  $^1J_{\text{C-F}} = 258.8$  Hz,  $\text{CF}_2\text{H}$ ).

Anal. calcd for  $\text{C}_6\text{H}_{11}\text{F}_2\text{NO}$ : C, 47.68; H, 7.34; N, 9.27. Found: C, 47.81; H, 7.29; N, 9.21.

**(S)-Benzyl 3-((difluoromethoxy)pyrrolidine-1-carboxylate (2l).**

Compound **1l** (50g, 0.226 mol, 1 eq) was dissolved in 400 mL of acetonitrile and copper iodide (8.6 g, 0.045 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (52.3 g, 0.294 mol, 1.3 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. As next, acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solids were filtered out. The re-

sulting solution of EtOAc was evaporated and the crude of **2l** was purified by column chromatography (Hexane : EtOAc - 7 : 1,  $R_f$  = 0.6). Yield = 84.8% (52 g).

The product is a transparent colorless oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , Ethyl acetate inside):  $\delta$  = 2.06 (br. s, 2H), 3.50-3.58 (m, 4H), 4.79 (m, 1H), 5.11 (s, 2H), 6.20 (t, 1H,  $^2J_{\text{H-F}} = 74.8$  Hz,  $\text{CF}_2\text{H}$ ), 7.28-7.34 (m, 5H).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta$  = -83.1 (d, 2F,  $^2J_{\text{H-F}} = 72.5$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 31.6, 32.4, 43.6, 44.0, 52.0, 52.3, 66.9, 72.5, 73.1, 115.8 (t,  $^1J_{\text{C-F}} = 273.0$  Hz,  $\text{CF}_2\text{H}$ ), 127.9, 128.0, 128.5, 136.8, 154.8.

Anal. calcd for  $\text{C}_{13}\text{H}_{15}\text{F}_2\text{NO}_3$ : C, 57.56; H, 5.57; N, 5.16. Found: C, 57.70; H, 5.61; N, 5.11.

### (S)-3-(Bifluoromethoxy)pyrrolidine (**3l**).

Compound **2l** (52 g, 0.192 mol, 1 eq) was dissolved in 300 mL of methanol and 5 g of 5% -Pd/C was added thereto. The mixture was stirred under 30 atm of  $\text{H}_2$  for 36 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated under vacuum. The product **3l** was purified by distillation under vacuum (18 Torr, 51 °C). Yield = 55.2% (14.5 g).

The product is a transparent colorless liquid,  $[\alpha]_{\text{D}} = -9.03$  (EtOH,  $c = 72.925$  mmol/L).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 1.87-2.00 (m, 3H), 2.81-3.11 (m, 4H), 4.71 (br. s, 1H), 6.17 (t, 1H,  $^2J_{\text{H-F}} = 74.6$  Hz,  $\text{CF}_2\text{H}$ ).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta$  = -82.1 (d, 2F,  $^2J_{\text{H-F}} = 74.1$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 33.6, 45.5, 54.0, 75.8, 116.0 (t,  $^1J_{\text{C-F}} = 258.4$  Hz,  $\text{CF}_2\text{H}$ ).

Anal. calcd for  $\text{C}_5\text{H}_9\text{F}_2\text{NO}$ : C, 43.79; H, 6.62; N, 10.21. Found: C, 43.65; H, 6.65; N, 10.15.

### (R)-Benzyl 3-(difluoromethoxy)pyrrolidine-1-carboxylate (**2m**).

Compound **1m** (40 g, 0.181 mol, 1 eq) was dissolved in 400 mL of acetonitrile and copper iodide (6.9 g, 0.036 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (41.9 g, 0.235 mol, 1.3 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solids were filtered out. The resulting solution of EtOAc was evaporated and the crude of **2m** was purified by column chromatography (Hexane : EtOAc - 7 : 1,  $R_f$  = 0.6). Yield = 82.6% (40.5 g).

The product is a transparent colorless oil.  $^1\text{H}$  NMR (500 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  = 2.08 (br. s, 2H), 3.55-3.57 (m, 4H), 4.79 (m, 1H), 5.12 (s, 2H), 6.21 (t, 1H,  $^2J_{\text{H-F}} = 74.4$  Hz,  $\text{CF}_2\text{H}$ ), 7.29-7.35 (m, 5H).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta$  = -83.0 (d, 2F,  $^2J_{\text{H-F}} = 74.5$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125 MHz,  $\text{CDCl}_3$ ):  $\delta$  = 31.2, 31.9, 43.2, 43.5, 51.5, 51.9, 66.5, 72.2, 72.9, 115.5 (t,  $^1J_{\text{C-F}} = 261.9$  Hz,  $\text{CF}_2\text{H}$ ), 127.5, 128.1, 136.4, 155.3.

Anal. calcd for  $C_{13}H_{15}F_2NO_3$ : C, 57.56; H, 5.57; N, 5.16. Found: C, 57.41; H, 5.52; N, 5.13.

**(R)-3-(Difluoromethoxy)pyrrolidine (3m).**

Compound **2m** (40.5 g, 0.149 mol, 1 eq) was dissolved in 300 mL of methanol and 5 g of 5% -Pd/C was added thereto. The mixture was stirred under 30 atm of  $H_2$  for 36 h at room temperature. Then Pd/C was filtered out and the mixture was concentrated under vacuum. The product **3m** was purified by distillation under vacuum (18 Torr, 51 °C). Yield = 63.5% (13 g).

The product is a transparent colorless liquid,  $[\alpha]_D = +6.31$  (EtOH,  $c = 72.925$  mmol/L).  $^1H$  NMR (500 MHz,  $CDCl_3$ ):  $\delta = 1.95$ -2.05 (m, 2H), 2.92-3.14 (m, 3H), 3.43 (s, 2H), 4.75 (s, 1H), 6.20 (t, 1H,  $^2J_{H-F} = 74.4$  Hz).

$^{19}F$  NMR (376 MHz,  $CDCl_3$ ):  $\delta = -82.3$  (d, 2F,  $^2J_{H-F} = 74.5$  Hz).

$^{13}C\{^1H\}$  NMR (125 MHz,  $CDCl_3$ ):  $\delta = 33.3, 45.1, 53.5, 75.4, 116.0$  (t,  $^1J_{C-F} = 295.8$  Hz,  $CF_2H$ ).

Anal. calcd for  $C_5H_9F_2NO$ : C, 43.79; H, 6.62; N, 10.21. Found: C, 43.72; H, 6.70; N, 10.15.

**Benzyl 3-(difluoromethoxy)azetidine-1-carboxylate (2n).**

Compound **1n** (44.7g, 0.218 mol, 1 eq) was dissolved in 400 mL of acetonitrile and copper iodide (8.2g, 0.043 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (57.65 g, 0.324 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solid was filtered out. Afterwards, the resulting solution of EtOAc was evaporated and the crude of **2n** was purified by column chromatography (Hexane : EtOAc - 7 : 1,  $R_f = 0.6$ ). Yield = 86.5% (48 g).

The product is a transparent yellowish oil.  $^1H$  NMR (400 MHz,  $CDCl_3$ ):  $\delta = 4.06$ -4.09 (m, 2H), 4.26-4.31 (m, 2H), 4.93-4.94 (m, 1H), 5.11 (s, 2H), 6.24 (t, 1H,  $^2J_{H-F} = 73.0$  Hz,  $CF_2H$ ), 7.36 (br. s, 5H).

$^{19}F$  NMR (376 MHz,  $CDCl_3$ ):  $\delta = -84.9$  (d, 2F,  $^2J_{H-F} = 74.9$  Hz).

$^{13}C\{^1H\}$  NMR (125 MHz,  $CDCl_3$ ):  $\delta = 56.8, 62.1$  (t,  $66.3$  J = 6.5 Hz), 66.8, 115.3 (t,  $^1J_{C-F} = 260.6$  Hz,  $CF_2H$ ), 127.9, 128.0, 128.4, 136.3, 156.1.

Anal. calcd for  $C_{12}H_{13}F_2NO_3$ : C, 56.03; H, 5.09; N, 5.45. Found: C, 55.89; H, 5.16; N, 5.39.

**3-(Difluoromethoxy)azetidine (3n).**

Compound **2n** (48 g, 0.187 mol, 1 eq) was dissolved in 300 mL of methanol and 5 g of 5% -Pd/C was added thereto. The mixture was stirred under 30 atm of  $H_2$  for 36 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated under vacuum. The product **3n** was purified by distillation under vacuum (18 Torr, 47 °C). Yield = 53.5% (12.3 g).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The product is a transparent colorless liquid.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta = 2.08$  (s, 1H), 3.70-3.71 (m, 4H), 4.91 (quintet, 1H,  $^3J = 7.0$  Hz), 6.21 (t, 1H,  $^2J_{\text{H-F}} = 73.9$  Hz,  $\text{CF}_2\text{H}$ ).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -84.1$  (d, 2F,  $^2J_{\text{H-F}} = 74.4$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (125 MHz,  $\text{CDCl}_3$ ):  $\delta = 54.3, 66.3, 115.4$  (t,  $^1J_{\text{C-F}} = 261.1$  Hz,  $\text{CF}_2\text{H}$ ).

Anal. calcd for  $\text{C}_4\text{H}_7\text{F}_2\text{NO}$ : C, 39.03; H, 5.73; N, 11.38. Found: C, 39.11; H, 5.79; N, 11.34.

### Benzyl 2-(difluoromethoxy)acetate (2o).

Compound **1o** (50.15 g, 0.302 mol, 1 eq) was dissolved in 500 mL of acetonitrile and copper iodide (11.5 g, 0.06 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (80.65 g, 0.453 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the solid phase was filtered out. The resulting solution of EtOAc was evaporated and the crude of **2o** was purified by column chromatography (Hexane : EtOAc - 1 : 7,  $R_f = 0.7$ ). Yield = 81.2% (53g).

The product is a transparent colorless liquid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 4.44$  (s, 2H), 5.21 (s, 2H), 6.34 (t, 1H,  $^2J_{\text{H-F}} = 73.9$  Hz,  $\text{CF}_2\text{H}$ ), 7.36 (br. s, 5H).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -86.6$  (d, 2F,  $^2J_{\text{H-F}} = 73.6$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta = 59.9, 67.3, 115.3$  (t,  $^1J_{\text{C-F}} = 272.0$  Hz,  $\text{CF}_2\text{H}$ ), 128.5, 128.8, 134.9, 167.6.

Anal. calcd for  $\text{C}_{10}\text{H}_{10}\text{F}_2\text{O}_3$ : C, 55.56; H, 4.66; F, 17.58. Found: C, 55.67; H, 4.71; F, 17.52.

### 2-(Difluoromethoxy)acetic acid (3o).

Compound **2o** (53 g, 0.245 mol, 1 eq) was dissolved in 300 mL of methanol and 7 g of 5%-Pd/C was added thereto. The mixture was stirred under 30 atm of  $\text{H}_2$  for 72 h at room temperature. After completion of the reaction Pd/C was filtered out and the reaction mixture was concentrated under vacuum. The product **3o** was dissolved in conc. solution of sodium bicarbonate and the obtained solution was washed three times with EtOAc. Then the solution was carefully acidified with conc. solution of citric acid and extracted three times with EtOAc. Combined organic layer was washed twice with brine, dried over magnesium sulphate and concentrated under vacuum to give the product **3o**. Yield = 64.1% (19.8 g).

The product is a white solid, mp = 46-48 °C.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 4.51$  (s, 2H), 6.36 (t, 1H,  $^2J_{\text{H-F}} = 75.9$  Hz,  $\text{CF}_2\text{H}$ ), 10.62 (br. s, 1H).

$^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -86.8$  (d, 2F,  $^2J_{\text{H-F}} = 87.9$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta = 59.1, 115.2$  (t,  $^1J_{\text{C-F}} = 285.6$  Hz,  $\text{CF}_2\text{H}$ ), 173.7.

Anal. calcd for  $\text{C}_3\text{H}_4\text{F}_2\text{O}_3$ : C, 28.58; H, 3.20. Found: C, 28.52; H, 3.10.

**1,4-Benzyl 4-(difluoromethoxy)cyclohexanecarboxylate (2p) (Mixture of diastereomers).**

Compound **1p** (as a mixture of *cis*- and *trans*- isomers 7:3) (42 g, 0.179 mol, 1 eq) was dissolved in 500 mL of acetonitrile and copper iodide (6.85 g, 0.36 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (47.9 g, 0.269 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solids were filtered out. The resulting solution of EtOAc was evaporated and the crude of **2p** was purified by column chromatography (Hexane : EtOAc - 1 : 7,  $R_f = 0.6$ ). Yield = 73.4% (37.4 g).

The product is a transparent colorless thick oil.  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.47\text{-}2.42$  (m, 8H), 4.06, 4.33 (s, 1H), 5.10 (br. s, 1H), 5.26 (s, 2H), 6.21 (t, 1H,  $^2J_{\text{H-F}} = 73.9$  Hz,  $\text{CF}_2\text{H}$ ), 7.34 (br. s, 5H).

$^{19}\text{F NMR}$  (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -83.6$  (d, 2F,  $^2J_{\text{H-F}} = 73.7$  Hz),  $-83.1$  (d, 2F,  $^2J_{\text{H-F}} = 73.7$  Hz).

$^{13}\text{C NMR}\{^1\text{H}\}$  (100 MHz,  $\text{DMSO-}d_6$ ) (mixture of diastereomers):  $\delta = 14.0, 22.1, 23.2, 26.3, 29.3, 31.3, 40.8, 65.5, 74.3, 117.2$  (t,

$^1J_{\text{C-F}} = 253.0$  Hz,  $\text{CF}_2\text{H}$ ), 117.4 (t,  $^1J_{\text{C-F}} = 253.0$  Hz,  $\text{CF}_2\text{H}$ ), 127.8, 128.0, 128.5, 136.3, 136.4, 174.2, 174.2.

Anal. calcd for  $\text{C}_{15}\text{H}_{18}\text{F}_2\text{O}_3$ : C, 63.37; H, 6.38. Found: C, 63.22; H, 6.43.

**1,4-*cis*-(difluoromethoxy)cyclohexanecarboxylic acid (3p).**

Compound **2p** (30 g, 0.106 mol, 1 eq) was dissolved in 300 mL of methanol and 6.5 g of 5%-Pd/C was added thereto. The mixture was stirred under 30 atm of  $\text{H}_2$  for 72 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated under vacuum. The crude was dissolved in conc. solution of sodium bicarbonate and the obtained solution was washed three times with EtOAc. Then the solution was carefully acidified with conc. solution of citric acid and extracted three times with EtOAc. Combined organic layer was washed twice with brine, dried over magnesium sulphate and concentrated under vacuum and recrystallized from petroleum ether – ethyl acetate (10:1) to give the product **3p**. Yield = 60% (12.3 g). Compound **3p** was obtained as a pure *cis*-isomer.

The product is a white solid, mp = 117-119 °C.  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.45\text{-}1.61$  (m, 4H), 2.07-2.10 (m, 4H), 2.32-2.37 (m, 1H), 4.09 (br. s, 1H), 6.23 (t, 1H,  $^2J_{\text{H-F}} = 74.6$  Hz,  $\text{CF}_2\text{H}$ ), 11.0 (br. s, 1H).

$^{19}\text{F NMR}$  (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -81.2$  (d, 2F,  $^2J_{\text{H-F}} = 76.4$  Hz).

$^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta = 28.3, 31.6, 41.4, 73.0, 116.2$  (t,  $^1J_{\text{C-F}} = 258.4$  Hz,  $\text{CF}_2\text{H}$ ), 181.6.

Anal. calcd for  $\text{C}_8\text{H}_{12}\text{F}_2\text{O}_3$ : C, 49.48; H, 6.23; F, 19.57. Found: C, 49.33; H, 6.19.

**Benzyl 3-(difluoromethoxy)cyclobutanecarboxylate (2q) (Mixture of diastereomers).**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
Compound **1q** (12.2 g, 0.059 mol, 1 eq) was dissolved in 200 mL of acetonitrile and copper iodide (2.25 g, 0.012 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (15.8 g, 0.089 mol, 1.5 eq) in 100 mL acetonitrile was added dropwise over a period of 45 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solid was filtered out. The resulting solution of EtOAc was evaporated and the crude of **2q** was purified by column chromatography (Hexane : EtOAc - 1 : 7,  $R_f = 0.6$ ). Yield = 73.9% (11.2g).

11  
12  
13  
14  
The product is a transparent colorless oil.  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta = 2.40\text{-}2.47$  (m, 2H),  $2.57\text{-}2.59$  (m, 2H),  $2.71\text{-}2.75$  (m, 1H),  $4.51\text{-}4.57$  (m, 1H),  $5.11$  (s, 2H),  $6.13$  (t, 1H,  $^2J_{\text{H-F}} = 74.0$  Hz,  $\text{CF}_2\text{H}$ ),  $7.34$  (br. s, 5H).

15  
16  
 $^{19}\text{F NMR}$  (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -83.6$  (d, 2F,  $^2J_{\text{H-F}} = 72.0$  Hz).

17  
18  
 $^{13}\text{C}\{^1\text{H}\}\text{NMR}$  (125.7 MHz,  $\text{CDCl}_3$ ):  $\delta = 29.9, 34.4, 63.9, 66.7, 115.7$  (t,  $^1J_{\text{C-F}} = 259.8$  Hz,  $\text{CF}_2\text{H}$ ),  $128.3, 128.4, 128.7, 136.0, 173.7$ .

19  
20  
Anal. calcd for  $\text{C}_{13}\text{H}_{14}\text{F}_2\text{O}_3$ : C, 60.93; H, 5.51. Found: C, 61.03; H, 5.45.

### 21 22 23 **3-(Difluoromethoxy)cyclobutanecarboxylic acid (3q) (Mixture of diastereomers).**

24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
Compound **2q** (3.5 g, 0.014 mol, 1 eq) was dissolved in 100 mL of methanol and 0.9 g of 5% -Pd/C was added thereto. The mixture was stirred under 30 atm of  $\text{H}_2$  for 36 h at room temperature. Then Pd/C was filtered out and the reaction mixture was concentrated under vacuum. The product **3q** was dissolved in conc. solution of sodium bicarbonate and the obtained solution was washed three times with EtOAc. Then the solution was carefully acidified with conc. solution of citric acid and extracted three times with EtOAc. Combined organic layer was washed twice with brine, dried over magnesium sulphate and concentrated under vacuum to give the product **3q**. Yield = 57.3% (1.3 g).

35  
36  
37  
38  
39  
40  
41  
42  
The product is a transparent colorless oil.  $^1\text{H NMR}$  (400 MHz,  $\text{CDCl}_3$ ) (mixture of diastereomers):  $\delta = 2.43\text{-}2.48$  (m, 2H),  $2.61\text{-}2.63$  (m,  $2\text{H}_{\text{butyl}}$ ),  $2.71\text{-}2.76$  (m, 1H),  $4.55\text{-}4.59$  (m, 1H),  $6.15$  (t, 1H,  $^2J_{\text{H-F}} = 74.0$  Hz,  $\text{CF}_2\text{H}$ ),  $12.0$  (s, 1H).

43  
44  
 $^{19}\text{F NMR}$  (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -83.9$  (d, 2F,  $^2J_{\text{H-F}} = 71.6$  Hz).

45  
46  
 $^{13}\text{C}\{^1\text{H}\}\text{NMR}$  (125 MHz,  $\text{CDCl}_3$ ):  $\delta = 29.3, 31.6, 33.7, 63.1$  (t,  $J = 5.6$  Hz),  $115.1$  (t,  $^1J_{\text{C-F}} = 261.2$  Hz,  $\text{CF}_2\text{H}$ ),  $181.1$ .

47  
48  
Anal. calcd for  $\text{C}_6\text{H}_8\text{F}_2\text{O}_3$ : C, 43.38; H, 4.85. Found: C, 43.49; H, 4.91.

### 49 50 51 52 53 54 **Tert-butyl (S)-3-(methoxycarbonyl)-3-(difluoromethoxy)propylcarbamate (2r).**

55  
56  
57  
58  
59  
60  
Compound **1r** (3.2 g, 0.014 mol, 1 eq) was dissolved in 50 mL of acetonitrile and copper iodide (0.5 g, 0.0026 mol, 0.2 eq) was added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (2.95 g, 0.017 mol, 1.2 eq) in 20 mL acetonitrile was added dropwise over a period of 30 minutes. The reaction mixture was heated for additional 30 minutes at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solids were filtered out. The

1 resulting solution of EtOAc was evaporated and the crude of **2r** was purified by column chromatography (Hexane : EtOAc - 3 : 1,  
2  $R_f = 0.65$ ). Yield = 79.8% (3.1 g).

3  
4 The product is a transparent colorless oil,  $[\alpha]_D = -20.2$  (MeOH,  $c = 35.3$  mmol/L).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.45$  (s, 9H),  
5 1.99-2.08 (m, 2H), 3.27 (br. s, 2H), 3.79 (s, 3H), 4.65-4.68 (m, 2H), 6.36 (t, 1H,  $^2J_{\text{H-F}} = 74.6$  Hz).

6  
7  $^{19}\text{F}$  NMR (376 MHz,  $\text{CDCl}_3$ ):  $\delta = -85.0$  (dd, F,  $^2J_{\text{F-F}} = 160.2$  Hz,  $^2J_{\text{H-F}} = 73.4$  Hz),  $-83.9$  (dd, F,  $^2J_{\text{F-F}} = 158.5$  Hz,  $^2J_{\text{H-F}} = 71.8$  Hz).

8  
9  $^{13}\text{C}\{^1\text{H}\}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta = 27.7, 28.4, 32.4, 52.6, 69.8, 80.0, 115.5$  (t,  $^1J_{\text{C-F}} = 272.0$  Hz,  $\text{CF}_2\text{H}$ ), 155.8, 170.7.

10  
11 Anal. calcd for  $\text{C}_{11}\text{H}_{19}\text{F}_2\text{NO}_5$ : C, 46.64; H, 6.76; N, 4.94. Found: C, 46.78; H, 6.70; N, 5.02.

12  
13  
14  
15 **(2S)-2-(difluoromethoxy)-4-[(1,1-dimethylethoxy)carbonylamino]-butanoic acid (3r).**

16  
17 Compound **2r** (1 g, 0.0035 mol, 1 eq) was dissolved in 20 mL of THF and the solution of  $\text{LiOH}\cdot\text{xH}_2\text{O}$  (0.178 g, 0.0042 mol, 1.2 eq)  
18 in 5 mL of water was added all at once at 0 °C. The reaction mixture was heated to room temperature and was stirred for 4 h. Af-  
19 terwards, the reaction mixture was diluted with 25 mL of EtOAc and 20 mL of water. The organic layer was separated, washed  
20 twice with 20 mL of water. Combined water solution was washed once with 25 mL of EtOAc, carefully acidified with 20 mL of  
21 conc. solution of citric acid and extracted three times with 30 mL of EtOAc. Combined organic layer was washed with brine, dried  
22 with magnesium sulphate and concentrated under vacuum to give compound **3r**. Yield = 57.9% (0.55 g).

23  
24 The product is a white solid, mp = 96-99 °C,  $[\alpha]_D = -21.3$  (MeOH,  $c = 37.14$  mmol/L).  $^1\text{H}$  NMR (500 MHz,  $\text{DMSO}-d_6$ ):  $\delta = 1.37$  (s,  
25 9H), 1.76-1.91 (m, 2H), 3.01 (br. s, 2H), 4.50-4.52 (m, 1H), 6.70 (t, 1H,  $^2J_{\text{H-F}} = 76.0$  Hz,  $\text{CF}_2\text{H}$ ), 6.87 (br. s, 1H), 13.20 (br. s, 1H).

26  
27  $^{19}\text{F}$  NMR (376 MHz,  $\text{DMSO}-d_6$ ):  $\delta = -82.3$  (d, 2F,  $^2J_{\text{H-F}} = 75.5$  Hz).

28  
29  $^{13}\text{C}\{^1\text{H}\}$  NMR (125 MHz,  $\text{DMSO}-d_6$ ):  $\delta = 28.2, 32.0, 36.0, 71.5, 77.7, 116.9$  (t,  $^1J_{\text{C-F}} = 280.5$  Hz,  $\text{CF}_2\text{H}$ ), 155.6, 171.3.

30  
31 Anal. calcd for  $\text{C}_{10}\text{H}_{17}\text{F}_2\text{NO}_5$ : C, 44.61; H, 6.36; N, 5.20. Found: C, 44.72; H, 6.30; N, 5.17.

32  
33  
34  
35 **(2S,4R)-1-tert-butyl 2-methyl 4-(difluoromethoxy)pyrrolidine-1,2-dicarboxylate (2s).**

36  
37 Compound **1s** (7.5 g, 0.031 mol, 1 eq) was dissolved in 100 mL of acetonitrile and copper iodide (1.18 g, 0.0062 mol, 0.2 eq) was  
38 added thereto. The mixture was heated to 50 °C and the solution of 2-fluorosulfonyl-2,2-difluoroacetic acid (6.6 g, 0.037 mol, 1.2  
39 eq) in 30 mL acetonitrile was added dropwise over a period of 30 minutes. The reaction mixture was kept for additional 30 minutes  
40 at 50 °C. Then acetonitrile was evaporated, the rest was dissolved in EtOAc and the insoluble solids were filtered out. The resulting  
41 solution of EtOAc was evaporated and the crude of **2s** was purified by column chromatography (Hexane : EtOAc - 1 : 4,  $R_f = 0.7$ ).  
42 Yield = 84.2% (7.4 g).

43  
44 The product is a transparent colorless oil,  $[\alpha]_D = +20.2$  (MeOH,  $c = 33.87$  mmol/L).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta = 1.46$  (s, 3H),  
45 1.51 (s, 6H), 2.21-2.26 (m, 1H), 2.45-2.50 (m, 1H), 3.59-3.78 (m, 2H), 3.79 (s, 3H), 4.39-4.50 (m, 1H), 4.90 (s, 1H), 6.28 (t, 1H,  
46  $^2J_{\text{H-F}} = 74.8$  Hz,  $\text{CF}_2\text{H}$ ).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -84.0- -82.8 (m, 2F).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 28.2, 28.3, 36.2, 37.2, 52.1, 52.3, 71.2, 71.9, 80.5, 115.6 (t, <sup>1</sup>J<sub>C-F</sub> = 262.8 Hz, CF<sub>2</sub>H), 153.4, 154.1, 172.6.

Anal. calcd for C<sub>12</sub>H<sub>19</sub>F<sub>2</sub>NO<sub>5</sub>: C, 48.83; H, 6.49; N, 4.74. Found: C, 48.68; H, 6.45; N, 4.69.

**(2*S*,4*R*)-1-(tert-butoxycarbonyl)-4-(difluoromethoxy)pyrrolidine-2-carboxylic acid (3s).**

Compound **2s** (2.75 g, 0.0093 mol, 1 eq) was dissolved in 40 mL of THF and the solution of LiOH·xH<sub>2</sub>O (0.47 g, 0.0112 mol, 1.2 eq) in 10 mL of water was added all at once at 0 °C. The reaction mixture was heated to room temperature and was stirred for 4 h. Afterwards, the reaction mixture was diluted with 75 mL of EtOAc and 40 mL of water. The organic layer was separated, washed twice with 20 mL of water. Combined water solution was washed once with 50 mL of EtOAc, carefully acidified with 35 mL of conc. solution of citric acid and extracted three times with 40 mL of EtOAc. Combined organic layer was washed with brine, dried over magnesium sulphate and concentrated under vacuum to give compound **3s**. Yield = 61.1% (1.6 g).

The product is a white solid, mp = 70-75 °C, [α]<sub>D</sub> = -36.6 (EtOH, c = 35.55 mmol/L). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 1.14 (s, 3H), 1.45 (s, 6H), 2.23-2.47 (m, 2H), 3.56-3.68 (m, 2H), 4.35-4.42 (m, 1H), 4.83 (s, 1H<sub>idine</sub>), 6.21 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 73.5 Hz, CF<sub>2</sub>H), 8.70 (br. s, 1H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -84.0 - -82.9 (m, 2F).

<sup>13</sup>C{<sup>1</sup>H} NMR (125 MHz, CDCl<sub>3</sub>): δ = 28.2, 28.3, 35.5, 37.2, 52.1, 52.5, 71.2, 71.5, 81.0, 81.8, 115.6 (t, <sup>1</sup>J<sub>C-F</sub> = 262.6 Hz, CF<sub>2</sub>H), 153.6, 155.7, 177.8.

Anal. calcd for C<sub>11</sub>H<sub>17</sub>F<sub>2</sub>NO<sub>5</sub>: C, 46.97; H, 6.09; N, 4.98. Found: C, 47.09; H, 6.15; N, 4.91.

**(*S*)-3-(difluoromethoxy)pyrrolidin-2-one (5).**

Compound **2r** (1.5 g, 0.0053 mol, 1 eq) was dissolved in 5 mL of DCM and cooled to 0 °C using ice bath. Afterwards trifluoroacetic acid (5 mL, 0.0653 mol, 12.3 eq) was added all at once. The reaction mixture was stirred for 30 minutes at 0 °C and then basified with conc. solution of potassium carbonate. Obtained suspension was extracted three times with 30 mL of EtOAc. Combined organic layer was washed with brine, dried with magnesium sulphate and concentrated under vacuum. The crude product was purified by flash-chromatography (Hexane : EtOAc - 5 : 1) to give compound **5**. Yield = 52.5% (0.42 g).

The product is a white solid, mp = 57-61 °C, [α]<sub>D</sub> = -72.9 (MeOH, c = 66.18 mmol/L). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ = 2.16-2.53 (m, 2H), 3.28-3.45 (m, 2H), 4.67 (t, 1H, <sup>3</sup>J = 8.1 Hz), 6.48 (t, 1H, <sup>2</sup>J<sub>H-F</sub> = 75.8 Hz), 7.50 (br. s, 1H).

<sup>19</sup>F NMR (376 MHz, CDCl<sub>3</sub>): δ = -84.3 (dd, F, <sup>2</sup>J<sub>H-F</sub> = 158.5 Hz, <sup>2</sup>J<sub>H-F</sub> = 75.7 Hz), -83.4 (dd, F, <sup>2</sup>J<sub>F-F</sub> = 158.5 Hz, <sup>2</sup>J<sub>H-F</sub> = 73.5 Hz).

<sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>): δ = 28.7, 38.8, 70.8, 115.8 (t, <sup>1</sup>J<sub>C-F</sub> = 258.4 Hz, CF<sub>2</sub>H), 173.7.

Anal. calcd for C<sub>5</sub>H<sub>7</sub>F<sub>2</sub>NO<sub>2</sub>: C, 39.74; H, 4.67; N, 9.27. Found: C, 39.80; H, 4.71; N, 9.33.

## ASSOCIATED CONTENT

### Supporting Information

Detailed experimental procedures and characterization data for all new compounds. This material is available free of charge *via* the Internet at <http://pubs.acs.org>.

## AUTHOR INFORMATION

### Corresponding Authors

\* E-mail:

[iva108@gmail.com](mailto:iva108@gmail.com) (VI),

[viktori@cbmm.lodz.pl](mailto:viktori@cbmm.lodz.pl) (VI),

[Pavel.Mykhailiuk@gmail.com](mailto:Pavel.Mykhailiuk@gmail.com) (PM),

[Pavel.Mykhailiuk@mail.enamine.net](mailto:Pavel.Mykhailiuk@mail.enamine.net) (PM).

## REFERENCES

- (a) Kirk, K. L. *J. Fluorine Chem.* **2006**, *127*, 1013–1029; (b) Hagmann, W. K. *J. Med. Chem.* **2008**, 4359–4369; (c) Begue, J.-P.; Bonnet-Delpont, D.; *Bioorganic and Medicinal Chemistry of Fluorine*, John Wiley & Sons, 1th edn., **2008**, 366, ISBN-10: 0470278307; (d) Gouverneur, V.; Müller, K.; *Fluorine in Pharmaceutical and Medicinal Chemistry*, Imperial College Press **2012**, ISBN-10: 1-84816-634-6; (e) Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soshonok, V. A.; Izawa, K.; Liu, H. *Chem. Rev.* **2016**, *116*, 422–518.
- (a) Isanbor, C.; O'Hagan, D. *J. Fluorine Chem.* **2006**, *127*, 303–319; (b) Jeschke, P. *ChemBioChem* **2004**, *5*, 570–589; (c) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881–1886; (d) Böhm, H.-J.; Banner, D.; Bendels, S.; Kansy, M.; Kuhn, B.; Müller, K.; Obst-Sander, U.; Stahl, M. *ChemBioChem* **2004**, *5*, 637–643; (e) Shah, P.; Westwell, A. D. *J. Enzyme Inhib. Med. Chem.* **2007**, *22*, 527–540; (f) Liu, P.; Sharon, A.; Chu, C. K. *J. Fluorine Chem.* **2008**, *129*, 743–766.
- Silverman, R. B.; *The organic chemistry of drug design, and drug action*, Elsevier Academic Press, 2<sup>nd</sup> edn., **2004**, 617, ISBN 0-12-643732-7; (b) Palmer, M.; Honek, J. F.; Dieckmann, T. *Biochemical Pharmacology*, John Wiley & Sons, 1<sup>th</sup> edn., **2012**, 428, ISBN-13: 978-0-470-17445-6.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
4. For isosteric substitution of hydrogen with fluorine, see: (a) Meanwell, N. A. *J. Med. Chem.* **2011**, *54*, 2529–2591; (b) Wang, J.; Sanchez-Roselló, M.; Acena, J. L.; del Pozo, C.; Sorochinsky, A. E.; Fustero, S.; Soloshonok, V. A.; Liu, H. *Chem. Rev.* **2014**, *114*, 2432–2506; (c) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320–330.
  5. For fluorine enhanced catalysts, see: (a) Bahmanyar, S.; Houk, K. N.; Martin, H. J.; List, B. *J. Am. Chem. Soc.* **2003**, *125*, 2475–2479; (b) Chandler, C. L.; List, B. *J. Am. Chem. Soc.* **2008**, *130*, 6737–6739; (c) Adam, W.; Curci, R.; Edwards, J. O. *Acc. Chem. Res.* **1989**, *22*, 205–211; (d) Mello, R.; Fiorentino, M.; Fusco, C.; Curci, R. *J. Am. Chem. Soc.* **1989**, *111*, 6749–6757.
  6. For fluorine enhanced catalysts, see: (a) Armstrong, A.; Hayter, B. R. *Chem. Commun.* **1998**, 621–622; (b) Armstrong, A.; Ahmed, G.; Dominguez-Fernandez, B.; Hayter, B. R.; Wailes, J. S. *J. Org. Chem.* **2002**, *67*, 8610–8617; (c) Denmark, S. E.; Wu, Z.; Crudden, C. M.; Matsushashi, H. *J. Org. Chem.* **1997**, *62*, 8288–8289; (d) Denmark, S. E.; Matsushashi, H. *J. Org. Chem.* **2002**, *67*, 3479–3486; (e) Yang, D.; Wong, M. K.; Yip, Y. C. *J. Org. Chem.* **1995**, *60*, 3887–3889; (f) D. Yang, *Acc. Chem. Res.* **2004**, *37*, 497–505; (g) Brown, D. S.; Marples, B. A.; Smith, P.; Walton, L. *Tetrahedron*, **1995**, *51*, 3587–3606.
  7. For our ongoing medicinal chemistry program on fluorine, see: (a) Mykhailiuk, P. K.; Afonin, S.; Palamarchuk, G. V.; Shishkin, O. V. Ulrich, A. S. Komarov, I. V. *Angew. Chem. Int. Ed.* **2008**, *47*, 5765–5767; (b) Kubyshkin, V. S.; Mykhailiuk, P. K.; Afonin, S.; Ulrich, A. S.; Komarov, I. V. *Org. Lett.* **2012**, *14*, 5254–5257; (c) Kubyshkin, V. S.; Afonin, S.; Kara, S.; Budisa, N.; Mykhailiuk, P. K.; Ulrich, A. S. *Org. Biomol. Chem.* **2015**, *13*, 3171–3181; (d) P. K. Mykhailiuk *Chem. Eur. J.* **2014**, *17*, 4942–4947; (e) Bandak, D.; Babii, O.; Vasiuta, R.; Komarov, I. V.; Mykhailiuk, P. K. *Org. Lett.* **2015**, *17*(2), 226–229; (f) Mykhailiuk P. K. *Angew. Chem. Int. Ed.* **2015**, *22*, 6558–6561; (g) Iaroshenko, V. O.; Ostrovskiy, D.; Petrosyan, A.; Mkrtchyan, S.; Villinger, A.; Langer, P. *J. Org. Chem.* **2011**, 2899–2903; (h) Iaroshenko, V. O.; Sevenard, D. V.; Kotljarov, A. V.; Volochnyuk, D. M.; Tolmachev, A. O.; Sosnovskikh, V. Ya. *Synthesis* **2009**, 731–740; (i) Slobodyanyuk, E. Y.; Artamonov, O. S.; Shishkin, O. V.; Mykhailiuk, P. K. *Eur. J. Org. Chem.* **2014**, 2487–2495;
  8. (a) Chauret, N.; Guay, D.; Li, C.; Day, S.; Silva, J.; Blouin, M.; Ducharme, Y.; Yergey, J. A.; Nicoll-Griffith, D. A. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 2149–2152; (b) Ohmine, T.; Katsube, T.; Tsuzaki, Y.; Kazui, M.; Kobayashi, N.; Komai, T.; Hagihara, M.; Nishigaki, T.; Iwamoto, A.; Kimura, T.; Kashiwase, H.; Yamashita, M.; *Bioorg. Med. Chem. Lett.* **2002**, *12*, 739–742; (c) Takahata, M.; Mitsuyama, J.; Yamashiro, Y.; Yonezawa, M.; Araki, H.; Todo, Y.; Minami, S.; Watanabe, Y.; Narita, H. *Antimicrob. Agents Chemother.* **1999**, *43*, 1077–1084.
  9. For conformational preferences of OCF<sub>2</sub>-R substituents, see: (a) Praly, J.-P.; Brard, L.; Descotes, G.; Toupet, L. *Tetrahedron* **1989**, *45*, 4141–4152.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
10. Lovering, F.; Bikker, J.; Humblet, C. *J. Med. Chem.*, **2009**, *52* (21), 6752–6756.
  11. (a) Ni, C.; Hu, J. *Synthesis* **2014**, *46*, 0842–0863; (b) Zafrani, Y.; Sod-Moriah, G.; Segall, Y. *Tetrahedron* **2009**, *65*, 5278–5283; (c) Zheng, J.; Li, Y.; Zhang, L.; Hu, J.; Meuzelaar, G. J.; Federsel, H.-J. *Chem. Commun.* **2007**, 5149–5151; (d) Fier, P. S.; Hartwig, J. F. *Angew. Chem. Int. Ed.* **2013**, *52*(7), 2092–2095; (e) Hartwig, J.; Fier, P. **2014**, WO 2014107380 A1 Jul 10, 2014; (f) Xiaoxi L.; Zhiqiang, W. *Org. Biomol. Chem.*, **2015**, *13*, 3432–3437.
  12. (a) Tortelli, V.; Calini, P.; Zompatori, A.; Antenucci, E. **2007**, EP 1801090 A2 Jun 27, 2007.
  13. For the transformation OCO, OCS to OCF<sub>2</sub>, see: (a) Slassi, A.; Dove, P. **2014**, WO 2014026282 A1 Feb 20, 2014; (b) Dmowski, W.; Kaminski, M. *J. Fluorine Chem.* **1983**, *23*(3), 207–18; (c) Dolbier, W. R. Jr.; Wang, F.; Tang, X.; Thomson, C. S.; Wang, L. *J. Fluorine Chem.* **2014**, *160*, 72–76.
  14. O'Neill, G. J.; Bulka, R. J. **1993**, EP 562858 A1 Sep 29, 1993.
  15. T. Miller, G. Thanassi, J. W. *J. Org. Chem.* **1960**, *25*, 2009–2012.
  16. Chen, Q.-Y.; Wu, S.-W. *J. Fluorine Chem.* **1989**, *44*, 433–440.
  17. Chen, Q.; Wu, S. *J. Org. Chem.* **1989**, *54*, 3023–3027.
  18. (a) Zheng, J.; Lin, J.-H.; Yu, L.-Y.; Wei, Y.; Zheng, X.; Xiao, J.-C. *Org. Lett.* **2015**, *17*, 6150–6153; (b) Gulevich, A. V.; Gevorgyan, V.; *Angew. Chem., Int. Ed.* **2013**, *52*, 1371–1373; (c) Selander, N.; Fokin, V. V. *J. Am. Chem. Soc.*, **2012**, *134* (5), 2477–2480; (d) Helan, V.; Gulevich, A. V. Gevorgyan, A. *Chem. Sci.*, **2015**, *6*, 1928–1931; (e) Werner, H. *Angew. Chem. Int. Ed.* **2010**, *49*, 4714–4728; (f) Zinn, F. K.; Viciu, M. S.; Nolan, S. P. *Annu. Rep. Prog. Chem., Sect. B*, **2004**, *100*, 231–249; (g) Whited, M T.; Grubbs, R. H. *Acc. Chem. Res.* **2009**, 1607–1616.
  19. (a) Takahira, Y.; Morizawa, Y. *J. Am. Chem. Soc.*, **2015**, *137* (22), 7031–7034; (b) Feng, Z.; Min, Q.-Q.; Zhang, X. *Org. Lett.*, **2016**, *18* (1), 44–47.
  20. (a) Tomashenko, O. A.; Grushin, V. V. *Chem. Rev.*, **2011**, *111* (8), 4475–4521; (b) Ye, Y.; Sanford, M. S.; *Synlett* **2012**, *23*, 2005–2013; (c) Mormino, M. G.; Fier, P. S.; Hartwig, J. F. *Org. Lett.* **2014**, *16*, 1744–1747.
  21. (a) For preparation of **1r**, see: DeWit, M. A.; Gillies, E. R. *Org. Biomol. Chem.*, **2011**, *9*, 1846–1854; (b) For preparation of **1s**, see: Grygorenko, O. O.; Komarov, I. V.; Cativiela, C. *Tetrahedron: Asymetry*, **2009**, *20* (12), 1433–1436.
  22. For our previous interest in fluorinated prolines and their use in peptide studies, see the literature 7a-c as well as the following literature sources: (a) Mykhailiuk, P. K.; Shishkina, S. V.; Shishkin, O. V.; Zaporozhets, O. A.; Komarov, I. V. *Tetrahedron* **2011**, *67*, 3091–3097; (b) Kubyshkin, V. S.; Mykhailiuk, P. M.; Afonin, S.; Grage, S. L.; Komarov, I. V.; Ulrich, A. S. *J. Fluorine Chem.* **2013**, *152*, 136–143.
  23. Dolbier, W. R. *Guide to Fluorine NMR for Organic Chemists*, John Wiley & Sons, 1th edn., **2009**, 272, ISBN: 978-0-470-1934