

## #-Carboline Amides as Intrinsic Directing Groups for C(sp<sup>2</sup>)-H Functionalization

Hel ne M.-F Viart, Andreas Bachmann, William Kayitare, and Richmond Sarpong

*J. Am. Chem. Soc.*, **Just Accepted Manuscript** • Publication Date (Web): 01 Jan 2017

Downloaded from <http://pubs.acs.org> on January 1, 2017

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

# $\beta$ -Carboline Amides as Intrinsic Directing Groups for C(sp<sup>2</sup>)-H Functionalization

Hélène M.-F. Viart,<sup>†</sup> Andreas Bachmann, William Kayitare and Richmond Sarpong\*

Department of Chemistry, University of California, Berkeley, California 94720, United States

**ABSTRACT:** Many site-selective palladium catalyzed C-H functionalization methods require directing groups. We report here  $\beta$ -carboline amides as intrinsic directing groups for C(sp<sup>2</sup>)-H functionalization. Various substrates including the natural product alaniobussinine and the marinacarboline core structure were functionalized using carboline-directed  $\delta$ -C(sp<sup>2</sup>)-H alkynylations. This transformation proceeds under mild conditions and is compatible with a wide variety of  $\beta$ -arylethylamines.  $\delta$ -Alkynylation of  $\beta$ -arylethylamines via a six-membered palladacycle is favored over  $\gamma$ -C(sp<sup>2</sup>)-H bond functionalization when both positions are accessible. The versatility of  $\beta$ -carboline amides as directing groups is evidenced by other  $\delta$ -C(sp<sup>2</sup>)-H functionalizations such as alkenylation, arylation, and C-N bond formation.

## 1) INTRODUCTION

The emergence of powerful transition metal-catalyzed C-H functionalization methods over the last decade and a half has provided new tools for chemical synthesis. Despite continuing efforts to identify traceless directing groups that can be introduced and removed under the conditions used in the functionalization reaction,<sup>1</sup> most selective C-H functionalization methods exploit intrinsic reactivity differences between C-H bonds in a substrate, coupled with the ability of auxiliaries (directing groups) to direct metallation at specific positions. In the majority of cases, it is necessary to introduce the auxiliary prior to C-H activation and then remove it at the end of the functionalization. Therefore, typical drawbacks of this approach are an increased number of synthetic manipulations, as well as the challenges inherent in introducing and removing the directing group. Generally, directing groups are appended to functional groups common in complex molecules, where C-H functionalization technology is increasingly employed.<sup>2</sup>


Drawing inspiration from substrate-controlled (diastereoselective) processes, where the controlling element for stereoselectivity is inherent to the substrate, we envisioned an analogous scenario for directed C-H functionalization, where the directing group is resident in the substrate and in the product. In this way, the requirement for removing the directing group post C-H functionalization is obviated. For this purpose, we have been especially drawn to directing groups based on pyridine that utilize 2-point binding.

The picolinamide and 8-aminoquinoline amide, first reported by Daugulis for the palladium-catalyzed  $\gamma$ - and  $\beta$ -arylation of aliphatic amine and carboxylic acid derivatives, respectively,<sup>3</sup> have emerged as a privileged directing group. However, the removal of these directing groups has historically been challenging. Several solutions to this challenge are beginning to appear. For example, recently, Chen and coworkers report-

ed two variants of the picolinamide and 8-aminoquinoline amide directing groups: the TBS-protected methylene hydroxyl group at the ortho position of the picolinamide (PAR),<sup>4</sup> and 8-amino-5-methoxyquinoline (MQ) directing groups,<sup>5</sup> which are easily cleaved through intramolecular acyl transfer under acidic conditions or with ceric ammonium nitrate under mild conditions, respectively.


On the contrary, instances where a picolinamide directing group is employed in C-H functionalization and retained in the targeted product are rare. Only a limited number of examples of bidentate directing groups that are found in or easily derived to afford a target compound have been described. Examples are mostly found for C(sp<sup>3</sup>)-H functionalizations in the synthesis and derivatizations of peptides and peptidomimetics, where the auxiliary can be either an amino acid itself,<sup>6</sup> or transformed into an amino acid derivative. For instance, the 2-methoxyiminoacetyl (MIA) directing group developed by Fan and Ma has been transformed into a glycine moiety by hydrogenation,<sup>7</sup> and the  $\alpha$ -amino oxazolonyl directing group developed by Shi and coworkers was opened and derivatized into amides in a mild, one-pot procedure.<sup>8</sup>

We envisioned  $\beta$ -carboline amides as inherent directing groups that afford many of the advantages of these previously described directing groups. Significantly,  $\beta$ -carbolines and their associated derivatives, which may be accessed in a single step, are found in a number of natural products and pharmaceutically relevant molecules (Figure 1).<sup>9</sup> As such, these 'functional groups' that are innate to the targeted compound may be employed as directing groups for C-H functionalization.



**Figure 1.** Examples of natural products and biologically active compounds featuring the  $\beta$ -carboline amide motif.

In the context of C–H functionalizations directed by  $\beta$ -carboline amides, we envisioned alkynylations to be particularly useful. Alkynes are versatile building blocks that may be readily transformed into a wide range of functional groups and substituents. Their importance is evidenced by the success of the Sonogashira cross-coupling reaction. Thus, C–H functionalization methods to complement this well-established cross-coupling methodology<sup>10</sup> would be highly useful.




While directed  $\gamma$ -C(sp<sup>2</sup>)-H alkynylations abound,<sup>11</sup> to the best of our knowledge, there has only been one report describing the palladium-catalyzed *ortho* C(sp<sup>2</sup>)-H alkynylation of aminoalkyl arene derivatives at the  $\delta$  position, which relies on an oxalamide directing group.<sup>12</sup> As reported herein, using a  $\beta$ -carboline amide directing group, we have achieved selective palladium-catalyzed  $\gamma$ - and  $\delta$ -C(sp<sup>2</sup>)-H alkynylations via five- and six-membered palladacycle intermediates.<sup>12–14</sup> Our method has been applied to the syntheses of derivatives of the natural products alangiobussinine and marinacarboline.

## 2) RESULTS AND DISCUSSION

Our investigations commenced with studies on the directed alkylation of alangiobussinine (**2a**), which was prepared in a single step from the known  $\beta$ -carboline-1-methyl ester **1** by activation of the ester moiety using 2-hydroxypyridine under aerobic conditions (Eq 1).<sup>15</sup> Our preparation of **2a** from **1** is an improvement in terms of step count and isolated yield over the previously reported synthesis of this natural product.<sup>16</sup> Applying this 2-hydroxypyridine mediated amidation method, a variety of other  $\beta$ -carboline amide derivatives were accessed in good yields from the corresponding methylesters. Alternative-

ly, La(OTf)<sub>3</sub> could also be employed for the amidation reactions (see the SI for details).<sup>17</sup>



Alangiobussinine was subjected to the conditions for Pd-catalyzed picolinamide-directed C(sp<sup>2</sup>)-H alkylation reported by Chen and coworkers.<sup>11b</sup> The desired alkylnated product (**4a**) was isolated in 41% yield (Table 1, entry 1). A survey of carboxylic acid additives known to play an important role in the proposed concerted palladation-deprotonation step,<sup>18</sup> showed that strong acids were detrimental, whereas acid additives with intermediate pKa (3.5–5) resulted in increased formation of the alkylnated product. *o*-Phenyl benzoic acid (*o*-PBA) has been shown to be an effective carboxylic acid promoter for C(sp<sup>3</sup>)-H arylation reactions. In our hands, *o*-PBA and PivOH led to comparable outcomes (Table 1, entries 2–3 and 18–19).<sup>11b,18</sup> A survey of various bases revealed that the nature of the counterion has a significant impact on the reaction outcome with K<sub>2</sub>CO<sub>3</sub> giving the best yield (entries 7–18). Surprisingly, silver salts, which are generally believed to act as halogen scavengers and are often beneficial in metal-catalyzed C–H functionalizations, shut down the reaction (entries 8 and 13). Finally, a clear solvent effect was observed: coordinating solvents such as DMA, THF, and dioxane impede the reaction, giving moderate yields, whereas chlorinated solvents were superior, with DCE providing the best yields (see the SI for details).


**Table 1.** Optimization of reaction conditions<sup>a</sup>

2a


3

4a

Entry	Base	Additive	Yield (%) <sup>b,c</sup>
1 <sup>d</sup>	KHCO <sub>3</sub>	<i>o</i> -PBA	41 <sup>e</sup>
2	KHCO <sub>3</sub>	<i>o</i> -PBA	72 (18)
3	KHCO <sub>3</sub>	PivOH	69 (26)
4	KHCO <sub>3</sub>	AcOH	69 (25)
5	KHCO <sub>3</sub>	TFA	13 (55)
6	KHCO <sub>3</sub>	DMBA	65 (22)
7	KHCO <sub>3</sub>	<i>o</i> -MBA	64 (27)
8	AgOAc	<i>o</i> -PBA	<5 (>95)
9	NaOAc	<i>o</i> -PBA	30 (56)
10	NaOTf	<i>o</i> -PBA	7 (64)
11	Cs <sub>2</sub> CO <sub>3</sub>	<i>o</i> -PBA	7 (80)
12	K <sub>2</sub> CO <sub>3</sub>	<i>o</i> -PBA	89 (<5)
13	Ag <sub>2</sub> CO <sub>3</sub>	<i>o</i> -PBA	4 (71)
14	<sup>t</sup> BuOK	<i>o</i> -PBA	23 (68)
15	NaHCO <sub>3</sub>	<i>o</i> -PBA	7 (83)
16	Na <sub>2</sub> CO <sub>3</sub>	<i>o</i> -PBA	63 (28)
17	KOAc	<i>o</i> -PBA	77 (15)
18	K <sub>2</sub> CO <sub>3</sub>	<i>o</i> -PBA	89 (5)



Under our optimized conditions, alkynylated alangiobus-sinine derivative **4a** was isolated in 78% yield (Eq 2) and its structure unambiguously confirmed by X-ray crystallographic analysis.<sup>19</sup>




Using the optimized conditions (1.2 equiv (bromoethynyl)triisopropylsilane, 10 mol% Pd(OAc)<sub>2</sub>, 0.2 equiv PivOH, 2.0 equiv K<sub>2</sub>CO<sub>3</sub>, DCE, 100 °C, 15 h),<sup>20</sup> a variety of β-carboline amide derivatives were successfully alkynylated at the δ position (Table 2). Both electron-deficient and electron-rich substrates bearing substituents at the *ortho*- or *meta*-positions were alkynylated in excellent yields favoring the less sterically-hindered *ortho*-position (**4b–4j**). In some cases, small amounts of dialkynylated products were formed and isolated when *meta*-substituted substrates were employed (see **4g** and **4j**), consistent with observations in other C–H alkynylation reactions.<sup>12,11c</sup> Phenethylamine substrates lacking *ortho*- or *meta*-substitution (**4k–4n**) gave mixtures of mono- and dialkynylated products under the standard conditions, but selectivity toward the mono- or di-alkynylated product could be achieved by varying the equivalents of the bromoalkyne (see the SI for experiments aimed at the selective synthesis of **4n<sub>mono</sub>** and **4n<sub>(o+o')di</sub>**). Various functional groups are tolerated under the reaction conditions, including bromide, nitrile, and trifluoromethyl (**4c**, **4l**, **4h**, and **4m**). Therefore, scaffolds displaying a bromide may be further functionalized using Pd<sup>0</sup> cross-coupling processes. Heteroaromatic compounds were alkynylated in good to excellent yields (see **4a**, **4u**). In particular, the presence of a protecting group on the indole nitrogen (see **2a**) is not required, in stark contrast with previously reported alkynylations of tryptamine derivatives.<sup>12</sup>


The vast majority of directed C–H functionalizations, including picolinamide-directed functionalizations, favor γ-functionalization. Presumably, a kinetically favored five-membered palladacycle predominates. However, the β-carboline amide directing group favors functionalization at the δ position via a six-membered palladacycle. We observe selectivity toward δ-functionalization when functionalization via five- and six-membered palladacycles is possible. For example, **4t<sub>mono</sub>** and **4t<sub>(o+o')di</sub>** were the sole products of C–H functionalization of the precursor substrate. The identity of the dialkynylated product (**4t<sub>(o+o')di</sub>**) was unambiguously confirmed by X-ray crystallographic analysis of the di-alkynylated product after cleavage of the TIPS protecting groups (Eq 3).<sup>19</sup> Furthermore, we have found that products that arise from functionalization at the γ-position require slightly more demanding conditions (e.g., for **4p–4s**, functionalizations occurred at 120 °C instead of 100 °C).

The β-carboline-directed C–H functionalization has proven to be highly general and versatile for C(sp<sup>2</sup>)–H alkynylation, and has been extended to alkene functionalization (see **4o**).

**Table 2. Substrate scope of the Pd-catalyzed C–H alkynylation reaction.<sup>a</sup>**




<sup>a</sup>Determined by NMR analysis of the crude mixture using 1,1,2,2-tetrachloroethane as internal standard. Isolated yields shown in parentheses. <sup>b</sup>2.0 equiv **3** used instead at 120 °C for 15 h. <sup>c</sup>at 120 °C for 15 h. <sup>d</sup>3.0 equiv **3**, 140 °C for 30 min by microwave heating.



To highlight the scalability of this reaction, the carboline amide derivative **2e** was alkynylated on 1.0 g (2.90 mmol) scale to afford **4e** in 80% isolated yield. Further derivatizations of **4e** were achieved on terminal alkyne **5e**, obtained in 88% yield by TBAF mediated cleavage of the TIPS group (Scheme 1). Alkyne **5e** was easily converted to triazole **6** by a copper-catalyzed ‘click’ cycloaddition,<sup>21</sup> hydrogenated to the corresponding alkane (**7**) using Pd/C, and converted to aldehyde **8** through an anti-Markovnikov hydration using Grotjahn’s ruthenium complex.<sup>22</sup>


### Scheme 1. Derivatizations of alkyne **5**.



To gain insight into the mechanism of the  $\beta$ -carboline directed functionalization, **2a** was subjected to stoichiometric palladium acetate in DMA in the presence of *tert*-butyl isonitrile. The six-membered palladacycle (**9**) was isolated in 51% yield and unambiguously characterized by X-ray crystallographic analysis (Scheme 2B).<sup>19</sup>

We propose that following C–H activation by a  $\text{Pd}^{\text{II}}$  species to generate a six-membered palladacycle similar to **9**, oxidative addition into the TIPS-protected acetylene bromide generates a  $\text{Pd}^{\text{IV}}$  intermediate, consistent with the observations of Shi and coworkers.<sup>11d</sup> Reductive elimination at this stage provides the alkynylated product. Finally, ligand exchange regenerates the requisite  $\text{Pd}^{\text{II}}$  species to perpetuate the catalytic cycle (Scheme 2A).<sup>23</sup>


### Scheme 2. A) Proposed mechanism; B) Evidence for C–H activation via six-membered palladacycle.



Because a number of pharmaceutically relevant compounds and natural products possess an N-alkylated  $\beta$ -carboline-1-carboxylic acid N-alkylamide (e.g., **2v**) or  $\beta$ -carboline-3-carboxylic acid N-alkylamide (e.g., **2w**) scaffold (see Figure 1), we sought to expand the carboline-directed alkynylations described above to **2v** and **2w** (Scheme 3). Both alkynylated products were successfully isolated. Of note, the presence of the methyl group on the carboline nitrogen slightly decreased


the efficiency of the reaction (**4a** vs **4v**), and higher temperatures were required to access **4w** in good yield.

### Scheme 3. Expansion of the $\text{C}(\text{sp}^2)$ –H alkylation reaction to other $\beta$ -carboline amide derivatives.



Gratifyingly, the  $\beta$ -carboline amide group directs other C–H functionalization reactions (Scheme 4). For example, using **2e**, arylated and alkenylated products were isolated in high yields, although more vigorous conditions were necessary (see **2e**→**10** and **2e**→**12**). A C–N bond forming reaction could also be achieved under mild conditions to afford the indoline derivative **11**.

### Scheme 4. Generalization of $\beta$ -carboline amide as directing group for various $\text{C}(\text{sp}^2)$ –H functionalizations



### 3) CONCLUSION

In conclusion, we report  $\beta$ -carboline amides as a new class of directing groups for a variety of C–H functionalizations that may proceed via five- or six-membered palladacycle intermediates. In particular, we report alkynylation reactions under mild conditions that achieve the functionalization of various  $\beta$ -arylethamine substrates and the natural product alangiobus-sinine. The synthetic utility of the method was highlighted by the direct derivatization of the alkynylated product into useful groups such as a triazole or aldehyde. The generality of the method is highlighted by the ability of the  $\beta$ -carboline amide to direct other functionalizations such as  $\delta$ -alkenylation,  $\delta$ -

arylation, and a C–N bond formation to generate an indoline derivative. Finally, we have demonstrated that  $\beta$ -carboline amides can also direct  $\gamma$ -alkynylation, albeit under slightly more demanding conditions, and exclusive  $\delta$ -alkynylation is achieved in the presence of a more accessible  $\gamma$ -C(sp<sup>2</sup>)–H bond.

## ASSOCIATED CONTENT

### Supporting Information

Experimental details and spectroscopic data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Crystallographic data (CIF, CIF, CIF)

Procedures, single crystal X-ray data, NMR spectra (PDF)

## AUTHOR INFORMATION

### Corresponding Author

\* [rsarpong@berkeley.edu](mailto:rsarpong@berkeley.edu)

### Present Addresses

† The Institute for Neurodegenerative Diseases, University of California, San Francisco, Sandler Neurosciences Center, 675 Nelson Rising Lane, San Francisco, CA 94143-0518

### Notes

The authors declare no competing financial interests.

## ACKNOWLEDGMENT

This work was supported by the NSF under the CCI Center for Selective C–H Functionalization (CHE-1205646). We are grateful to the Carlsberg Foundation for a postdoctoral scholarship to H.M.-F.V., and to the NSF CCI Center for Selective C–H Functionalization for a summer fellowship for W.K. We thank A. DiPasquale (UC Berkeley) for solving the crystal structures of **4a**, **5**<sub>(ortho)di</sub> and **9** (displayed with CYLView), supported by NIH Shared Instrumentation Grant (S10-RR027172).

## REFERENCES

- Jun, C.-H.; Lee, H.; Hong, J.-B. *J. Org. Chem.* **1997**, *62*, 1200–1201.
- Mo, F.; Dong, G. *Science* **2014**, *345*, 68.
- Bedford, R. B.; Coles, S. J.; Hursthouse, M. B.; Limmert, M. E. *Angew. Chem. Int. Ed.* **2003**, *42*, 112.
- Lightburn, T. E.; Dombrowski, M. T.; Tan, K. L. *J. Am. Chem. Soc.* **2008**, *130*, 9210.
- Grünanger, C. U.; Breit, B. *Angew. Chem. Int. Ed.* **2008**, *47*, 7346.
- Luo, J.; Preciado, S.; Larrosa, I. *J. Am. Chem. Soc.* **2014**, *136*, 4109.
- Zhang, F.-L.; Hong, K.; Li, T.-J.; Park, H.; Yu, J.-Q. *Science* **2016**, *351*, 252.
- For selected reviews and examples of complex molecules synthesis via C–H functionalization, see a) Feng, Y.; Chen, G. *Angew. Chem. Int. Ed.* **2010**, *49*, 958. b) Gutekunst, W. R.; Baran, P. S. *Chem. Soc. Rev.* **2011**, *40*, 1976, and references therein. c) McMurray, L.; O'Hara, F.; Gaunt, M. J. *Chem. Soc. Rev.* **2011**, *40*, 1885, and references therein. d) Ting, C. P.; Maimone, T. J. *Angew. Chem. Int. Ed.* **2014**, *53*, 3115. e) Dailler, D.; Danoun, G.; Baudoin, O. *Angew. Chem. Int. Ed.* **2015**, *54*, 4919.
- Zaitsev, V. G.; Shabashov, D.; Daugulis, O. *J. Am. Chem. Soc.* **2005**, *127*, 13154.
- a) He, G.; Chen, G. *Angew. Chem.* **2011**, *123*, 5298; *Angew. Chem. Int. Ed.* **2011**, *50*, 5192. b) He, G.; Zhao, Y.; Zhang, S.; Lu, C.; Chen, G. *J. Am. Chem. Soc.* **2012**, *134*, 3. c) Zhang, S.-Y.; He, G.; Zhao, Y.; Wright, K.; Nack, W. A.; Chen, G. *J. Am. Chem. Soc.* **2012**, *134*, 7313. d) Zhang, S.-Y.; He, G.; Nack, W. A.; Zhao, Y.; Li, Q.; Chen, G. *J. Am. Chem. Soc.* **2013**, *135*, 2124.
- a) He, G.; Zhang, S.-Y.; Nack, W. A.; Li, Q.; Chen, G. *Angew. Chem. Int. Ed.* **2013**, *52*, 11124. b) Wang, B.; Nack, W. A.; He, G.; Zhang, S.-Y.; Chen, G. *Chem. Sci.* **2014**, *5*, 3952. c) Zhang, S.-Y.; Li, Q.; He, G.; Nack, W. A.; Chen, G. *J. Am. Chem. Soc.* **2015**, *137*, 531.
- Gong, W.; Zhang, G.; Liu, T.; Giri, R.; Yu, J.-Q. *J. Am. Chem. Soc.* **2014**, *136*, 16940.
- Fan, M.; Ma, D. *Angew. Chem. Int. Ed.* **2013**, *52*, 12152.
- Chen, K.; Li, Z.-W.; Shen, P.-X.; Zhao, H.-W.; Shi, Z.-J. *Chem. Eur. J.* **2015**, *21*, 7389.
- a) Wang, K.-B.; Di, Y.-T.; Bao, Y.; Yuan, C.-M.; Chen, G.; Li, D.-H.; Bai, J.; He, H.-P.; Hao, X.-J.; Pei, Y.-H.; Jing, Y.-K.; Li, Z.-L.; Hua, H.-M. *Org. Lett.* **2014**, *16*, 4028. b) Diallo, A. O.; Mehri, H.; Iouzalén, L.; Plat, M. *Phytochemistry* **1995**, *40*, 975. c) Huang, H.; Yao, Y.; He, Z.; Yang, T.; Ma, J.; Tian, X.; Li, Y.; Huang, C.; Chen, X.; Li, W.; Zhang, S.; Zhang, C.; Ju, J. *J. Nat. Prod.* **2011**, *74*, 2122. d) Xiao, Z.; Liu, Z.; Lei, C.; Wang, J.; Chen, Z.; Han, G.; Xiao, Y.; He, H.; Chen, S. China Pat. Appl. CN103130800A, June, 05, 2013; e) Clark, M.P.; Lockwood, M. A.; Wagner, F.F.; Natchus, M.G.; Doroh, B.C.; Johnson, T.L.; Tahirovic, Y. A.; Wilson, L.; Wiseman, J. M.; Skudlarek, J. W. PCT Int. Appl. WO2009121063A2 October 01, 2009; f) Rottmann, M.; McNamara, C.; Yeung, B.K.S.; Lee, M.C.S.; Zou, B.; Russell, B.; Seitz, P.; Plouffe, D.M.; Dharia, N.V.; Tan, J.; Cohen, S.B.; Spencer, K.R.; González-Páez, G.E.; Lakshminarayana, S.B.; Goh, A.; Suwanarusk, R.; Jegla, T.; Schmitt, E.K.; Beck, H.-P.; Brun, R.; Nosten, F.; Renia, L.; Dartois, V.; Keller, T.H.; Fidock, D.A.; Winzeler, E.A.; Diagona, T.T. *Science* **2010**, *329*, 1175.
- a) Sonogashira, K.; Tohda, Y.; Hagihara, N. *Tetrahedron Lett.* **1975**, *16*, 4467. b) Negishi, E.; Anastasia, L. *Chem. Rev.* **2003**, *103*, 1979. c) Chinchilla, R.; Nájera, C. *Chem. Soc. Rev.* **2011**, *40*, 5084.
- For palladium-catalyzed  $\gamma$ -C(sp<sup>2</sup>)–H alkynylations, see: a) Tobisu, M.; Ano, Y.; Chatani, N. *Org. Lett.* **2009**, *11*, 3250. b) Zhao, Y.; He, G.; Nack, W. A.; Chen, G. *Org. Lett.* **2012**, *14*, 2948. c) Ano, Y.; Tobisu, M.; Chatani, N. *Org. Lett.* **2012**, *14*, 354. d) Ye, X.; Xu, C.; Wojtas, L.; Akhmedov, N. G.; Chen, H.; Shi, X. *Org. Lett.* **2016**, *18*, 2970.
- for cobalt-catalyzed  $\gamma$ -C(sp<sup>2</sup>)–H alkynylations, see: e) Sauermann, N.; González, M. J.; Ackermann, L. *Org. Lett.*, **2015**, *17*, 5316. f) Landge, V. G.; Midya, S. P.; Rana, J.; Shinde, D. R.; Balaraman, E. *Org. Lett.* **2016**, *18*, 5252.
- For ruthenium-catalyzed  $\gamma$ -C(sp<sup>2</sup>)–H alkynylations, see: g) Ano, Y.; Tobisu, M.; Chatani, N. *Synlett* **2012**, 2763.
- For rhodium-catalyzed  $\gamma$ -C(sp<sup>2</sup>)–H alkynylations, see: h) Xie, F.; Qi, Z.-S.; Yu, S.-J.; Li, X.-W. *J. Am. Chem. Soc.* **2014**, *136*, 4780. i) Feng, C.; Loh, T. P. *Angew. Chem. Int. Ed.* **2014**, *53*, 2722. j) Feng, C.; Feng, D.; Luo, Y.; Loh, T. P. *Org. Lett.* **2014**, *16*, 5956.
- For nickel-catalyzed  $\gamma$ -C(sp<sup>2</sup>)–H alkynylations, see: k) Liu, Y.-J.; Liu, Y.-H.; Yana, S.-Y.; Shi, B.-F. *Chem. Commun.* **2015**, *51*, 6388.
- For copper-catalyzed  $\gamma$ -C(sp<sup>2</sup>)–H alkynylations, see: l) Liu, Y.-J.; Liu, Y.-H.; Yin, X.-S.; Gu, W.-J.; Shi, B.-F. *Chem. Eur. J.* **2015**, *21*, 205.
- Guan, M.; Chen, C.; Zhang, J.; Zeng, R.; Zhao, Y. *Chem. Commun.* **2015**, *51*, 12103.
- C(sp<sup>3</sup>)–H functionalizations via a six-membered Pd-Cycle: a) Subba Reddy, B. V.; Rajender Reddy, L.; Corey, E. J. *Org. Lett.* **2006**, *8*, 3391. b) He, G.; Zhao, Y.; Zhang, S.; Lu, C.; Chen, G. *J. Am. Chem. Soc.* **2012**, *134*, 3. c) Nadres, E. T.; Daugulis, O. *J. Am. Chem. Soc.* **2012**, *134*, 7. d) Zhang, S.-Y.; He, G.; Nack, W.A.; Zhao, Y.; Li, Q.; Chen, G. *J. Am. Chem. Soc.* **2013**, *135*, 2124. e) Cui, W.; Chen, S.; Wu, J.-Q.; Zhao, X.; Hu, W.; Wang, H. *Org. Lett.* **2014**, *16*, 4288. f) Xu, J.-W.; Zhang, Z.-Z.; Rao, W.-H.; Shi, B.-F. *J. Am. Chem. Soc.* **2016**, *138*, 10750.
- C(sp<sup>2</sup>)–H functionalizations via a six-membered Pd-Cycle: a) Li, J.-J.; Mei, T.-S.; Yu, J.-Q. *Angew. Chem. Int. Ed.* **2008**, *47*, 6452. b) Jordan-Hore, J. A.; Johansson, C.; Gulias, M.; Beck, E. M.; Gaunt, M. J. *J. Am. Chem. Soc.* **2008**, *130*, 16184. c) Han, J.; Liu, P.; Wang, C.; Zhao, Y.-S. *Org. Lett.* **2014**, *16*, 5682. d) Wang, C.; Chen, C.-P.; Zhang, J.-Y.; Yao, Y.-M.; Zhao, Y.-S. *Angew. Chem. Int. Ed.* **2014**, *53*, 9884.
- a) Openshaw, H.T.; Whittaker, N. *J. Chem. Soc. C* **1969**, 89. b) Wang, L.-H.; Zipse, H. *Liebigs Ann. Chem.* **1996**, 1501.
- Baget, J.; Llona-Minguez, S.; Lang, S.; MacKay, S. P.; Suckling, C. J.; Sutcliffe, O. B. *Beilstein J. Org. Chem.* **2011**, *7*, 1407.

1 17) Morimoto, H.; Fujiwara, R.; Shimizu, Y.; Morisaki, K.;  
2 Ohshima, T. *Org. Lett.* **2014**, *16*, 2018.

3 18) Feng, Y.; Wang, Y.; Landgraf, B.; Liu, S.; Chen, G. *Org. Lett.*  
4 **2010**, *12*, 3414 and references therein.

5 19) Most H's removed for clarity. Atoms are shown at the 50%  
6 probability level.

7 20) While other bromoacetylene derivatives were surveyed, only  
8 the use of (bromoethynyl)triisopropylsilane resulted in productive  
9 reactivity, consistent with the observations of Chen and  
10 Chatani.(Refs. 11b,11c). Activation/cross-coupling mediated by other  
11 metal complexes may be more successful and are currently being  
12 investigated.

21) V. O. Rodionov; S. I. Presolski; S. Gardinier; Y.-H. Lim; M. G.  
Finn; *J. Am. Chem. Soc.* **2007**, *129*, 12696–12704.

22) D. B. Grotjahn and D. A. Lev; *J. Am. Chem. Soc.* 2004, *126*,  
12232–12233.

23) Migratory insertion of the bromo-alkyne unit followed by  $\beta$ -Br  
elimination, as proposed by Chatani (Refs. 11a, b), cannot be dis-  
counted.

## TOC GRAPHIC:

