


This article was downloaded by: [Brown University]

On: 15 January 2013, At: 19:05

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides, Nucleotides and Nucleic Acids

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/Incn20>

Perylene Diimide-Oligonucleotide Conjugates Constructed by Click Chemistry

Alexey V. Ustinov^a, Veronika V. Dubnyakova^a & Vladimir A. Korshun^a

^a Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, Moscow, Russia

Version of record first published: 10 Dec 2007.

To cite this article: Alexey V. Ustinov, Veronika V. Dubnyakova & Vladimir A. Korshun (2007): Perylene Diimide-Oligonucleotide Conjugates Constructed by Click Chemistry, *Nucleosides, Nucleotides and Nucleic Acids*, 26:6-7, 751-754

To link to this article: <http://dx.doi.org/10.1080/15257770701490936>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

PERYLENE DIIMIDE–OLIGONUCLEOTIDE CONJUGATES CONSTRUCTED BY CLICK CHEMISTRY

Alexey V. Ustinov, Veronika V. Dubnyakova, and Vladimir A. Korshun □

*Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences,
Moscow, Russia*

□ *Oligonucleotide conjugates were prepared by copper-catalyzed cycloaddition of acetylene-modified oligonucleotides to diazido derivative of perylene 3,4,9,10-tetracarboxylic acid diimide.*

Keywords Oligonucleotides; conjugates; click chemistry; perylene diimide; [3+2] cycloaddition

The copper-catalyzed cycloaddition reaction of azides to alkynes^[1,2] (Scheme 1) is a remarkable representative of so-called “click-chemistry”^[3] reactions that are useful for bioconjugate synthesis. The process is being carried out regioselectively in mild conditions, tolerated by a majority of functional groups, affords great opportunities for the preparation of modified oligonucleotides,^[4,5] and oligonucleotide immobilization.^[6]


Conjugates of oligonucleotides with photostable perylene diimide dyes were extensively studied because of their interesting ability to self-assembly.^[7–9] The perylene diimide moiety located in the middle of an oligonucleotide strand is known to increase its propensity to triplex formation.^[10,11]

Although phosphoramidite reagents^[9,10] were used to introduce perylene diimide into the strand, the preparation of conjugates with two strands of the same polarity flanking a single dye is not straightforward. Herein we would like to report a click chemistry approach to these products.


An introduction of reactive group into an oligonucleotide is required for conjugation to take place. Since the azido group is incompatible with

This research was supported by grants from the Russian Foundation for Basic Research (projects 06-04-81019 and 06-03-32426). We thank Vadim V. Shmanai, Sergey V. Gontarev, and Maksim V. Kvach for oligonucleotide synthesis; Andrei D. Malakhov, Igor A. Prokhorenko, and Mikhail V. Skorobogatyi for helpful discussions; and Larisa V. Grusintseva for technical assistance.

Address correspondence to Alexey V. Ustinov, Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, Miklukho-Maklaya 16/10, 117997 Moscow, Russia. E-mail: austinov@mail.ibch.ru



SCHEME 1 Copper-catalyzed cycloaddition of azides to alkynes.




Reagents and conditions: *i*) Propargylamine, MeOH, 60°C, 48 h; 4,4'-dimethoxytrityl chloride, Py, 0°C to rt, 12 h, 86%; *ii*) 2-cyanoethoxy-bisdiisopropylaminophosphine, diisopropylammonium tetrazole, rt, 12 h, 93%

SCHEME 2 Synthesis of acetylenic oligonucleotides

phosphoramidite synthesis because of Staudinger reaction,^[12] we decided to introduce a terminal alkyne using a dihydroxybutyramide^[13] reagent **1** prepared from propargylamine and (*R*)-2-hydroxybutyrolactone (Scheme 2).

Oligonucleotides (Table 1) were prepared from reagent **1** and solid support **2** by means of standard oligonucleotide synthesis.

Perylene diimide-derived diazide **3** prepared from perylenetetracarboxylic acid dianhydride in two steps (Scheme 3) was then coupled to oligonucleotides to give corresponding conjugates (Table 2). The reaction



Reagents and conditions: *i*) 2-(2-hydroxyethoxy)ethylamine, 130°C, 3 h, 94%; *ii*) (PhO)₂PON₃, Ph₃P, DEAD, THF, rt, 12 h, 98%.

SCHEME 3 Synthesis of perylene diimide-oligonucleotide conjugates.

TABLE 1 Oligonucleotides containing terminal alkyne (**X** = residue of **1**)

#	Sequence, 5' → 3'
ON1	X CA-TTA-CAT-CCA-GAC
ON2	X GT-CTG-GAT-GTA-ATG

TABLE 2 Yields of conjugates of acetylenic oligonucleotides with diazide **3**

Starting oligonucleotide	Oligonucleotide/azide	Products ^a			
		Monoadduct		Diadduct	
		Yield	MALDI ^b	Yield	MALDI ^b
ON1	1:1	36%	5029.8/5028.5	42%	9444.7/9439.4
ON1	2:1	—	—	41%	—
ON2	1:1	35%	5171.1/5170.6	35%	9729.2/9723.6
ON2	2:1	—	—	59%	—

^a Isolated yield (PAGE), determined by UV absorbance^b MALDI-TOF mass (M+H), found/calculated

was carried out in aq. DMSO under argon. Tris[(1-benzyltriazol-4-yl)methyl]amine^[14] was used as a ligand for Cu(I) species to decrease copper oxidation accompanied by free hydroxyl radical ($\cdot\text{OH}$) formation leading to oligonucleotide cleavage. Heating (95°C to rt, overnight) was required to facilitate dissolution of azide and increase overall yield of conjugation.

The conjugates can be isolated either by PAGE or by HPLC. It is possible to prepare diadducts of oligonucleotides **ON1** and **ON2** selectively using 2 equiv. of oligonucleotide.

We are planning to study the fluorescent properties of these conjugates as well as their use for assembly of DNA-nanostructures.^[15]

REFERENCES

1. Tornøe, C.W.; Christensen, C.; Meldal, M. Peptidotriazoles on solid phase: [1,2,3]-triazoles by regioselective copper(I)-catalyzed 1,3-dipolar cycloadditions of terminal alkynes to azides. *J. Org. Chem.* **2002**, 67(9), 3057–3064.
2. Rostovtsev, V.V.; Green, L.G.; Fokin, V.V.; Sharpless, K.B. A stepwise Huisgen cycloaddition process: Copper(I)-catalyzed regioselective ligation of azides and terminal alkynes. *Angew. Chem. Int. Ed.* **2002**, 41(14), 2596–2599.
3. Kolb, H.C.; Finn, M.G.; Sharpless, K.B. Click chemistry: Diverse chemical function from a few good reactions. *Angew. Chem. Int. Ed.* **2001**, 40(11), 2004–2021.
4. Gartner, Z.J.; Grubina, R.; Calderone, C.T.; Liu, D.R. Two enabling architectures for DNA-templated organic synthesis. *Angew. Chem. Int. Ed.* **2003**, 42(12), 1370–1375.
5. Ustinov, A.V.; Korshun, V.A. Oligonucleotides containing arylacetylene residues: Synthesis and post-synthetic modification via [3+2] cycloaddition. *Russ. Chem. Bull.* **2006**, 55(7), 1268–1274.

6. Devaraj, N.K.; Miller, G.P.; Ebina, W.; Kakaradov, B.; Collman, J.P.; Kool, E.T.; Chidsey, C.E.D. Chemoselective covalent coupling of oligonucleotide probes to self-assembled monolayers. *J. Am. Chem. Soc.* **2005**, 127(24), 8600–8601.
7. Abdalla, M.A.; Bayer, J.; Rädler, J.O.; Müllen, K. Synthesis and self-assembly of perylenediimide-oligonucleotide conjugates. *Angew. Chem. Int. Ed.* **2004**, 43(30), 3967–3970.
8. Zheng, Y.; Long, H.; Schatz, G.C.; Lewis, F.D. Duplex and hairpin dimer structures for perylene diimide-oligonucleotide conjugates. *Chem. Comm.* **2005**, 4795–4797.
9. Wagner, C.; Wagenknecht, H.-A. Perylene-3,4:9,10-tetracarboxylic acid diimide dye as an artificial DNA base surrogate. *Org. Lett.* **2006**, 8(19), 4191–4194.
10. Bevers, S.; O'Dea, T.P.; McLaughlin, L.W. Perylene- and naphthalene-based linkers for duplex and triplex stabilization. *J. Am. Chem. Soc.* **1998**, 120(42), 11004–11005.
11. Bevers, S.; Schutte, S.; McLaughlin, L.W. Naphthalene- and perylene-based linkers for the stabilization of hairpin triplexes. *J. Am. Chem. Soc.* **2002**, 122(25), 5905–5915.
12. Bräse, S.; Gil, C.; Knepper, K.; Zimmermann, V. Organic azides: An exploding diversity of a unique class of compounds. *Angew. Chem. Int. Ed.* **2005**, 44(33), 5188–5240.
13. Dioubankova, N.N.; Malakhov, A.D.; Stetsenko, D.A.; Gait, M.J.; Korshun, V.A. Phosphoramidites and solid supports based on N-substituted 2,4-dihydroxybutyramides: universal reagents for synthesis of modified oligonucleotides. *Tetrahedron*, **2006**, 62(29), 6762–6773.
14. Chan, T.R.; Hilgraf, R.; Sharpless, K.B.; Fokin, V.V. Polytriazoles as copper(I)-stabilizing ligands in catalysis. *Org. Lett.* **2004**, 6(17), 2853–2855.
15. Detailed procedures and compound characterization will be reported elsewhere.