

Communication

Enantioselective, Catalytic Fluorolactonization Reactions with a Nucleophilic Fluoride Source

Eric Woerly, Steven M. Banik, and Eric N. Jacobsen

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.6b09499 • Publication Date (Web): 06 Oct 2016

Downloaded from http://pubs.acs.org on October 6, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Enantioselective, Catalytic Fluorolactonization Reactions with a Nucleophilic Fluoride Source

Eric M. Woerly, Steven M. Banik, Eric N. Jacobsen*

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, United States

Supporting Information Placeholder

ABSTRACT: The enantioselective synthesis of 4fluoroisochromanones via chiral aryl iodide-catalyzed fluorolactonization is reported. This methodology uses HF•pyridine as a nucleophilic fluoride source with a peracid stoichiometric oxidant, and provides access to lactones containing fluorine-bearing stereogenic centers in high enantioand diastereoselectivity. The regioselectivity observed in these lactonization reactions is complementary to that obtained with established asymmetric electrophilic fluorination protocols.

The stereocontrolled construction of C-F bonds represents a frontier endeavor in synthetic chemistry, motivated in large part by the important ways that fluorine incorporation is known to modulate the physical and biological properties of organic molecules.¹ Electrophilic fluorine sources ("F⁺") such as Selectfluor, N-fluoropyridinium salts, and Nfluorobenzenesulfonimide (NFSI) have been used extensively in the enantiocontrolled generation of fluorine-bearing stereogenic centers,² most often via the intermediacy of enolate equivalents to produce α -fluorocarbonyl compounds.^{2a,3} Fluorofunctionalization of alkenes using F⁺ sources is another powerful approach to the enantioselective synthesis of fluorine-containing chiral compounds that allows for the synthesis of highly functionalized products from simple olefin-containing starting materials.^{4,5} In reported efforts directed toward enantioselective fluorolactonization reactions, electrophilic fluorinating reagents have been shown to afford γ-butyrolactone products with exocyclic fluoromethyl substituents in moderate-to-high enantioselectivity (Scheme 1, top).⁶ We hypothesized that a hypervalent iodine catalysis of a nucleophilic fluorination pathway ("F-") could provide a complementary approach to regioisomeric fluorolactone products containing a C-F stereogenic center (Scheme 1, bottom), in a manner analogous to that observed in recently described enantioselective acetoxylactonization reactions.⁷ Herein, we report the development of an enantioselective, catalytic fluorolactonization reaction for the preparation of 4-fluoroisochromanones in high enantio- and diastereoselectivity.

Scheme 1. Enantioselective fluorolactonizations with electrophilic and nucleophilic fluorinating agents



Scheme 2. Anchimeric assistance in 1,2-difluorination of styrenes with ArI-catalysts and an analogous fluorolactonization pathway



We reported recently a protocol for the aryl iodidecatalyzed diastereoselective 1,2-difluorination of both terminal and internal alkenes^{8,9} using HF-pyridine (pyr•9HF) as a nucleophilic fluorine source and meta-chloroperbenzoic acid (mCPBA) as a stoichiometric oxidant. These conditions were first developed by Shibata and coworkers in the context of catalytic aminofluorination reactions.¹⁰ Alkenes bearing proximal weakly Lewis basic groups were shown to undergo net anti-difluorination, while substrates lacking such functionality afforded syn-difluoride products. We proposed that properly positioned weakly nucleophilic groups (e.g. the o-NO₂ group in Scheme 2) can displace the aryl iodide from intermediate I to form an unstable bridged intermediate II. Invertive displacement of the neighboring group by fluoride led to the anti-diastereomeric outcome. The propensity for such anchimeric assistance in these reactions suggested that styrenes containing other nucleophilic neighboring groups such as an ortho-carboxylic acid or ester might undergo fluorolactonization reactions.¹¹ As outlined in the proposed catalytic pathway in Scheme 2, nucleophilic displacement of the aryliodo group in intermediate I with a carboxylate equivalent would lead to formation of fluoroisochromanone products. This approach would produce a fluorine-bearing stereogenic center with a syn relationship between the two newly formed bonds,¹² a stereochemical outcome distinct from prototypical bromo- and iodolactonizations.¹³ Furthermore, as members of the polyketide-derived 4-oxyisochromanone class of natural products are known to possess interesting biological activity,¹⁴ we were motivated by the possibility that an efficient, stereocontrolled route to 4-fluoro analogs may be of interest for biological or pharmacological applications.

Methyl benzoate derivative **1a** was evaluated as a model substrate for the proposed fluorolactonization reaction. Various classes of chiral aryl iodides were examined as potential catalysts, including chiral resorcinol derivatives such as **2a**-**2d**,^{7,15} which have found application previously in a variety of enantioselective alkene oxidation reactions.¹⁶ In the presence of catalysts **2a**-**d**, *m*CPBA as oxidant, and pyr•9HF as the fluorine source and acid promoter, **1a** was observed to undergo cyclization to fluoroisochromanone **3** as a single observable diastereomer (Table 1).¹⁷ The syn relative configuration was determined via X-ray crystallographic analysis, consistent with the reaction pathway outlined in Scheme 2.¹⁸

Catalysts **2c** and **2d** were found to impart significantly higher enantioselectivities than the corresponding benzylsubstituted analogs **2a–b** (entries 1-4). This observation stands in contrast to results obtained in the migratory geminal difluorination of β -substituted styrenes reported recently by our group,¹⁹ where that trend was reversed and the polarizable benzylic groups in para-substituted analogs of **2a–b** were shown to play a critical role in enhancing ee. It is evident that subtle yet fundamentally different factors are responsible for enantioinduction in these closely related reactions. Variation of the ester group of the catalyst had very little effect on the enantioselectivity of the fluorolactonization reaction, but measurably improved yields were obtained with benzyl ester catalysts **2a** and **2d** relative to their methyl ester counterparts. On the basis of these results, **2d** was selected as the optimal catalyst for further study.

Table 1. Optimization of the fluorolactonization reaction

	OR pyr	talyst (10 mol% PBA (1.2 equiv •9HF (2.8 equiv 0.4 M CH ₂ Cl ₂ –50 °C, 24 h			Ý.
entry	substrate	R	catalyst	ee (%) ^a	yield (%) ^b
1	1a	Me	2a	87	72
2	1a	Me	2b	-87	50
3	1a	Me	2c	-94	68
4	1a	Ме	2d	95	86 (68) ^c
5	1b	<i>i</i> Pr	2d	95	72
6	1c	Bn	2d	87	42
7	1d	Н	2d	86	95 (70) ^c

Conditions: substrate (0.10 mmol), catalyst (10 mol%), *m*CPBA (0.12 mmol), pyr•9HF (2.5 mmol HF) in CH₂Cl₂ (0.25 mL) cooled to –50 °C, 24 h. ^aEnantioselectivities were determined by GC or HPLC analysis with commercial chiral columns. ^bYields were measured by GC and are based on an internal standard. ^qIsolated yield on a 1 mmol scale.



Variation of the carboxylate equivalent (1a-c) or use of the free carboxylic acid 1d resulted in formation of fluorolactonization product 3, albeit with discernible changes in ee and yield (entries 4-7) and with methyl ester 1a affording the best results.

Table 2. Fluorolactonization of aryl-substituted substrates

		O OMe Me 4a-o	2d (10 mol%) mCPBA (1.2 equiv) pyr∙9HF (2.8 equiv) 0.4 M CH ₂ Cl ₂ −50 °C, 24 h		R 5a-o F Me	
_	entry	4	R	5	ee (%) ^a	yield (%) ^b
	1	4a	6-Me	5a	96	53
	2	4b	5-Me	5b	93	50
	3	4c	4-Me	5c	94	64
	4	4d	3-Me	5d	66	52
	5	4e	5-OCF ₃	5e	89	55
	6	4f	5-Br	5f	86	48
	7	4g	5-CI	5g	86	60
	8	4h	6-F	5h	83	67
	9	4i	5-F	5i	93	57
	10	4j	4-F	5j	80	61
	11	4k	3-F	5k	30	49
	12	41	5-CF ₃	51	73	60
	13	4m	4-CF ₃	5m	76	44
	14	4n	5-CO ₂ Me	5n	70	35
	15	4o	4-CO ₂ Me	5o	58	51

Conditions: substrate (1 mmol), catalyst (10 mol%), *m*CPBA (1.2 mmol), pyr•9HF (25 mmol HF) in CH₂Cl₂ (2.5 mL) cooled to -50 °C, 24 h. ^aEnantioselectivities were determined by GC analysis with commercial chiral columns. ^bIsolated yields are reported.

Table 3. Fluorolactonization of substituted alkene substrates



Conditions: substrate (1 mmol), catalyst (10 mol%), *m*CPBA (1.2 mmol), pyr•9HF (25 mmol HF) in CH₂Cl₂ (2.5 mL) cooled to –50 °C, 24 h. Enantioselectivities were determined by HPLC or GC analysis with commercial chiral columns. Isolated yields are reported. ^aAbsolute configuration was determined by X-ray crystallographic analysis; all other products are assigned by analogy.

The effect of arene substitution in the enantioselective fluorolactonization reaction catalyzed by **2d** is illustrated in Table 2. Alkyl, halide, or trifluoromethoxy substitution at the 6, 5, and 4-positions of **1** is generally well tolerated, with the fluorolactone products obtained in 80-96% ee (entries 1-3, 5-10). However, substrates bearing more electron-deficient trifluoromethyl or carbomethoxy groups underwent reaction in reduced yields and enantioselectivities (entries 12-15). In all cases, the fluorolactone product was obtained as a single diastereomer.²⁰

The new fluorolactonization reaction was also extended successfully to a variety of β -substituted styrene derivatives (Table 3). In general, substituents of varying size and functionality have little impact on reaction enantioselectivity. The Lewis basic ether and cyano substituents in **7f** and **7g** are observed not to alter the relative stereochemical outcome of the fluorolactonization reaction, despite the propensity toward anchimeric assistance pathways in these reactions as noted above. Catalyst control over stereoselectivity of the reaction was observed with **7h** and **7i**, with complementary diastereoselectivity obtained using **2d** or ent-**2d**.

As illustrated in Scheme 3, the regioselectivity observed in the fluorolactonization reactions with a nucleophilic fluoride source is opposite to that obtained with electrophilic reagents.²¹ Furthermore, as demonstrated with **1d**, electrophilic fluorolactonizations of disubstituted styrenes are observed to be poorly diastereoselective. The successful introduction of C-F stereocenters in a highly stereocontrolled manner is thus a significant feature of the new catalytic protocol.

Scheme 3. Comparison of the reactivity of electrophilic and nucleophilic fluorinating reagents



In conclusion, the enantio- and diastereoselective, catalytic synthesis of 4-fluoroisochromanones can be accomplished with HF-pyridine as a nucleophilic fluoride source. Readily accessible chiral aryl iodides catalyze fluorolactonization with generation of C–F stereogenic centers from simple styrene precursors. Ongoing efforts are directed toward exploring the scope of fluorofunctionalization reactions induced by hypervalent iodine(III) catalysis, as well as elucidating the basis of the subtle catalyst structural properties that control the enantioselectivity in these reactions.

ASSOCIATED CONTENT

Supporting Information. Experimental procedures and characterization data for new compounds are available free of charge via the internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

jacobsen@chemistry.harvard.edu

Funding Sources

No competing financial interests have been declared.

ACKNOWLEDGMENT

This work was supported by the NIH (GM043214), by an NSF pre-doctoral fellowship to S.M.B., and by a Suzanne and Bob Wright postdoctoral fellowship to E.M.W. from the Damon Runyon Cancer Research Foundation (DRG-2180-14). We thank Dr. Shao-Liang Zheng for crystal structure determination and Jennifer Wang for mass spectrometry analysis.

REFERENCES

- (a) Gillis, E.P.; Eastman, K.J.; Hill, M.D.; Donnelly, D.J.; Meanwell, N.A. *J. Med. Chem.* 2015, *58*, 8315. (b) Wang, J.; Sanchez-Rosello, M.; Acena, J.L.; del Pozo, C.; Sorochinsky, A.E.; Fustero, S.; Soloshonok, V.A.; Liu, H. *Chem. Rev.* 2014, *114*, 2432. (c) Purser, S.; Moore, P.R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* 2008, *37*, 320. (d) Hagmann, W.K. *J. Med. Chem.* 2008, *51*, 4359.
- (2) For reviews on this topic, see: (a) Yang, X.; Tao, W.; Phipps, R.J.; Toste, F.D. *Chem. Rev.* 2015, *115*, 826. (b) Champagne, P.A.; Desroches, J.; Hamel, J.-D.; Vandamme, M.; Paquin, J.F. *Chem. Rev.* 2015, *115*, 9073.
- (3) For seminal examples utilizing organocatalysts, see: (a) Shibata, N.; Suzuki, E.; Takeuchi, Y. J. Am. Chem. Soc. 2000, 122, 10728.
 (b) Cahard, D.; Audouard, C.; Plaquevent, J.-C.; Roques, N. Org. Lett. 2000, 2, 3699. (c) Kim, D.Y.; Park, E.J. Org. Lett. 2002, 4, 545. (d) Enders, D.; Hüttl, M.R. Synlett 2005, 6, 991. (e) Marigo, M.; Fielenbach, D.; Braunton, A.; Kjærsgaard, A.; Jørgensen, K. A. Angew. Chem. Int. Ed. 2005, 44, 3703. (f) Steiner, D.D.; Mase, N.; Barbas, C.F. III. Angew. Chem. Int. Ed. 2005, 44, 3706. (g) Beeson, T.D.; MacMillan, D.W.C. J. Am. Chem. Soc. 2005, 127, 8826.
- (4) For reviews on this topic, see: (a) Wolstenhulme, J.R.; Gouverneur, V. Acc. Chem. Res. 2014, 47, 3560. (b) Denmark, S.E.; Kuester, W. E.; Burk, M.T. Angew. Chem. Int. Ed. 2012, 51, 10938. (c) Cahard, D.; Xu, X.; Couve-Bonnaire, S.; Pannecoucke, X. Chem. Soc. Rev. 2010, 39, 558.
- (5) For specific examples, see: (a) Greedy, B.; Paris, J.-M.; Vidal, T.; Gouverneur, V. Angew. Chem. Int. Ed. 2003, 42, 3291. (b) Wilkinson, S.C.; Lozano, O.; Schuler, M.; Pacheco, M.; Salmon, R.; Gouverneur, V. Angew. Chem. Int. Ed. 2009, 48, 7083. (c) Rauniyar, V.; Lackner, A.D.; Hamilton, G.L.; Toste, F.D. Science 2011, 334, 1681. (d) Wolstenhulme, J.R.; Rosenqvist, J.; Lozano, O.; Ilupeju, J.; Wurz, N.; Engle, K.M.; Pidgeon, G.W.; Moore, P.R.; Sandford, G.; Gouverneur, V. Angew. Chem. Int. Ed. 2013, 52, 9796.

- (6) (a) Parmar, D.; Maji, M.S.; Rueping, M. Chem. Eur. J. 2014, 20, 83. (b) Egami, H.; Asada, J.; Sato, K.; Hashizume, D.; Kawato, Y.; Hamashima, Y. J. Am. Chem. Soc. 2015, 137, 10132.
- (7) (a) Fujita, M.; Yoshida, Y.; 'Miyata, K.; Wakisaka, A.; Sugimura, T. Angew. Chem. Int. Ed. 2010, 49, 7068. (b) Shimogaki, M.; Fujita, M.; Sugimura, T. Eur. J. Org. Chem. 2013, 7128.
- (8) Banik, S. M.; Medley, J. W.; Jacobsen, E. N. J. Am. Chem. Soc. 2016, 138, 5000.
- (9) Gilmour and coworkers independently reported a similar protocol for the 1,2-difluorination of terminal alkenes: Molnár, I.G.; Gilmour, R. J. Am. Chem. Soc. 2016, 138, 5004.
- (10) (a) Suzuki, S.; Kamo, T.; Fukushi, K.; Hiramatsu, T.; Tokunaga, E.; Dohi, T.; Kita, Y.; Shibata, N. *Chem. Sci.* **2014**, *5*, 2754. For an additional report utilizing HF-pyridine and *m*CPBA in catalytic fluorination reactions, see: (b) Kitamura, T.; Muta, K.; Oyamada, J. *J. Org. Chem.* **2015**, *80*, 10431.
- (11) For an example of an enantioselective fluorinative intramolecular cyclization reaction of alkenes with stoichiometric, chiral iodine(III) reagents, see: Kong, W.; Feige, P.; de Haro, T.; Nevado, C. Angew. Chem. Int. Ed. 2013, 52, 2469.
- (12) Alternatively, cyclization prior to nucleophilic fluorination would also provide 4-fluoroisochromanone products with *syn*configurations. Such a mechanism cannot be ruled out based on the data presented here. For reactions where cyclization is proposed to precede fluorination, see Sawaguchi, M.; Hara, S.; Fukuhara, T.; Yoneda, N. *J. Fluorine Chem.* **2000**, *104*, 277.
- (13) Chen, J.; Zhou, L.; Tan, C.K.; Yeung, Y.-Y. *J. Org. Chem.* **2012**, *77*, 999.
- (14) (a) Aldridge, D.C.; Galt, S.; Giles, D.; Turner, W.B. *J. Chem. Soc. C* 1971, 1623. (b) Zhang, W.; Krohn, K.; Draegar, S.; Schultz, B. *J. Nat. Prod.* 2008, *71*, 1078.
- (15) Uyanik, M.; Yasui, T.; Ishihara, K. Angew. Chem. Int. Ed. 2010, 49, 2175.
- (16) For reviews on this topic, see: (a) Yoshimura, A; Zhdankin, V.V. *Chem. Rev.* 2016, *116*, 3328. (b) Romero, R.M.; Woste, T.H.; Muniz, K. *Chem. Asian J.* 2014, *9*, 972. (c) Parra, A.; Reboredo, S. *Chem. Eur. J.* 2013, *19*, 17244.
- (17) Reducing the number of equivalents of pyr•9HF resulted in lower yields of 3, albeit with consistent enantioselectivities (1.1 equiv. pyr•9HF: 93% ee and 28% yield; 5.6 equiv. pyr•9HF: 93% ee and 83% yield). Reactions conducted at -20 °C led to the generation of 3 in 86% ee and 71% yield (yields determined by GC analysis).
- (18) For examples of syn 1,2 additions to alkenes promoted by hypervalent iodine reagents, see references 7, 8, and Hara, S.; Nakahigashi, J.; Ishi-I, K.; Sawaguchi, M.; Sakai, H.; Fukuhara, T.; Yoneda, N. *Synlett* **1998**, 495.
- (19) Banik, S.M.; Medley, J.W.; Jacobsen, E.N. *Science* **2016**, *353*, 51.
- (20) The remainder of the mass balance in these reactions can be attributed to unreacted starting material and to the formation of several uncharacterized, but readily separable, byproducts. For a discussion of background reactions in closely related catalytic oxylactonization reactions, see reference 7b
- (21) For an additional example of a racemic, regioselective fluorocyclization reaction with hypervalent iodine reagents, see: Geary, G.C.; Hope, E.G.; Stuart, A.M. Angew. Chem. Int. Ed. 2015, 54, 14911.

