## Total Synthesis of (—)-Polycavernoside A: Suzuki—Miyaura Coupling Approach

## Yusuke Kasai, Takanori Ito, and Makoto Sasaki\*

*Graduate School of Life Sciences, Tohoku University, 2-1-1 Katahira, Aoba-ku, Sendai 980-8577, Japan* 

masasaki@bios.tohoku.ac.jp

## Received May 9, 2012



A total synthesis of (-)-polycavernoside A, a marine lethal toxin isolated from the edible alga *Gracilaria edulis*, has been achieved via a convergent approach. The synthesis is highlighted by catalytic asymmetric syntheses of the two key fragments and their union through Suzuki-Miyaura coupling and Keck macrolactonization.

Polycavernoside A (1, Figure 1) was isolated in 1993 by Yasumoto and co-workers, together with a minor analogue polycavernoside B, as causative toxins for the fatal human poisoning that occurred in Guam in 1991 and in the Philippines in 2002-2003, due to ingestion of the edible red alga Gracilaria edulis (Polycavernosa tsudai).<sup>1,2</sup> The  $LD_{99}$  value of 1 in mice (ip) was estimated to be 200–400  $\mu$ g/kg. The planar structure of 1 was elucidated on the basis of extensive 2D NMR studies, which showed it to be a novel glycosylated macrolide.<sup>1</sup> The complete stereochemistry, including the absolute configuration, was established by the Murai group with their first total synthesis of (-)-polycavernoside A in 1998.<sup>3</sup> Thereafter, five congeners, polycavernosides A2, A3, B2, C, and C2, were isolated as minute constituents from G. edulis and structurally determined by Yotsu-Yamashita et al.<sup>4</sup> The structure of 1 consists



Figure 1. Structures of polycavernosides A and B.

of a 16-membered macrolactone adorned with tetrasubstituted tetrahydropyran and five-membered hemiacetal rings, a conjugated triene side chain appended at C15, and a highly *O*-methylated L-fucosyl-D-xylose unit attached at C5.

The complex molecular architecture of polycavernoside A, coupled with the lethal toxicity and very limited availability from natural sources, makes this natural substance an attractive target molecule for total synthesis. While four

LETTERS 2012 Vol. 14, No. 12 3186–3189

ORGANIC

<sup>(1)</sup> Yotsu-Yamashita, M.; Haddock, R. L.; Yasumoto, T. J. Am. Chem. Soc. 1993, 115, 1147–1148.

<sup>(2)</sup> Yotsu-Yamashita, M.; Yasumoto, T.; Yamada, S.; Bajarias, F. F. A.; Formeloza, M. A.; Romero, M. L.; Fukuyo, Y. *Chem. Res. Toxicol.* **2004**, *17*, 1265–1271.

<sup>(3)</sup> Fujiwara, K.; Murai, A.; Yotsu-Yamashita, M.; Yasumoto, T. J. Am. Chem. Soc. **1998**, 120, 10770–10771.

<sup>(4) (</sup>a) Yotsu-Yamashita, M.; Seki, T.; Paul, V. J.; Naoki, H.; Yasumoto, T. *Tetrahedron Lett.* **1995**, *36*, 5563–5566. (b) Yotsu-Yamashita, M.; Abe, K.; Seki, T.; Fujiwara, K.; Yasumoto, T. *Tetrahedron Lett.* **2007**, *48*, 2255–2259.

total syntheses<sup>3,5</sup> and several synthetic approaches<sup>6</sup> have been reported to date, its detailed mechanism of action still remains unresolved due to the lack of natural sample.<sup>7</sup> Herein, we report a new convergent total synthesis of polycavernoside A based on a Suzuki–Miyaura coupling strategy.

Our retrosynthetic analysis of polycavernoside A (1) is illustrated in Scheme 1. We envisioned that the known intermediate 2, which was used in all previous total syntheses,<sup>3,5</sup> would be derived from bis-pyran 3 through oxidation of the enol ether moiety to the C9–C10 diketone functionality followed by macrolactone formation. The cyclic enol ether 3, in turn, could be constructed by Suzuki–Miyaura coupling of an alkylborane generated from the *exo*-olefin 4 and enol triflate 5.<sup>8,9</sup> These two key





(5) (a) Paquette, L. A.; Barriault, L.; Pissarnitski, D. J. Am. Chem. Soc. 1999, 121, 4542–4543. (b) Paquette, L. A.; Barriault, L.; Pissarnitski, D.; Johnston, J. N. J. Am. Chem. Soc. 2000, 122, 619–631. (c) White, J. D.; Blakemore, P. R.; Browder, C. C.; Hong, J.; Lincoln, C. M.; Nagornyy, P. A.; Robarge, L. A.; Wardrop, D. J. J. Am. Chem. Soc. 2001, 123, 8593-8595. (d) Blakemore, P. R.; Browder, C. C.; Hong, J.; Lincoln, C. M.; Nagornyy, P. A.; Robarge, L. A.; Wardrop, D. J.; White, J. D. J. Org. Chem. 2005, 70, 5449–5460. (e) Woo, S. K.; Lee, E. J. Am. Chem. Soc. 2010, 132, 4564–4565.

(6) (a) Pérez-Balado, C.; Markó, I. E. *Tetrahedron* **2006**, *62*, 2331–2349. (b) Barry, C. S.; Bushby, N.; Harding, J. R.; Willis, C. L. Org. Lett. **2005**, *7*, 2683–2686. (c) Pérez-Balado, C.; Markó, I. E. *Tetrahedron Lett.* **2005**, *46*, 4887–4890. (d) Dumeunier, R.; Markó, I. E. *Tetrahedron Lett.* **2000**, *41*, 10219–10222.

(7) (a) Cagide, E.; Louzao, M. C.; Ares, I. R.; Vieytes, M. R.; Yotsu-Yamashita, M.; Paquette, L. A.; Yasumoto, T.; Botana, L. M. *Cell. Physiol. Biochem.* **2007**, *19*, 185–194. (b) Barriault, L.; Boulet, S. L.; Fujiwara, K.; Murai, A.; Paquette, L. A.; Yotsu-Yamashita, M. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 2069–2072.

(8) For selected reviews on Suzuki-Miyaura coupling, see:
(a) Miyaura, N.; Suzuki, A. Chem. Rev. 1995, 95, 2457-2483.
(b) Chemler, S. R.; Trauner, D.; Danishefsky, S. J. Angew. Chem., Int. Ed. 2001, 40, 4544-4568. (c) Suzuki, A. Angew. Chem., Int. Ed. 2011, 50, 6722-6737.

(9) For recent reviews, see: (a) Sasaki, M.; Fuwa, H. Synlett 2004, 1851–1874.
(b) Sasaki, M. Bull. Chem. Soc. Jpn. 2007, 80, 856–871.
(c) Sasaki, M.; Fuwa, H. Nat. Prod. Rep. 2008, 25, 401–426. (d) Fuwa, H. Bull. Chem. Soc. Jpn. 2010, 83, 1401–1420.

fragments **4** and **5**, of comparable complexity, would be constructed through catalytic asymmetric reactions.

The synthesis of the C1-C8 exo-olefin 4 started with the known enone 6, which is readily available in three steps from 1,3-propanediol.<sup>10</sup> Catalytic asymmetric hetero-Diels-Alder reaction between silvloxy diene 7, derived from 6 (TMSTOf, Et<sub>3</sub>N, room temperature), and aldehyde  $\mathbf{8}^{11}$  was promoted by the Jacobsen tridentate chromium-(III) catalyst 9 (3 mol %)<sup>12</sup> and afforded the desired cvcloadduct 10 (Scheme 2). Subsequent treatment with K<sub>2</sub>CO<sub>3</sub> in MeOH produced a 6:1 mixture of ketone 11a with the desired stereochemistry at C4 and its isomer 11b, which was separated by flash column chromatography on silica gel. Isomerization of the undesired 11b was realized by using DBU (toluene, room temperature) and gave a 10:1 mixture of 11a and 11b. After two cycles of isomerization, the desired 11a was obtained in 60% overall yield from 6. Luche reduction of 11a (NaBH<sub>4</sub>, CeCl<sub>3</sub>·7H<sub>2</sub>O, MeOH, -20 °C)<sup>13</sup> delivered alcohol **12** in 94% yield as a single stereoisomer at C5. At this stage, the enantiomeric excess (96% ee) and the absolute configuration of 12 were established by <sup>1</sup>H NMR analysis of its Mosher esters.<sup>14</sup> After protection as a triisopropylsilyl (TIPS) ether, selective deprotection of the TBS group and iodination of the resultant alcohol gave iodide 13 in 86% yield (three steps). Treatment of 13 with t-BuOK furnished the exo-olefin 4, which was directly used in the next coupling reaction (see Scheme 4).

The synthesis of the C9–C16 enol triflate **5** commenced with (*R*)-(+)-citronellal (97.2% ee), which was converted into alcohol **14** via a six-step sequence<sup>15</sup> (Scheme 3). Ozonolysis of the double bond gave aldehyde **15** in 92% yield. Catalytic asymmetric allylation of **15** with prenyl bromide was carried out using the chiral sulfonamide ligand **16** and following the procedure of Kishi and co-workers;<sup>16</sup> it provided alcohol **17** (31%) and the corresponding diol (69%) as single stereoisomers, respectively. Reprotection of the primary alcohol of the latter as its TBS ether afforded **17** in 86% yield. Asymmetric dihydroxylation of the terminal alkene using (DHQ)<sub>2</sub>PYR as a chiral ligand<sup>17</sup> led to triol **18** in 97% yield. Selective protection of the primary alcohol as its *tert*-butyldiphenylsilyl (TBDPS) ether, removal of the TBS group under acidic conditions

(16) Kurosu, M.; Lin, M.-H.; Kishi, Y. J. Am. Chem. Soc. 2004, 126, 12248–12249.

(17) (a) Crispino, G. A.; Jeong, K.-S.; Kolb, H. C.; Wang, Z.-M.; Xu, D.; Sharpless, K. B. J. Org. Chem. 1993, 58, 3785–3786. (b) Kolb, H. C.; VanNieuwenhze, M. S.; Sharpless, K. B. Chem. Rev. 1994, 94, 2483–2547.

<sup>(10)</sup> Robertson, J.; Dallimore, J. W. P.; Meo, P. Org. Lett. 2004, 6, 3857–3859.

<sup>(11)</sup> Keck, G. E.; Wager, C. A.; Wager, T. T.; Savin, K. A.; Covel, J. A.; McLaws, M. D.; Krishnamurthy, D.; Cee, V. J. *Angew. Chem., Int. Ed.* **2001**, *40*, 231–234.

<sup>(12)</sup> Dossetter, A. G.; Jamison, T. F.; Jacobsen, E. N. Angew. Chem., Int. Ed. 1999, 38, 2398–2400.

<sup>(13) (</sup>a) Luche, J. L. J. Am. Chem. Soc. 1978, 100, 2226–2227.
(b) Gemal, A. L.; Luche, J. L. J. Am. Chem. Soc. 1981, 103, 5454–5459.

<sup>(14) (</sup>a) Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. J. Am. Chem. Soc. **1991**, 113, 4092–4096. (b) Hoye, T. R.; Jeffrey, C. S.; Shao, F.

*Nat. Protoc.* **2007**, *2*, 2451–2458. See the Supporting Information for details. (15) Lee, E.; Lee, Y. R.; Moon, B.; Kwon, O.; Shim, M. S.; Yun, J. K.

<sup>(15)</sup> Lee, L., Lee, T. K., Moon, B., Kwon, O., Shini, W. S., Tun, J. K. J. Org. Chem. **1994**, *59*, 1444–1456.





(CSA, MeOH), and oxidative lactonization of the resultant 1,5-diol with TEMPO and  $PhI(OAc)_2^{18}$  delivered lactone **19** in 76% yield for the three steps. The residual secondary alcohol was then protected as its triethylsilyl (TES) ether to give **20** in quantitative yield which, upon treatment with KHMDS and PhNTf<sub>2</sub> (HMPA, THF, -78 °C), furnished the desired enol triflate **5**.

With the requisite fragments in hand, we next turned our attention to their union through Suzuki–Miyaura coupling.<sup>8,9</sup> After extensive experimentation, Suzuki– Miyaura coupling of the alkylborane generated from *exo*-olefin **4** (9-BBN-H, THF, 0 °C to room temperature) with enol triflate **5** was achieved under the conditions which used aqueous  $Cs_2CO_3$  as a base, PdCl<sub>2</sub>(dppf)· CH<sub>2</sub>Cl<sub>2</sub> as a catalyst, and Ph<sub>3</sub>As as a coligand in DMF/ THF at room temperature, affording the desired coupled product **3** in high yield.<sup>19</sup> The cyclic enol ether moiety of **3** was then oxidized with *m*-chloroperoxybenzoic acid (*m*-CPBA) in CH<sub>2</sub>Cl<sub>2</sub>/MeOH to afford an inconsequential epimeric mixture of alcohols **21a** and **21b** in 66 and 25% overall yields from **13**, respectively. These alcohols were Scheme 3. Synthesis of the C9-C16 Enol Triflate 5



individually oxidized to the same ketone **22** in high yields.<sup>20</sup> Selective cleavage of the TES ether and oxidative removal of the PMB group afforded diol **23** in 98% yield for the two steps. The primary alcohol of diol **23** was selectively oxidized with TEMPO/PhI(OAc)<sub>2</sub> in CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O (10:1)<sup>21</sup> to provide the corresponding carboxylic acid in 97% yield. Subsequent macrolactonization proved to be more difficult than expected. After extensive experimentation, it was finally found that the best result was obtained by using Keck protocol<sup>22</sup> employing DCC, pyridine, and PPTS<sup>23</sup> under reflux to afford macrolactone **24** in quantitative yield.<sup>24</sup> The primary TBDPS ether was selectively cleaved with buffered HF·pyridine, and oxidation of the resulting alcohol with Dess–Martin periodinane<sup>25</sup> followed by

(21) (a) van den Bos, L. J.; Codée, J. D. C.; van der Toorn, J. C.; Boltje, T. J.; van Boom, J. H.; Overkleeft, H. S.; van der Marel, G. A. *Org. Lett.* **2004**, *6*, 2165–2168. (b) Epp, J. B.; Widlanski, T. S. *J. Org. Chem.* **1999**, *64*, 293–295.

<sup>(18)</sup> Hansen, T. M.; Florence, G. J.; Lugo-Mas, P.; Chen, J.; Abrams, J. N.; Forsyth, C. J. *Tetrahedron Lett.* **2003**, *44*, 57–59.

<sup>(19)</sup> Suzuki–Miyaura coupling of the phosphate counterpart resulted in a low yield of 3  $[Pd(PPh_3)_4, PdCl_2(dppf) \cdot CH_2Cl_2, or PdCl_2-(dppf) \cdot CH_2Cl_2/Ph_3As]$  presumably due to the low reactivity.

<sup>(20)</sup>  $10\beta$ -Alcohol **21a** was labile even under mild acidic conditions. For example, Swern oxidation of **21a** caused partial cleavage of the methyl acetal and resulted in a low yield of ketone **22** (37%).

<sup>(22)</sup> Boden, E. P.; Keck, G. E. J. Org. Chem. 1985, 50, 2394-2395.

<sup>(23)</sup> Kageyama, M.; Tamura, T.; Nantz, M. H.; Roberts, J. C.; Somfai, P.; Whritenour, D. C.; Masamune, S. J. Am. Chem. Soc. 1990, 112, 7407–7408.

<sup>(24)</sup> Fujiwara et al. reported that Yamaguchi lactonization of a substrate closely related to 23 proceeded smoothly to give the corresponding 16-membered macrolactone in good yield. However, macrolactonization of a hydroxy acid derived from 23 under Yamaguchi or Shiina conditions resulted in only a low yield of 24.

<sup>(25) (</sup>a) Dess, D. B.; Martin, J. C. J. Org. Chem. 1983, 48, 4155–4156.
(b) Dess, D. B.; Martin, J. C. J. Am. Chem. Soc. 1991, 113, 7277–7287.

Scheme 4. Fragment Coupling and Completion of the Synthesis of Polycavernoside A



Takai reaction (CHI<sub>3</sub>, CrCl<sub>2</sub>)<sup>26</sup> led to (*E*)-vinyliodide **25** in 61% yield (three steps). Treatment of **25** with TFA/THF/ $H_2O$  (1:2:2)<sup>3</sup> effected deprotection of the TIPS ether and methyl acetal with concomitant reconstruction of the five-membered hemiacetal to afford alcohol **2** in 98% yield.

Finally, glycosylation of **2** with thioglycoside  $26^{5c,d}$  under the influence of NBS<sup>27</sup> (41%), oxidative cleavage of the benzyl ether with DDQ (68%), and Stille coupling with the known dienylstannane  $27^{5a,b}$  (45%) completed the synthesis of (–)-polycavernoside A (1).

In summary, an efficient total synthesis of polycavernoside A has been accomplished via a convergent approach (a longest linear sequence of 29 steps and 2.4% overall yield from commercially available (R)-(+)-citoronellal). The synthesis features catalytic asymmetric syntheses of the C1–C8 and C9–C16 fragments and their assembly through Suzuki–Miyaura coupling and Keck macrolactonization to form the macrolactone core. Our convergent fragment coupling strategy allows for the preparation of sufficient quantities of polycavernoside A itself and its analogues for detailed biological studies. Further studies along this line are underway and will be reported in due course.

Acknowledgment. We thank Takasago International Co. for providing us (R)-(+)-citronellal. This work was supported by Grants-in-Aid from Scientific Research (A) (No. 21241050) and Scientific Research (C) (No. 22603001) from Japan Society for the Promotion of Sciences (JSPS) and the Tohoku University Global COE program "International Center of Research & Education for Molecular Complex Chemistry".

**Supporting Information Available.** Experimental procedures, characterization data, and copies of <sup>1</sup>H and <sup>13</sup>C NMR spectra for all new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

<sup>(26)</sup> Takai, K.; Nitta, K.; Utimoto, K. J. Am. Chem. Soc. 1986, 108, 7408–7410.

<sup>(27)</sup> Nicolaou, K. C.; Seitz, S. P.; Papahatjis, D. P. J. Am. Chem. Soc. **1983**, 105, 2430–2434.

The authors declare no competing financial interest.