

Enantioselective Dichlorination of Allylic Alcohols

K. C. Nicolaou,* Nicholas L. Simmons, Yongcheng Ying, Philipp M. Heretsch, and Jason S. Chen

Department of Chemistry and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, California 92037, United States

Department of Chemistry and Biochemistry, University of California—San Diego, La Jolla, California 92093, United States

S Supporting Information

ABSTRACT: The development of an enantioselective allylic alcohol dichlorination catalyzed by dimeric cinchona alkaloid derivatives and employing aryl iododichlorides as chlorine sources is reported. Reaction optimization, exploration of the substrate scope, and a model for stereoinduction are presented.

Despite tremendous advances in the development of chiral methods, asymmetric olefin dichlorination¹ remains a challenging problem. Such a reaction would be useful for the synthesis of oligo- and polychlorinated compounds.² Originally discovered from the membranes of freshwater algae,³ chlorosulfolipids have gained significant attention since 2001⁴ due to their putative link to diarrhetic shellfish poisoning. Their biosynthesis likely proceeds through a series of site-selective and stereoselective enzymatic chlorinations of unfunctionalized positions on sulfolipid precursors.⁵ In contrast, synthetic chemists generally can control chlorination only at functionalized sites. Evaluation of existing synthetic approaches¹ and the structure of chlorosulfolipid cytotoxin **1** (the most complex member of the family; see Figure 1)⁶ suggested a need for asymmetric olefin dichlorination methods, especially those applicable to allylic alcohols. Most recent approaches have relied on substrate control of stereochemistry and required substrate derivatization (e.g., epoxidation of the olefin^{1e–g} or esterification of an allylic alcohol^{1h}). In the Snyder group's total synthesis of napyradiomycin A1,^{1c} there is an isolated example of a practical enantioselective olefin dichlorination employing a stoichiometric chiral auxiliary. Although the above methods have been employed in total syntheses of several chlorosulfolipids,⁷ there remains a need for additional asymmetric dichlorination methods. We present herein the development of an enantioselective dichlorination of allylic alcohols.

As shown in Scheme 1, olefin dichlorination is a challenging reaction to render enantioselective. The reaction proceeds through an initial electrophilic chlorination (see **2**) to form chloronium species **3**. Even if this process is rendered facial-selective,⁸ there remains a regioselectivity challenge in the subsequent nucleophilic chlorination; attack of the two chloronium positions of homochiral species **3** leads to opposite antipodes of **4**. Additionally, the configurational stability of chloronium species **3** may be degraded by reversibility and/or direct chloronium transfer to another molecule of the olefinic substrate (**2**).⁹

In view of these potential challenges, we selected *trans*-cinnamyl alcohol (**5**, Table 1) as a model substrate. The benzylic nature of the intermediate chloronium species would enforce a





Figure 1. Molecular structure of chlorosulfolipid cytotoxin **1** and opportunities for asymmetric allylic alcohol dichlorination.

Scheme 1. Mechanism of Olefin Dichlorination and Challenges for Enantioselectivity




regiocontrolled chloride attack. The hydroxyl moiety might hydrogen bond with a catalyst or reagent, rigidifying the system and potentially improving stereocontrol. Our first clue suggesting a direction for catalyst screening came from observations that electrophilic halogenation is dramatically accelerated by tertiary amines.¹⁰ Screening of common amine catalysts (e.g., proline and imidazolidinone derivatives,¹¹ small peptides,¹² and cinchona alkaloids^{1a,8,13}) and a range of chlorine sources (e.g., Cl_2 gas, Et_4NCl_3 , and PhICl_2) revealed the dimeric cinchona alkaloid derivative $(\text{DHQ})_2\text{PHAL}$ (commonly employed as a ligand for Sharpless asymmetric dihydroxylation) and PhICl_2 as a uniquely promising reagent combination for further studies.

Interestingly, the quality of PhICl_2 proved to be critical, with the use of PhICl_2 generated from PhI and NaOCl (Chlorox bleach or a solution from Sigma–Aldrich) resulting in poor reproducibility. However, employing PhICl_2 freshly prepared from PhI and Cl_2 gas yielded consistent results. Furthermore, catalyst aging in the presence of PhICl_2 degraded both reactivity and selectivity.

Received: March 23, 2011

Published: May 04, 2011

Table 1. Screening of Reaction Conditions for Enantioselective Dichlorination^a

entry	ArICl ₂	solvent	temp (°C)	ee (%) ^b
1	PhICl ₂	CH ₂ Cl ₂	25	22
2	PhICl ₂	CH ₂ Cl ₂	0	23
3	PhICl ₂	CH ₂ Cl ₂	−40	41
4	PhICl ₂	CH ₂ Cl ₂	−78	82
5	PhICl ₂	THF	−78	<5
6	PhICl ₂	EtOAc	−78	10
7	PhICl ₂	Et ₂ O	−78	<5
8	PhICl ₂	CH ₂ Cl ₂ :hexanes (1:1)	−78	75
9	PhICl ₂	CH ₂ Cl ₂ :PhMe (1:1)	−78	75
10	<i>o</i> -Me(C ₆ H ₄)ICl ₂	CH ₂ Cl ₂	−78	56
11	<i>p</i> -Ph(C ₆ H ₄)ICl ₂	CH ₂ Cl ₂	−78	85
12	<i>p</i> - <i>t</i> -Bu(C ₆ H ₄)ICl ₂	CH ₂ Cl ₂	−78	73

^a Reactions were performed on 7 mg (50 μmol) scale and run to completion (TLC analysis). ^b Determined by chiral HPLC analysis.

Therefore, since the uncatalyzed reaction is very slow, the catalyst was added last to a reaction mixture already containing the substrate and PhICl₂. An initial addition of 10 mol % catalyst followed by slow addition of another 10 mol % catalyst gave the best results.

Under these conditions, (DHQD)₂PHAL-catalyzed dichlorination of *trans*-cinnamyl alcohol (**5**) by PhICl₂ at ambient temperature (Table 1, entry 1) afforded dichloride **6** in 22% ee. Lowering the temperature (entries 2–4) prolonged reaction times but improved enantioselectivity. A solvent screen (entries 4–7) revealed CH₂Cl₂ to be uniquely suitable. Mixed solvents containing CH₂Cl₂ could also be used (entries 8 and 9), but the reactions became more sluggish (likely due to reduced solubility of the catalyst and PhICl₂), and selectivities were not improved. An exploration of alternative aryl iododichlorides (entries 10–12) revealed some effects on the level of enantioselectivity, with the use of *p*-Ph(C₆H₄)ICl₂ (entry 11) delivering dichloride **6** in 85% ee.

Scaling up the reaction conditions in entry 11 of Table 1 and reducing the amount of aryl iododichloride to 1.6 equiv gave dichloride **6** in 63% yield and 81% ee (see entry 1a, Table 2). Under otherwise identical conditions, but employing the pseudoenantiomeric catalyst (DHQD)₂PHAL, dichloride *ent*-**6** was obtained in 58% yield and 61% ee. As shown in Table 2, the reaction was tolerant of some changes in electronics (entries 1a–e). Very electron-deficient olefins failed to react. Electron-rich cinnamyl substrates underwent rapid dichlorination, but the products were prone to epimerization, chloride elimination, and other decomposition reactions, presumably due to facile benzylic S_N1-type reactions. This is not a limitation of the method but of compound stability; racemic reference samples were similarly labile. Naphthyl substrates reacted in the same

Table 2. Generality and Scope of Enantioselective Dichlorination^a

entry	substrate	product	yield (%) ^b	ee (%) ^c
1a			63	81
1b			65	44
1c ^d	1a: X = H (5) 1b: X = Me 1c: X = CF ₃	1a: X = H (6) 1b: X = Me 1c: X = CF ₃	75	48
1d	1d: X = Cl	1d: X = Cl (7)	81	71
1e	1e: X = F	1e: X = F	73	72
2			84	74
3			66	47
4			63	68
5 ^e			90	43
6			35	25 ^f
7			32 ^g	<5 ^g
8 ^e			48	43 ^h
9 ^e			59	54

^a Reactions were performed on 40–50 mg scale using 20 mol % of (DHQD)₂PHAL and 1.6 equiv of *p*-Ph(C₆H₄)ICl₂ in CH₂Cl₂ (0.05 M) at −78 °C and run to completion (TLC analysis). ^b Isolated yield after flash column chromatography. ^c Determined by chiral HPLC analysis. ^d Reaction performed at −40 °C. ^e PhICl₂ used in place of *p*-Ph(C₆H₄)ICl₂. ^f Absolute configuration not determined. ^g Incomplete reaction. Yield and ee determined after desilylation. ^h Determined by NMR analysis of the corresponding Mosher ester.

manner (entries 2 and 3). Steric congestion was tolerated both on the aryl ring (entry 4) and near the allylic alcohol (entry 5). Importantly, the reaction is stereospecific. Thus, as shown in entry 6, although the efficiency and enantioselectivity of the reaction of *cis*-cinnamyl alcohol (**8**) left much to be desired, it proceeded to give the opposite diastereoisomer (**9**) as compared with the dichlorination of the *trans* isomer (contrast with entry 1a). Furthermore, the allylic alcohol proved to be a critical substrate feature; masking of this moiety as a TES ether (entry 7) abolished enantioselectivity.

We also investigated the suitability of our reaction conditions on selected non-cinnamyl substrates in order to ensure that we were developing a generally useful catalytic system. We decided to investigate the reactions of monobenzylylated *cis*- and *trans*-butenediol since differentially protected hydroxyl moieties might be useful for further elaboration of the products.¹⁴ The use of *p*-Ph(C₆H₄)ICl₂ resulted in impractically long reaction times




Figure 2. Molecular structures and ORTEPs of *p*-nitrobenzoate esters **7'** and **ent-10'**.

at $-78\text{ }^{\circ}\text{C}$, so the more reactive oxidant PhICl_2 was employed instead. Pleasingly, as shown in entries 8 and 9 of Table 2, the reactions proceeded with efficiency and enantioselectivity comparable to those of cinnamyl substrates.

Enantiopure dichlorination products **7** (entry 1d, Table 2) and **ent-10** (see entry 9, Table 2) (obtained by preparative chiral HPLC) were converted into *p*-nitrobenzoate esters **7'** and **ent-10'** (Figure 2). X-ray crystallographic analysis of the *p*-nitrobenzoates allowed assignment of their absolute configurations. The other dichlorination products of *trans*-cinnamyl alcohols were presumed to have the same absolute configuration as **7**; however, we currently have no experimental proof of these assignments.

Since the dichlorination is stereospecific, it likely proceeds through the intermediacy of a chloronium species in the manner outlined in Scheme 1. Furthermore, the relative configuration of the products eliminates the possibility of anchimeric assistance in chloronium attack (see **3** $\rightarrow$ **4**, Scheme 1). The sense of absolute stereoselection is consistent with preferential chloronium formation on the face of the olefin that would react in a Sharpless asymmetric dihydroxylation employing $(\text{DHQ})_2\text{PHAL}$ as a ligand. We propose a chloronium ion transfer from an electrophilic chlorinating reagent generated through attack of the aryl iododichloride by one of the quinuclidine nitrogens of the catalyst (see **11**, Figure 3).^{10,15} Chloronium delivery might then proceed in a manner analogous to that proposed by Corey and Noe¹⁶ for the Sharpless asymmetric dihydroxylation. Since an unmasked allylic alcohol is required, we postulate the presence of a hydrogen bond to one of the phthalazine nitrogens of the catalyst (see **11**, Figure 3). Consistent with this hypothesis, $(\text{DHQ})_2\text{AQN}$ (Figure 3), lacking the phthalazine nitrogens, provides minimal stereoselection (10% ee in favor of **ent-6**, compare with 85% ee in favor of **6**; entry 11, Table 1). We acknowledge that the




Figure 3. Proposed stereoselection model (**11**) and molecular structure of $(\text{DHQ})_2\text{AQN}$.

outlined model is speculative; ongoing studies will test and refine this working model.

In conclusion, we have developed an enantioselective dichlorination of allylic alcohols employing the dimeric cinchona alkaloid derivative $(\text{DHQ})_2\text{PHAL}$ [or its pseudoenantiomer $(\text{DHQD})_2\text{PHAL}$] and aryl iododichlorides. Further screening of catalyst and iododichloride modifications is under way and promises to lead to improved enantioselectivity and generality.

■ ASSOCIATED CONTENT

S Supporting Information. Experimental procedures and characterization data for key compounds (CIF, PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

kcn@scripps.edu

■ ACKNOWLEDGMENT

We thank Drs. D. H. Huang, L. Pasternack, and R. Chadha for NMR spectroscopic and X-ray crystallographic assistance. This work was supported by the National Institutes of Health (R01ES013314), the Skaggs Institute for Research, the National Science Foundation (predoctoral fellowship to N.L.S.), and the Deutsche Akademie der Naturforscher Leopoldina (postdoctoral fellowship to P.M.H.).

■ REFERENCES

- (1) Enantioselective approaches: (a) Julia, S.; Ginebreda, A. *Tetrahedron Lett.* **1979**, *20*, 2171. (b) Adam, W.; Mock-Knoblach, C.; Saha-Möller, C. R.; Herderich, M. *J. Am. Chem. Soc.* **2000**, *122*, 9685. (c) Snyder, S. A.; Tang, Z.-Y.; Gupta, R. *J. Am. Chem. Soc.* **2009**, *131*, 5744. (d) Snyder, S. A.; Treitler, D. S.; Brucks, A. P. *J. Am. Chem. Soc.* **2010**, *132*, 14303. Diastereoselective approaches: (e) Iranpoor, N.; Firouzabadi, H.; Azadi, R.; Ebrahimzadeh, F. *Can. J. Chem.* **2006**, *84*, 69. (f) Yoshimitsu, T.; Fukumoto, N.; Tanaka, T. *J. Org. Chem.* **2009**, *74*, 696. (g) Denton, R. M.; Tang, X.; Przeslak, A. *Org. Lett.* **2010**, *12*, 4678. Diastereoselective approach: (h) Shibuya, G. M.; Kanady, J. S.; Vanderwal, C. D. *J. Am. Chem. Soc.* **2008**, *130*, 12514.

- (2) (a) Gribble, G. W. *Acc. Chem. Res.* **1998**, *31*, 141. (b) Kladi, M.; Vagias, C.; Roussis, V. *Phytochem. Rev.* **2004**, *3*, 337.
- (3) (a) Elovson, J.; Vagelos, P. R. *Proc. Natl. Acad. Sci. U.S.A.* **1969**, *62*, 957. (b) Elovson, P. R.; Vagelos, P. R. *Biochemistry* **1970**, *9*, 3110. (c) Haines, T. H.; Pousada, M.; Stern, B.; Mayers, G. L. *Biochem. J.* **1969**, *113*, 565.
- (4) Ciminiello, P.; Fattorusso, E.; Forino, M.; Di Rosa, M.; Ianaro, A.; Poletti, R. *J. Org. Chem.* **2001**, *66*, 578.
- (5) Mooney, C. L.; Haines, T. H. *Biochemistry* **1973**, *12*, 4469.
- (6) Ciminiello, P.; Dell'Aversano, C.; Fattorusso, E.; Forino, M.; Magno, S.; Di Rosa, M.; Ianaro, A.; Poletti, R. *J. Am. Chem. Soc.* **2002**, *124*, 13114.
- (7) (a) Nilewski, C.; Geisser, R. W.; Carreira, E. M. *Nature* **2009**, *457*, 573. (b) Bedke, D. K.; Shibuya, G. M.; Pereira, A.; Gerwick, W. H.; Haines, T. H.; Vanderwal, C. D. *J. Am. Chem. Soc.* **2009**, *131*, 7570. (c) Bedke, D. K.; Shibuya, G. M.; Pereira, A. R.; Gerwick, W. H.; Vanderwal, C. D. *J. Am. Chem. Soc.* **2010**, *132*, 2542. (d) Yoshimitsu, T.; Fukumoto, N.; Nakatani, R.; Kojima, N.; Tanaka, T. *J. Org. Chem.* **2010**, *75*, 5425. (e) Umezawa, T.; Shibata, M.; Kaneko, K.; Okino, T.; Matsuda, F. *Org. Lett.* **2011**, *13*, 904. (f) Yoshimitsu, T.; Nakatani, R.; Kobayashi, A.; Tanaka, T. *Org. Lett.* **2011**, *13*, 908.
- (8) (a) Whitehead, D. C.; Yousefi, R.; Jaganathan, A.; Borhan, B. *J. Am. Chem. Soc.* **2010**, *132*, 3298. (b) Yousefi, R.; Whitehead, D. C.; Mueller, J. M.; Staples, R. J.; Borhan, B. *Org. Lett.* **2011**, *13*, 608.
- (9) Denmark, S. E.; Burk, M. T.; Hoover, A. J. *J. Am. Chem. Soc.* **2010**, *132*, 1232.
- (10) Zhang, W.; Xu, H.; Xu, H.; Tang, W. *J. Am. Chem. Soc.* **2009**, *131*, 3832.
- (11) (a) Brochu, M. P.; Brown, S. P.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2004**, *126*, 4108. (b) Halland, N.; Braunton, A.; Bachmann, S.; Marigo, M.; Jørgensen, K. A. *J. Am. Chem. Soc.* **2004**, *126*, 4790. (c) Amatore, M.; Beeson, T. D.; Brown, S. P.; MacMillan, D. W. C. *Angew. Chem., Int. Ed.* **2009**, *48*, 5121.
- (12) Colby Davie, E. A.; Mennen, S. M.; Xu, Y.; Miller, S. J. *Chem. Rev.* **2007**, *107*, 5759.
- (13) (a) Tommaso, M.; Hiemstra, H. *Synthesis* **2010**, 1229. (b) Wack, H.; Taggi, A. E.; Hafez, A. M.; Drury, W. J., III; Lectka, T. *J. Am. Chem. Soc.* **2001**, *123*, 1531. (c) France, S.; Wack, H.; Taggi, A. E.; Hafez, A. M.; Wagerle, T. R.; Shah, M. H.; Dusich, C. L.; Lectka, T. *J. Am. Chem. Soc.* **2004**, *126*, 4245. (d) Bartoli, G.; Bosco, M.; Carlone, A.; Locatelli, M.; Melchiorre, P.; Sambra, L. *Angew. Chem., Int. Ed.* **2005**, *44*, 6219.
- (14) β,γ -Dichloroalcohols are amenable to Dess–Martin oxidation and elaboration of the resulting aldehyde without loss of configurational stability.^{7d}
- (15) Denmark, S. E.; Beutner, G. L. *Angew. Chem., Int. Ed.* **2008**, *47*, 1560.
- (16) (a) Corey, E. J.; Noe, M. C. *J. Am. Chem. Soc.* **1993**, *115*, 12579. (b) Corey, E. J.; Noe, M. C. *J. Am. Chem. Soc.* **1996**, *118*, 319. (c) Corey, E. J.; Noe, M. C. *J. Am. Chem. Soc.* **1996**, *118*, 11038.