🛇 Cite This: Org. Lett. XXXX, XXX, XXX–XXX

Enantioselective Aza-Reformatsky Reaction with Ketimines

Aitor Maestro, Edorta Martinez de Marigorta, Francisco Palacios,*[®] and Javier Vicario*[®]

Departamento de Química Orgánica I, Facultad de Farmacia, University of the Basque Country, UPV/EHU Paseo de la Universidad 7, 01006 Vitoria-Gasteiz, Spain

Supporting Information

ABSTRACT: Here, an enantioselective aza-Reformatsky reaction using acyclic ketimine substrates is presented. Using α -phosphorated ketimines as electrophilic substrates and a simple BINOL-derived ligand, phosphorated analogues of aspartic acid holding chiral tetrasubstituted carbons are efficiently obtained with excellent enantioselectivity through an asymmetric organocatalytic Reformatsky-type reaction.

Organic


Letters



The phosphorated analogues of aspartic acid have been used for the synthesis of phosphorus-containing enantiopure tetrasubstituted β -lactams.

eformatsky reaction is a valuable synthetic process that is Kwidely used for the effective formation of C–C bonds. This transformation consists of a nucleophilic addition, in general to aldehyde or ketone substrates, of a zinc enolate, which is generated in situ through a metal insertion into activated carbon halides.² The discovery that organozinc species can be used as a metal source³ offered more convenient homogeneous conditions for this reaction, and since then, many efforts have been made in developing catalytic systems for the enantioselective Reformatsky reaction using ketones and aldehydes.⁴ Imines are, however, suitable substrates for this reaction, providing access to β -amino acid derivatives in a simple synthetic protocol. The first catalytic enantioselective imino-Reformatsky reaction was reported by Cozzi in 2006 using in situ preformed aldimines and 20-30% of Nmethylephedrine as a chiral ligand (Scheme 1).⁵

Scheme 1. Precedents in Enantioselective Aza-Reformatsky Reaction



Although not strictly catalytic, more recently Ando et al. have reported the synthesis of α,α -difluoro- β -lactams by an imino-Reformatsky protocol promoted by 75% of aminoalcohol ligands.⁶ Pedro and Vila et al. have performed successful imino-Reformatsky processes with cyclic imines using 20% of prolinol-derived ligand.⁷ Remarkably, their catalytic system is equally effective when cyclic ketimines are used as substrates, which allows the synthesis of quaternary cyclic β -aminoesters with excellent ees (Scheme 1).

Due to the poor electrophilic character of ketimine carbons and the additional steric hindrance on the substrate, the formation of tetrasubstituted carbons from ketimines is always a challenging task. In addition, the enantiotopic faces of ketimines are not as easily discriminated as those of aldimines when asymmetric synthesis is sought.⁸ These obstacles are vitally important when acyclic substrates are used. In this context, we reported recently the organocatalyzed nucleophilic addition of cyanide or nitromethane to phosphorated ketimines⁹ and the diastereoselective addition of organometallic reagents to TADDOL-derived α -iminophosphonates¹⁰ for the asymmetric preparation of tetrasubstituted α -aminophosphonate derivatives.

In this case, we were intrigued about the possibility of accessing phosphorated analogues of aspartic acid holding tetrasubstituted carbons, using α -ketiminophosphonates as electrophile substrates in an asymmetric aza-Reformatsky reaction (Scheme 1). As far as we know, there are no examples of such a reaction using acyclic ketimines as substrates.

Moreover, it is well-known that α -aminophosphonates are an important class of compounds that can behave as stable analogues of the transition state of peptide cleavage. Due to this fact, they often inhibit enzymes implicated in proteolysis processes, and for this reason, α -aminophosphonic acid

Received: October 17, 2019

Acyclic ketimines

derivatives and their phosphapeptides show an assorted biological activity.¹¹ The biological activity of drugs in general¹² and α -aminophosphonic acids in particular¹³ is known to be strongly dependent on its absolute configuration, and for this reason, the development of new methods for the preparation of enantioenriched α -aminophosphonate derivatives is an important task in organic and medicinal chemistry.

Based on the zinc intermediates proposed by Noyori¹⁴ and inspired by the mechanisms for other enantioselective Reformatsky processes proposed by Cozzi⁵ and Feringa,¹⁵ initially we chose several diols and β -aminoalcohols as chiral inductors in the aza-Reformatsky reaction of α -ketiminophosphonates (Figure 1). Our required imine electrophile substrates can be easily obtained by a formal oxidation of the parent tertiary α -aminophosphonates as reported.⁹




Figure 1. Ligands tested in this study.

For the optimization of the reaction conditions, first we studied the reaction of dimethyl α -iminophosphonate 1 with ethyl iodoacetate in the presence of an organozinc reagent and 20 mol % of racemic BINOL I. Depending on the organozinc reagent used in the reaction, besides the desired product 2, compounds 3 and 4 were observed in the reaction mixture, resulting from a direct nucleophilic addition of organozinc reagent or the reduction of an imine bond, respectively (Table 1).

First, when diethylzinc was used as the metal source in dichloromethane, the reaction proceeded fast, but only reduced α -aminophosphonate 4 was observed (Table 1,

Table 1. Optimization of the Reaction Conditions

(MeO) ₂ F ز 1	Ph cat. I-	₂Et, ZnR₂ /, rt, 2 h	(MeO)	× .	NHTs ₂P R (MeO) Ö 3	NHTs 2 ^P Ph Ö 4
entry	solvent	R	cat.	conversion/%	2:3:4 ^c	ee/% ^d
1	$CH_2Cl_2^{a}$	Et	I	100	0:0:100	-
2	$CH_2Cl_2^{a}$	Me	Ι	100	0:100:0	-
3	$CH_2Cl_2^{\ b}$	Et	Ι	100	0:0:100	-
4	$CH_2Cl_2^{\ b}$	Me	Ι	100	90:10:0	-
5	toluene ^b	Me	I	100	90:10:0	-
6	DMF ^b	Me	Ι	100	90:10:0	-
7	THF ^b	Me	Ι	100	95:5:0	-
8	CH ₃ CN ^b	Me	I	100	97:3:0	-
9	CH ₃ CN ^b	Me	II	100	97:3:0	64
10	CH ₃ CN ^b	Me	III	100	97:3:0	99
11	CH ₃ CN ^b	Me	IV	100	97:3:0	38
12	CH ₃ CN ^b	Me	v	100	97:3:0	97

^{*a*}Under N₂ atmosphere. ^{*b*}Under dry air atmosphere. ^{*c*}Determined by ¹H NMR. ^{*d*}Determined by chiral HPLC.

entry 1). Then, in order to minimize the ratio of β -hydride transfer that leads to the formation of the reduced product, dimethylzinc was used as metal source, but only quaternary α -aminophosphonate **3** (R = Me) was observed, which presumably results from the nucleophilic addition of dimethylzinc to the imine electrophile.

It is known that diakylzinc species under an oxygen atmosphere are able to form the very reactive alkylperoxides (RZnOOR),¹⁶ which are excellent radical initiators. In fact, some enantioselective Reformatsky reactions have been reported, making use of organozinc reagents in the presence of air or oxygen.^{5,15,17} Consequently, the reaction was performed in dichloromethane under a dry air atmosphere using diethylzinc as the metal source, but although the reaction proceeded equally fast, again only reduced α -aminophosphonate 4 was observed (Table 1, entry 3). Switching to dimethylzinc as the metal source, the formation of aza-Reformatsky product 2 was observed together with 10% of 1,2addition product 3 (Table 1, entry 4). Due to the unfeasibility of the β -hydride transfer reaction, no formation of reduced compound 4 was noticed in this case. Although the same ratio 2:3 was obtained when the reaction was performed in low polar or strong polar solvents such as toluene or N,Ndimethylformamide, respectively (Table 1, entries 5 and 6), the amount of 2 could be increased using tetrahydrofurane or acetonitrile as solvents (Table 1, entries 7 and 8).

Next, we tested enantiopure (R)-BINOL (II) as a chiral ligand, under the optimal reaction conditions, but only modest enantioselectivity was obtained (Table 1, entry 9). To our delight, when anthracenyl-substituted BINOL ligand III was used in the same conditions, very high enantioselectivity was observed (Table 1, entry 10). Surprisingly, the use of (S)-VAPOL (IV) resulted in decreased enantiomeric excess compared to simple (R)-BINOL (II) (Table 1, entry 11). Finally, we even tested β -aminoalcohol derivative V that showed very good enantioselection (Table 1, entry 12).

With the optimal reaction conditions in our hands, we then extended the scope of the enantioselective aza-Reformatsky reaction to differently substituted α -ketiminophosphonates 1 (Figure 2). Excellent enantiomeric excesses were obtained in all cases. The reaction tolerates the presence of para- and metaalkyl substituents at the aromatic ring (Figure 2, 2b-c) as well as strong electron-donating groups at the para position of the aromatic imine (Figure 1, 2d). Several halogen-substituted aromatic ketimines were also successfully used in the reaction, including para-substituted aromatic rings containing fluorine, chlorine, or bromine (Figure 2, 2e-g), meta-substituted aromatic rings holding fluorine and chlorine (Figure 2, 2hi), as well as ortho-fluorophenyl-substituted ketimines (Figure 2, 2j). Moreover, dichloro- and difluoro-phenyl-substituted imines showed excellent enantioselectivities (Figure 2, 2k-m), and even perfluorophenyl-substituted substrates were tested with success using the same catalytic system (Figure 2, 2n).

Furthermore, an excellent result was observed for imine 10 holding an aromatic ring that is substituted with a chlorine and a strong electron-donating methoxy group. Besides, good reactivity and excellent enantioselectivity are also obtained using aromatic imines substituted with electron-withdrawing groups such as *p*-nitro or *p*-trifluoromethyl substituents (Figure 2, 2p-q). The reaction can be also extended to the use of imines holding biphenyl or heteroaromatic substituents (Figure 2, 2r-s). Finally, in order to further deprotect selectively the ester group, benzyl iodoacetate was used as a




Figure 2. Scope of the reaction.


reagent in the aza-Reformatsky reaction with our α iminophosphonate substrates, again with an excellent result (Figure 2, **2t**).

Next, with the purpose of illustrating the applications of our enantioenriched substrates, some synthetic transformations of α -aminophosphonate derivatives **2** were accomplished. A selective deprotection of the phosphonate group can be easily performed in a few hours using dimethylphosphonate derivative **2a** in the presence of trimethylsilyl bromide¹⁸ to afford α -aminophosphonic acid derivative **5** in very good yield.

For the selective deprotection of the ester group, we chose benzylic ester **2t** as the substrate. A vigorous stirring of compound **2t** under hydrogen atmosphere in the presence of a catalytic amount of palladium yielded the expected phosphorylated β -amino acid derivative **6** in excellent yield. Finally, activation of the carboxylic group in **6** with *N*,*N*-dicyclohexylcarbodiimide (DCC) in the presence of dimethylaminopyridine (DMAP) led to the formation of phosphorated β -lactam derivative **7** bearing a tetrasubstituted stereocenter (Scheme 2).

In order to authenticate the synthetic potential of our asymmetric methodology, a multigram-scale reaction was performed for the synthesis of α -aminophosphonate **2t**.

Scheme 2. Synthetic Applications of the Phosphorated Analogues of Aspartic Acid 2



Thus, 1.00 g of ketimine 1a reacted with benzyl iodoacetate in the presence of 20 mol % of BINOL ligand III to yield the aza-Reformatsky product 2t in 84% yield and 87% ee. With the aim of determining the absolute configuration of the stereogenic carbon of the major enantiomer in our substrates, using enantioenriched compound 2t, a substantial amount of the enantiopure β -lactam 7 was prepared. A crystal of the major enantiomer of 7 was isolated and submitted to X-ray diffraction analysis, revealing an S absolute configuration of the quaternary stereocenter (Figure 3).




Figure 3. X-ray structure of lactam 7 showing its absolute configuration (gray: C, white: H, red: O, blue: N, purple: P).

In summary, although successful enantioselective Reformatsky reactions have been described during the last years using acyclic aldehydes, aldimines, or ketones, we report here the first example of an enantioselective aza-Reformatsky reaction using acyclic ketimines as electrophile substrates. Phosphorated analogues of aspartic acid holding tetrasubstituted carbons are efficiently obtained with excellent enantioselectivity, using α -ketiminophosphonates, with a simply functionalized BINOL ligand as the chiral inductor. In addition, besides that phosphorus-containing enantiopure tetrasubstituted β lactams have not been reported so far, we describe here the first asymmetric synthesis of phosphorylated β -lactams using a catalytic approach.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.or-glett.9b03669.

Thermal ellipsoid plot for 7 and full experimental details, characterization, and copies of ¹H, ¹³C, and ³¹P NMR spectra for compounds **1**, **2**, **5**, **6**, and 7 and ¹⁹F NMR for compounds **1e**,**h**,**j**,**l**,**m**,**n**,**q** and **2e**,**h**,**j**,**l**,**m**,**n**,**q** (PDF)

Accession Codes

CCDC 1957474 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Authors

*E-mail: francisco.palacios@ehu.eus. *E-mail: javier.vicario@ehu.eus.

ORCID

Francisco Palacios: 0000-0001-6365-0324 Javier Vicario: 0000-0002-8976-0991

Author Contributions

The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Financial support by Ministerio de Ciencia, Innovación y Universidades (RTI2018-101818-B-I00, MCIU/AEI/FEDER, UE), and Gobierno Vasco (GV, IT 992-16) is gratefully acknowledged. The authors are thankful for the technical and human support provided by SGIker (UPV/EHU/ERDF, EU).

REFERENCES

(1) Reformatsky, S. Neue Synthese zweiatomiger einbasischer Säuren aus den Ketonen. Ber. Dtsch. Chem. Ges. **1887**, 20, 1210.

(2) For some reviews, see: (a) Ocampo, R.; Dolbier, W. R., Jr The Reformatsky reaction in organic synthesis. Recent advances. *Tetrahedron* **2004**, *60*, 9325. (b) Orsini, F.; Sello, G. Transition Metals-Mediated Reformatsky Reactions. *Curr. Org. Synth.* **2004**, *1*, 111. (c) Baba, A.; Yasuda, M.; Nishimoto, Y. Zinc enolates: the Reformatsky and Blaise reactions. In *Comprehensive Org. Synth.*; Knochel, P., Molander, G. A., Eds.; Elsevier: Amsterdam, 2014; pp 523–542.

(3) (a) Adrian, J. C.; Snapper, M. L. Multiple Component Reactions: An Efficient Nickel-Catalyzed Reformatsky-Type Reaction and Its Application in the Parallel Synthesis of β -Amino Carbonyl Libraries. J. Org. Chem. **2003**, 68, 2143. (b) Kanai, K.; Wakabayashi, H.; Honda, T. Rhodium-Catalyzed Reformatsky-Type Reaction. Org. Lett. **2000**, 2, 2549.

(4) (a) Cozzi, P. G. Reformatsky Reactions Meet Catalysis and Stereoselectivity. *Angew. Chem., Int. Ed.* 2007, *46*, 2568. (b) Fernández-Ibañez, M. A.; Macía, B.; Alonso, D. A.; Pastor, I. M. Recent Advances in the Catalytic Enantioselective Reformatsky Reaction. *Eur. J. Org. Chem.* 2013, *2013*, 7028. (c) Pellissier, H. Recent developments in the asymmetric Reformatsky-type reaction. *Beilstein J. Org. Chem.* 2018, *14*, 325.

(5) Cozzi, P. G. A Catalytic Enantioselective Imino-Reformatsky Reaction. *Adv. Synth. Catal.* **2006**, 348, 2075.

(6) (a) Tarui, A.; Nishimura, H.; Ikebata, T.; Tahira, A.; Sato, K.; Omote, M.; Minami, H.; Miwa, Y.; Ando, A. Ligand-Promoted Asymmetric Imino-Reformatsky Reaction of Ethyl Dibromofluoroacetate. Org. Lett. **2014**, *16*, 2080. (b) Tarui, A.; Ikebata, T.; Sato, K.; Omote, M.; Ando, A. Enantioselective synthesis of α,α -difluoro- β lactams using amino alcohol ligands. Org. Biomol. Chem. **2014**, *12*, 6484.

(7) (a) De Munck, L.; Vila, C.; Muñoz, M. C.; Pedro, J. R. Catalytic Enantioselective Aza-Reformatsky Reaction with Cyclic Imines. *Chem.* - *Eur. J.* **2016**, *22*, 17590. (b) De Munck, L.; Sukowski, V.; Vila, C.; Muñoz, M. C.; Pedro, J. R. Catalytic enantioselective-aza-Reformatsky reaction with seven-membered cyclic imines dibenzo[b₁f][1,4]-oxazepines. *Org. Chem. Front.* **2017**, *4*, 1624.

(8) (a) Quaternary Stereocentres: Challenges and Solutions for Organic Synthesis; Christoffers, J., Baro, A., Eds.; Wiley-VCH: Weinheim, 2006. (b) Shimizu, M. Construction of Asymmetric Quaternary Carbon Centers with High Enantioselectivity. Angew. Chem., Int. Ed. 2011, 50, 5998.

(9) (a) Vicario, J.; Ezpeleta, J. M.; Palacios, F. Asymmetric Cyanation of α -Ketiminophosphonates Catalyzed by Cinchona Alkaloids. Enantioselective Synthesis of Tetrasubstituted α -Aminophosphonic Acid Derivatives From Trisubstituted α -Aminophosphonates. Adv. Synth. Catal. 2012, 354, 2641. (b) Vicario, J.; Ortiz, P.; Ezpeleta, J. M.; Palacios, F. Asymmetric Synthesis of Functionalized Tetrasubstituted α -Aminophosphonates through Enantioselective Aza-Henry Reaction of Phosphorylated Ketimines. J. Org. Chem. 2015, 80, 156.

(10) Vicario, J.; Ortiz, P.; Palacios, F. Synthesis of Tetrasubstituted α -Aminophosphonic Acid Derivatives from Trisubstituted α -Aminophosphonates. *Eur. J. Org. Chem.* **2013**, 2013, 7095.

(11) (a) Hiratake, J.; Oda, J. Aminophosphonic and Aminoboronic Acids as Key Elements of a Transition State Analogue Inhibitor of Enzymes. *Biosci., Biotechnol., Biochem.* **1997**, *61*, 211. (b) Kafarski, P.; Lejczak, B. Aminophosphonic Acids of Potential Medical Importance. Curr. Med. Chem.: Anti-Cancer Agents **2001**, *1*, 301. (c) Berlicki, L.; Kafarski, P. Computer-Aided Analysis and Design of Phosphonic and Phosphinic Enzyme Inhibitors as Potential Drugs and Agrochemicals. Curr. Org. Chem. **2005**, *9*, 1829. (d) Van der Jeught, K.; Stevens, C. V. Direct Phosphonylation of Aromatic Azaheterocycles. Chem. Rev. **2009**, *109*, 2672.

(12) (a) Nguyen, L. A.; He, H.; Pham-Huy, C. Chiral Drugs: An Overview. *Int. J. Biomedical Sci.* **2006**, *2*, 85. (b) Kasprzyk-Hordern, B. Pharmacologically active compounds in the environment and their chirality. *Chem. Soc. Rev.* **2010**, *39*, 4466.

(13) (a) Kafarski, P.; Lejczak, B.; Szewczyk, J. Optically active 1aminoalkanephosphonic acids. Dibenzoyl-L-tartaric anhydride as an effective agent for the resolution of racemic diphenyl 1-aminoalkanephosphonates. Can. J. Chem. 1983, 61, 2425. (b) Atherton, F. R.; Hassall, C. H.; Lambert, R. W. Synthesis and structure-activity relationships of antibacterial phosphonopeptides incorporating (1aminoethyl)phosphonic acid and (aminomethyl)phosphonic acid. J. Med. Chem. 1986, 29, 29. (c) Drag, M.; Pawelczak, M.; Kafarski, P. Stereoselective synthesis of 1-aminoalkanephosphonic acids with two chiral centers and their activity towards leucine aminopeptidase. Chirality 2003, 15, S104. (d) Orsini, F.; Sello, G.; Sisti, M. Aminophosphonic acids and derivatives. Synthesis and biological applications. Curr. Med. Chem. 2010, 17, 264. (e) Kudzin, Z. H.; Kudzin, M. H.; Drabowicz, J. S. F.; Stevens, C. V. Aminophosphonic Acids - Phosphorus Analogues of Natural Amino Acids. Part 1: Syntheses of α -Aminophosphonic Acids. Curr. Org. Chem. 2011, 15, 2015. (f) Debrouwer, W.; Hertsen, D.; Heugebaert, T. S. A.; Boydas, E. B.; Van Speybrouck, V.; Catak, S.; Stevens, C. V. Tandem Addition of Phosphite Nucleophiles Across Unsaturated Nitrogen-Containing Systems: Mechanistic Insights on Regioselectivity. J. Org. Chem. 2017, 82, 188.

(14) Kitamura, M.; Suga, S.; Niwa, M.; Noyori, R. Self and Nonself Recognition of Asymmetric Catalysts. Nonlinear Effects in the Amino Alcohol-Promoted Enantioselective Addition of Dialkylzincs to Aldehydes. J. Am. Chem. Soc. **1995**, *117*, 4832.

(15) (a) Fernández-Ibáñez, M. A.; Maciá, B.; Minnaard, A. J.; Feringa, B. L. Catalytic enantioselective reformatsky reaction with aldehydes. *Angew. Chem., Int. Ed.* **2008**, 47, 1317. (b) Fernández-Ibáñez, M. A.; Maciá, B.; Minnaard, A. J.; Feringa, B. L. Catalytic enantioselective Reformatsky reaction with ketones. *Chem. Commun.* **2008**, 2571. (c) Fernández-Ibáñez, M. A.; Maciá, B.; Minnaard, A. J.; Feringa, B. L. Catalytic enantioselective Reformatsky reaction with ortho-substituted diarylketones. *Org. Lett.* **2008**, *10*, 4041.

(16) (a) Lewínski, J.; Śliwiński, W.; Dranka, M.; Justyniak, I.; Lipkowski, J. Reactions of [ZnR₂(L)] Complexes with Dioxygen: A New Look at an Old Problem. *Angew. Chem., Int. Ed.* 2006, 45, 4826.
(b) Lewínski, J.; Ochal, Z.; Bojarski, E.; Tratkiewicz, E.; Justyniak, I.; Lipkowski, J. First Structurally Authenticated Zinc Alkylperoxide: A Model System for the Epoxidation of Enones. *Angew. Chem., Int. Ed.* 2003, 42, 4643. (c) Bertrand, M.; Feray, L.; Gastaldi, S. Synthesis and selective transformation of nitrogen-containing compounds via radical

pathways. C. R. Chim. **2002**, *5*, 623 and references therein. . (17) (a) Tanaka, T.; Hayashi, M. Catalytic Enantioselective Reformatsky Reaction of Alkyl Iodoacetate with Aldehydes Catalyzed by Chiral Schiff Base. Chem. Lett. 2008, 37, 1298. (b) Wolf, C.; Moskowitz, M. Bisoxazolidine-Catalyzed Enantioselective Reformatsky Reaction. J. Org. Chem. 2011, 76, 6372.

(18) McKenna, C. E.; Higa, M. T.; Cheung, N. H.; McKenna, M. C. The facile dealkylation of phosphonic acid dialkyl esters by bromotrimethylsilane. *Tetrahedron Lett.* **1977**, *18*, 155.