Rapid Assembly of the Polyhydroxylated β -Amino Acid Constituents of Microsclerodermins C, D, and E

Thomas Hjelmgaard,[†] Sophie Faure,[†] Pascale Lemoine,[‡] Bernard Viossat,[‡] and David J. Aitken* $^{*,\$}$

Laboratoire de Synthèse et Etude de Systèmes d'Intérêt Biologique, SEESIB (UMR 6504 - CNRS), Université Blaise Pascal - Clermont-Ferrand 2, 24 avenue des Landais, 63177 Aubière cedex, France, Laboratoire de Cristallographie et RMN Biologiques (UMR 8015 - CNRS), Université René Descartes - Paris 5, 4 avenue de l'Observatoire, 75270 Paris cedex 06, France, and Laboratoire de Synthèse Organique et Méthodologie, ICMMO (UMR 8182 - CNRS), Université Paris-Sud 11, Bât. 420, 15 rue Georges Clemenceau, 91405 Orsay cedex, France

David.Aitken@u-psud.fr

Received December 11, 2007

LETTERS 2008 Vol. 10, No. 5 841–844

ORGANIC

ABSTRACT



AETD constituent of Microsclerodermin E: R = (E)-CH=CH(p-C₆H₄OEt)

A very short and efficient synthesis of protected derivatives of APTO and AETD, the complex polyhydroxylated β -amino acid residues present in microsclerodermins C, D, and E, is described. The targets are obtained in only five steps, in 23% and 16% overall yields, respectively. The key transformation involves the completely diastereoselective two-carbon homologation of appropriately selected intermediate chiral sulfinimines.

The microsclerodermins A–I and two dehydro derivatives constitute a family of complex cyclic peptides isolated from the lithistid sponges *Microscleroderma* sp. and *Theonella* sp. which possess antifungal and antitumor activities.¹ The family is characterized by a 23-atom macrocyclic ring comprising six amino acid residues, three of which are common. The more complex residues, which vary within the family, are a modified tryptophan, an unusual 3-aminopyrrolidone-4-acetic acid moiety, and a polyhydroxylated

10.1021/ol702962z CCC: \$40.75 © 2008 American Chemical Society Published on Web 02/07/2008

 β -amino acid residue containing four or five contiguous chiral centers. In microsclerodermins C and D (Figure 1) as well as their dehydro counterparts which contain a dehydropyrrolidone moiety, the latter constituent is (2*S*,3*R*,4*S*,5*S*,7*E*)-3-amino-8-phenyl-2,4,5-trihydroxyoct-7-enoic acid (APTO). In microsclerodermin E (Figure 1) this component is (2*S*,3*R*,4*S*,5*S*,7*E*,9*E*)-3-amino-10-(4-ethoxyphenyl)-2,4,5-trihydroxydeca-7,9-dienoic acid (AETD) (Figure 1).

To date, only the total synthesis of microsclerodermin E has been reported,² in which the AETD fragment was obtained in 16 steps and 10% overall yield. Recently, 12-step synthetic pathways to APTO (12% overall) and AETD (9% overall) derivatives were disclosed.³ In related studies,

[†] Université Blaise Pascal - Clermont-Ferrand 2.

[‡] Université René Descartes - Paris 5.

[§] Université Paris-Sud 11.

^{(1) (}a) Bewley, C. A.; Debitus, C.; Faulkner, D. J. J. Am. Chem. Soc. **1994**, *116*, 7631. (b) Schmidt, E. W.; Faulkner, D. J. *Tetrahedron* **1998**, *54*, 3043. (c) Qureshi, A.; Colin, P. L.; Faulkner, D. J. *Tetrahedron* **2000**, *56*, 3679. (d) Erdogan, I.; Tanaka, J.; Higa, T. *FABAD J. Pharm. Sci.* **2000**, *25*, 7.

⁽²⁾ Zhu, J.; Ma, D. Angew. Chem., Int. Ed. 2003, 42, 5348.

⁽³⁾ Shuter, E. C.; Duong, H.; Hutton, C. A.; McLeod, M. D. Org. Biomol. Chem. 2007, 5, 3183.



Figure 1. Microsclerodermins C, D, and E, with the APTO and AETD fragments highlighted in blue and red, respectively.

some synthetic approaches to building blocks for the construction of microsclerodermins A and B have appeared.⁴

Herein, we present our synthetic efforts in this field, providing a rapid and efficient construction of protected APTO and AETD derivatives. The retrosynthetic analysis for protected APTO, the first target, is outlined in Scheme 1. We envisaged construction of the α -hydroxy- β -amino acid



structural feature by stereocontrolled two-carbon homologation of chiral sulfinimine A2 with anion A1.⁵ Sulfinimine A2 should be accessible via a differentiated dialdehyde synthon A3, for which a convenient precursor should be the readily available 2-deoxy-D-ribose acetonide 1.⁶ Initial studies of the Wittig reaction of acetonide 1 with PhCH=PPh₃ provided the olefin 2 as an inseparable mixture of E/Z-isomers. Complete *E*-selectivity was achieved using Horner–Wadsworth–Emmons methodology (Scheme 2).



Best results (see Supporting Information) were obtained using ^{*n*}BuLi as base, which afforded pure **2** in 52% yield. A modified Swern oxidation⁷ of hydroxy acetonide **2** provided aldehyde **3** in 85% yield. Ensuing epimerization of the α -stereocenter with K₂CO₃ in methanol using an adaptation of the literature method⁸ gave aldehyde **4** with the requisite stereochemistry in 83% yield. Aldehyde **4** was then converted into chiral sulfinimines **5a** (50%), **5b** (74%), and **5c** (69%) by reaction with (*R*)-(-)-*p*-toluenesulfinamide, (*R*)-(+)-*tert*butanesulfinamide, and (*S*)-(-)-*tert*-butanesulfinamide, respectively, in the presence of Ti(OEt)₄.⁹

In support of the planned two-carbon homologation, we were encouraged by a recent report that the addition of enolates of α -hydroxyacetates bearing a bulky *O*-protecting group (Boc) to simple chiral sulfinimines proceeded in a highly diastereoselective manner.¹⁰ This observation was exploited in an illustrative synthesis of the taxol side chain with a 2*R*,3*S* configuration. Thus, we examined this reported procedure with a view to obtaining protected APTO fragments **7** with a 2*S*,3*R* configuration by reaction of sulfinimines **5** with protected α -hydroxyacetates **6** (Table 1). Initially, the products were obtained in disappointingly low yields, but inversing the reagent addition order (addition of sulfinimine to the enolate) provided a dramatic improvement. 'Bu-sulfinimines **5b** and **5c** (entries 3, 4, 6) provided much higher selectivities and better yields than *p*-tolylsulfinimine

^{(4) (}a) Sasaki, S.; Hamada, Y.; Shioiri, T. *Tetrahedron Lett.* **1997**, *38*, 3013. (b) Sasaki, S.; Hamada, Y.; Shioiri, T. *Synlett* **1999**, 453. (c) Sasaki, S.; Hamada, Y.; Shioiri, T. *Tetrahedron Lett.* **1999**, *40*, 3187. (d) Shioiri, T.; Sasaki, S.; Hamada, Y. *ARKIVOC* **2003**, 103. (e) Chandrasekhar, S.; Sultana, S. S. *Tetrahedron Lett.* **2006**, *47*, 7255.

⁽⁵⁾ For reviews on chiral sulfinimines, see: (a) Davis, F. A.; Zhou, P.; Chen, B.-C. *Chem. Soc. Rev.* **1998**, *27*, 13. (b) Ellman, J. A.; Owens, T. D.; Tang, T. P. *Acc. Chem. Res.* **2002**, *35*, 984. (c) Zhou, P.; Chen, B.-C.; Davis, F. A. *Tetrahedron* **2004**, *60*, 8003. (d) Brinner, K.; Ellman, J. A. In *Enantioselective Synthesis of* β -*Amino Acids*, 2nd ed.; Juaristi, E., Soloshonok, V. A., Eds.; Wiley-VCH: New Jersey, 2005; Chapter 8.

⁽⁶⁾ Barbat, J.; Gelas, J.; Horton, D. *Carbohydr. Res.* **1983**, *116*, 312. (7) Hjelmgaard, T.; Søtofte, I.; Tanner, D. J. Org. Chem. **2005**, *70*, 5688

⁽⁸⁾ Servi, S. J. Org. Chem. 1985, 50, 5865.

⁽⁹⁾ Davis, F. A.; Prasad, K. R.; Carroll, P. J. J. Org. Chem. 2002, 67, 7802.

⁽¹⁰⁾ Wang, Y.; He, Q.-F.; Wang, H.-W.; Zhou, X.; Huang, Z.-Y.; Qin, Y. J. Org. Chem. 2006, 71, 1588.



^{*a*} As judged by NMR. ^{*b*} Complex mixtures obtained. ^{*c*} Only one isomer observed. ^{*d*} Reagent decomposed. ^{*e*} Determined by X-ray crystallography. All = allyl; Tr = trityl.

5a (entries 1, 2). The system appeared very sensitive to substituent effects: changing the enolate from **6** (All/Boc) to **6** (Me/Boc) improved the stereoselectivity to effective completeness, whereas the enolate from **6** (Me/Tr) decomposed in the same conditions (entries 3-5). The reactions of **6** (Me/Boc) with **5b** and **5c** thus satisfyingly provided pure APTO fragments **7b** and **7c** (R = 'Bu, R¹ = Me, Pg = Boc) as single diastereomers in 66% and 94% yield, respectively (entries 4, 6).

The absolute stereochemistry of **7b** and **7c** was determined by means of single-crystal X-ray diffraction (Figure 2). To our surprise we found that **7b** (derived from (R)-(+)-*tert*butanesulfinamide) possessed the 2R,3S configuration, whereas **7c** (derived from (S)-(-)-*tert*-butanesulfinamide) possessed the desired 2S,3R configuration!¹¹

The formation of **7c** from **5c** cannot be explained rationally by a 6/4 bicyclic transition-state model. A more comfortable explanation for the observed stereoselectivity might be a nonchelated reaction pathway (Figure 3a), proposed earlier for similar systems.¹² Alternatively, a 6/6 chelated bicyclic chairlike transition state where the most bulky groups are in equatorial positions can be proposed (Figure 3b). Both these models are in accordance with the observed steric demands of our system: the sulfinimine moiety and the *O*-protecting group of **6** need to be as bulky as possible, while only the diminutive methyl ester of **6** gives complete selectivity.

It is noteworthy that the stereochemical outcome of this two-carbon homologation is controlled by the sulfinyl group only and thus proceeds independently of other neighboring chiral centers in the molecule. This chiral predominance phenomenon was previously noted for sulfinimine onecarbon homologation with cyanide.⁹



Figure 2. X-ray structures of 7b (top) and 7c (bottom).

Having developed this short stereoselective pathway to the protected APTO fragment of microsclerodermins C and D, we then turned our attention to the synthesis of the AETD fragment of microsclerodermin E (Scheme 3). In contrast to the results obtained in the synthesis of olefin 2, all attempts to use Horner–Wadsworth–Emmons methodology proved fruitless for the synthesis of olefin 9 from 1. After numerous explorations (see Supporting Information), access to the desired alcohol was only found possible through use of phosphonium bromide 8 in benzene solution in the presence of KHMDS as base.³ The product was obtained as an E/Z-



Figure 3. Two plausible models, (a) nonchelated and (b) chelated, for interactions of enolate 6 (Me/Boc) with 5c leading to the observed selectivity of 7c.

⁽¹¹⁾ These stereochemical observations are in contradiction with the chemical structures presented in ref 10 and, in consequence, are in disaccord with arguments presented therein to account for the stereoselectivity. (12) Fujisawa, T.; Kooriyama, Y.; Shimizu, M. *Tetrahedron Lett.* **1996**, *37*, 3881.



mixture which was easily separated by column chromatography to furnish the desired pure (*E*)-**9** in 49% yield as well as 19% of pure (*Z*)-**9**. Swern oxidation of (*E*)-**9** (78%) followed by isomerization with K_2CO_3 (78%) provided aldehyde **11**. The latter was converted into sulfinimine **12** using (*S*)-(-)-*tert*-butanesulfinamide in 70% yield. Reaction

with methyl *O*-Boc-hydroxyacetate **6** (Me/Boc) according to our optimized conditions then provided the protected AETD fragment **13** in 79% yield as a single diastereomer. Compound **13** showed spectral data which were very similar to those of **7c**.

In conclusion, we have established very short and effective syntheses of protected APTO **7c** and protected AETD **13**. These proceed in 23% and 16% overall yield, respectively, in only 5 steps from the readily available acetonide-protected 2-deoxy-D-ribose **1**. This approach compares favorably with the existing pathways for the preparation of comparable intermediates for the synthesis of microsclerodermins C, D, and E and is furthermore not restricted to small-scale synthesis.

Acknowledgment. We thank the French Ministry for Research for financial support.

Supporting Information Available: Full experimental procedures, characterization data, and NMR spectra for all intermediates and final products; optimization data for the synthesis of alcohols **2** and **9**; CIF files for **7b** and **7c**. This material is available free of charge via the Internet at http://pubs.acs.org.

OL702962Z