Accepted Manuscript

Prediction of photophysical properties of pyrimidine chromophores using Taguchi method

Charline Denneval, Sylvain Achelle, Christine Baudequin, Françoise Robin-le Guen

PII: S0143-7208(14)00218-6

DOI: 10.1016/j.dyepig.2014.05.030

Reference: DYPI 4402

To appear in: Dyes and Pigments

Received Date: 21 March 2014

Revised Date: 20 May 2014

Accepted Date: 22 May 2014

Please cite this article as: Denneval C, Achelle S, Baudequin C, Robin-le Guen F, Prediction of photophysical properties of pyrimidine chromophores using Taguchi method, *Dyes and Pigments* (2014), doi: 10.1016/j.dyepig.2014.05.030.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Prediction of photophysical properties of pyrimidine chromophores using Taguchi method.

Charline Denneval,¹ Sylvain Achelle,^{2*} Christine Baudequin,¹ and Françoise Robin-le Guen.²

¹ Normandie Univ. COBRA, UMR 6014 et FR 3038; Univ. Rouen; INSA Rouen; CNRS,

IRCOF, 1 rue Tesnière, 76821 Mont Saint Aignan Cedex, France

² Institut des Sciences Chimiques de Rennes UMR CNRS 6226, IUT de Lannion, rue Edouard

Branly, BP 30219, F22302 Lannion Cedex, France

Corresponding author: sylvain.achelle@univ-rennes1.fr Tel: +33 (0)2 96 46 94 48 Fax : + 33 (0)2 96 46 93 54.

Abstract

This paper presents the photophysical properties of a series of pyrimidine chromophores. The influence of three parameters has been studied: the nature of the electron-donating group and of the π -conjugated linker as well as the substituted position of the pyrimidine core (2, 4 and 6 position). Taguchi design of experiment combined with analysis of variance methodologies have been employed. This approach has permitted to highlight the main factor affecting the absorption and emission wavelengths of the studied compounds. Absorption maxima of this family of compounds can be predicted with relatively good precision (± 18 nm). The emission maxima can be also predicted, however the precision is lower (± 39 nm). Taguchi methodology has been also employed with these families of compounds to optimize the intramolecular charge transfer estimated by emission solvatochromy.

Key words: Pyrimidine, Photophysical Properties, Intramolecular Charge Transfer, Design of Experiment, Dyes

1. Introduction

Donor-acceptor π -conjugated (D- π -A) materials, also called push-pull materials, with both electron-donating and electron-withdrawing groups linked by a π -conjugated bridge have been extensively studied during the past decades [1]. In such molecules, Intramolecular Charge Transfer (ICT) that occurs can be easily illustrated by two limiting resonance forms (aromatic and quinoid/zwitterionic arrangements) [2]. Push-pull structures have found useful applications in fluorescent dyes and sensors [3], sensitizers for Dye-Sensitized Solar-Cells (DSSCs) [4] and emitters for Organic Light-Emitting Diodes (OLED) [5]. This molecular design is also typical of second and third order Non-Linear Optical (NLO) chromophores [6]. Second order NLO materials have found applications in green laser obtained from red sources through frequency doubling, in second-harmonic generation microscopy or in terahertz wave

generation [7]. Third order NLO, in particular two-photon absorption (TPA) materials have also attracted considerable attention due to their applications in photodynamic therapy, confocal microscopy, optical power limiting, 3D data storage and microfabrication [8]. In case of second order NLOphores, the NLO answer ($\mu\beta$) is directly related to the intensity of the ICT phenomena [9].

Pyrimidine, which is a highly π -deficient heterocycle, can be used as electron-withdrawing part in push-pull structure. The ability of protonation, hydrogen bond formation and chelation of the nitrogen atoms of the pyrimidine ring are also of great importance: such derivatives could be therefore used for the formation of supramolecular assemblies and as sensors. The use of pyrimidine ring in the structure of π -conjugated materials has been recently reviewed [10]. During the past decade, hundreds of pyrimidine chromophores have been designed. In particular numerous arylpyrimidine derivatives have been studied as fluorescent dyes [11]. Moreover, it should be noted that 4,6-di(arylvinyl)pyrimidine is now a well established structure of TPA dyes [12]. Recently, the second order NLO properties of 4arylvinylpyrimidine have been also studied [13].

Even if the solvent polarity affects the absorption spectra of push-pull compounds [14], the emission solvatochromism is generally more intense. Broad structureless red shifted emission is observed on increasing the solvent polarity along with a successive decrease in the fluorescence intensity. This solvatochromic behavior results from the stabilization of the highly polar emitting state by polar solvents [15]. Therefore studying the emission solvatochromism is a way to evaluate ICT upon excitation in push-pull structures. Different scales have been described for quantification of solvent polarity [16], one of them is the Dimroth-Reichardt ($E_T(30)$) polarity scale based on the solvatochromic method [17].

Taguchi robust design method is a combination of mathematical and statistical techniques used in an empirical study [18]. The Taguchi design of experiment method has become a

standard method to enhance the quality of products and processes. In principle, the Taguchi's design of experiments is used to get information such as main effects and interaction effects of design parameters from minimum number of experiments. It consists in an experimental strategy in which the effects of multiple factors are studied simultaneously by running tests with the core factors at various levels. In the Taguchi approach, experiments are designed using special orthogonal arrays that allow the parameter space to be examined by performing a limited number of experiments. It is therefore economical for characterizing a complicated process. Fewer experiments are required to study all levels of all input parameters, and filters out some effects due to statistical variation. Taguchi design of experiments method has been widely used in chemistry for the optimization of synthetic procedure [19]. However, only few examples of statistical studies dedicated to the photophysical data of series of dyes have been described. Bureš and coworkers used statistical treatment to rationalize the photophysical data of a series of heterocyclic compounds [20], and more recently, Chen *et al* described the optimization of luminescence properties of materials by design of experiments methodologies [21]

The aim of the work described in this paper is to predict the absorption and emission maxima and to optimize ICT, estimated by the importance of emission solvatochromism, of a series of pyrimidine chromophores (Figure 1). The influence of three parameters has been studied: the nature of the electron-donating group and of the π -conjugated linker as well as the substituted position of the pyrimidine core (2, 4 and 6 position).

<Figure1>

2. Materials and method

2.1. method

In order to predict the absorption and emission maxima and to optimize ICT in pyrimidine chromophores, three parameters have been studied: the nature of the electron-donating group,

the nature of the π -conjugated linker and the substituted position of the pyrimidine core. For each parameter, three levels have been considered (Table 1). The orthogonal array of L₉ type was used and is represented in Table 2, where column D is used as control to check if there are interactions between factors A-C. The corresponding compounds are represented in Figure 2. It should be noted that in case of compound **3**, for solubility reasons, the methoxy groups have been replaced by hexyloxy group. This modification is supposed to have a negligible influence on the photophysical parameters studied.

<Table 1>

<Table 2>

<Figure 2>

2.2. General experimental

All solvents were reagent grade. The starting materials were purchased from Sigma-Aldrich or Acros and used without further purification. For air and moisture-sensitive reactions, all glassware pieces were flame-dried and cooled under nitrogen. NMR spectra were recorded in CDCl₃, on a Bruker Advance 300 spectrometer (¹H at 300 MHz and ¹³C at 75 MHz). The chemical shifts δ are reported in ppm relative to the residual solvent (CHCl₃). The coupling constants *J* are given in Hz. In the ¹H NMR spectra, the following abbreviations are used to describe the peak patterns: s (singlet), d (doublet), dd (doublet of doublet), t (triplet), m (multiplet). In the ¹³C NMR spectra, the nature of the carbons (C, CH, CH₂ or CH₃) was determined by performing a JMOD experiment. Melting points (°C) were measured on a Kofler hot-stage with a precision of 2 degrees (± 2°C). High-resolution mass analyses were performed at the "Centre Régional de Mesures Physiques de l'Ouest" (CRMPO, University of Rennes1) using a Bruker MicroTOF-Q II apparatus. Analytical thin layer chromatographies (TLC) were performed on silica gel plates (Merck 60F254) and compounds were visualized

by irradiation with UV light at 254 and 365 nm. Flash chromatography was performed using silica Acros SI 60 (60-200 mesh ASTM). UV-Visible spectra were recorded on a Varian Cary 50 scan spectrophotometer. Fluorescence spectroscopy studies were conducted on a Varian Cary Eclipse spectrophotometer. Compounds were excited at their absorption maxima (band of lowest energy) for recording the emission spectra. All solutions were measured with optical densities below 0.1. Fluorescence quantum yield (\pm 10%) determined relative to quinine sulfate in 1 M H₂SO₄ ($\Phi_F = 0.54$) or harmane in 0.1 M H₂SO₄ as standard ($\Phi_F = 0.83$). Stokes shifts were calculated considering the lowest energetic absorption band.

2.3. Synthesis

Compounds 1 [22], 2 [23], 3 [24], 5 [13a], 6 [11a], 9 [22] have been synthesized according to reported methods. The synthetic routes to chromophores 4, 7, 8 and 12 are reported in schemes 1-4.

<Scheme 1>

<Scheme 2>

<Scheme 3>

<Scheme 4>

2.4. General Knoevenagel condensation procedure

A stirred mixture of methylpyrimidine derivative (1.5 mmol) and corresponding aldehyde (1.5 mmol) in aqueous sodium hydroxide (5 M, 15 mL) containing Aliquat 336 (64 mg, 0.15 mmol) was heated under reflux for 3 h. The mixture was allowed to cool, the precipitate was filtered off, and washed with water.

2.5.Synthesis of 4

Compound **4** was synthesized according to general Knoevenagel condensation procedure and the crude product was purified by column chromatography (SiO₂, AcOEt/Petroleum Ether, 1:1); yellow solid; 216 mg; yield 64%; mp : 149-152°C; ¹H NMR (300 MHz, CDCl₃) δ 3.01 (s, 6H), 6.71 (d, 1H, *J* = 9.0 Hz), 7.01 (t, 1H, *J* = 4.8 Hz), 7.03 (d, 1H, *J* = 16.2 Hz), 7.53 (d, 2H, *J* = 9.0 Hz), 7.92 (d, 1H, *J* = 16.2 Hz), 8.65 (d, 2H, *J* = 4.8 Hz); ¹³C NMR and JMOD (75 MHz, CDCl₃) δ 165.7 (C), 156.9 (CH), 151.1 (C), 138.5 (CH), 129.1 (C), 124.1 (C), 122.5 (CH), 117.6 (CH), 112.1 (CH), 40.2 (CH₃); Anal. Calcd for C₁₄H₁₅N₃: C, 74.64; H, 6.71; N, 18.65. Found C, 74.51; H, 6.95; N, 18.37.

2.6. Synthesis of **10**

Compound **10** was synthesized according to general Knoevenagel condensation procedure and the crude product was purified by column chromatography (SiO₂, AcOEt/Petroleum Ether, 1:1); cream solid; 172 mg; yield 44%; ¹H NMR (300 MHz, CDCl₃) δ 7.12 (t, 1H, *J* = 4.8 Hz), 7.22 (d, 1H, *J* = 16.2 Hz), 7.48 (d, 1H, *J* = 8.7 Hz), 7.52 (d, 2H, *J* = 8.7 Hz), 7.91 (d, 1H, *J* = 16.2 Hz), 8.72 (d, 2H, *J* = 4.8 Hz); ¹³C NMR and JMOD (75 MHz, CDCl₃) δ 164.7 (C), 157.1 (CH), 136.8 (CH), 134.9 (C), 132.0 (CH), 129.0 (CH), 128.1 (CH), 123.1 (C), 118.7 (CH); Anal. Calcd for C₁₂H₉BrN₂: C, 55.20; H, 3.47; N, 10.73. Found C, 55.15; H, 3.56; N, 10.57.

2.7. General Suzuki cross-coupling reaction procedure

A stirred mixture of halogeno derivative (1 mmol), corresponding phenylboronic acid (1.2 mmol), $Pd(PPh_3)_4$ (58 mg, 0.05 mmol), aqueous 1 M sodium carbonate (1 mmol, 1 mL) and ethanol (1 mL) in degassed toluene (10 mL) was heated under nitrogen for 24 h (40h in case of **12**). The reaction mixture was cooled, filtered, and dissolved with a mixture of EtOAc and water 1:1 (50 mL) and the organic layer separated. The aqueous layer was extracted with EtOAc (2 × 25 mL). The combined organic extracts were dried with MgSO₄ and the solvents evaporated.

2.8. Synthesis of 7

Compound **7** was synthesized according to general Suzuki cross coupling reaction procedure and the crude product was purified by column chromatography (SiO₂, AcOEt/Petroleum Ether, 1:1); yellow solid; 296 mg; yield 70%; mp : 182-186°C; ¹H NMR (300 MHz, CDCl₃) δ 7.10-7.03 (m, 3H), 7.16-7.12 (m, 7H), 7.31-7.28 (m, 4H), 7.52 (d, 2H, *J* = 8.7 Hz), 7.62 (d, 2H, *J* = 8.4 Hz), 7.69 (d, 2H, *J* = 8.4 Hz), 8.03 (d, 1H, *J* = 16.2 Hz), 8.74 (d, 2H, *J* = 4.8 Hz); ¹³C NMR and JMOD (75 MHz, CDCl₃) δ 165.0 (C), 157.1 (CH), 147.6 (C), 147.5 (C), 141.3 (C), 137.8 (CH), 134.5 (C), 134.1 (C), 129.3 (CH), 128.2 (CH), 127.6 (CH), 127.0 (CH), 126.9 (CH), 124.6 (CH), 123.7 (CH), 123.1 (CH), 118.4 (CH); HRMS (ESI+, ASAP) m/z calculated for C₃₀H₂₄N₃ [M+H]⁺ 426.1970, found 426.1968.

2.9.Synthesis of 11

Compound **11** was synthesized according to general Suzuki Cross coupling reaction procedure and the crude product was purified by column chromatography (SiO₂, AcOEt/Petroleum Ether, 1:9); pale yellow solid; 250 mg; yield 70%; mp : 147-149°C; ¹H NMR (300 MHz, CDCl₃) δ 7.20-7.10 (m, 8H), 7.37-7.31 (m, 4H), 7.65 (s, 1H), 7.94 (d, 2H, *J* = 8.7 Hz),8.96 (s, 1H); ¹³C NMR and JMOD (75 MHz, CDCl₃) δ 165.1 (C), 161.5 (C), 158.9 (CH), 151.3 (C), 146.7 (C), 129.7 (CH), 128.4 (CH), 127.5 (C), 125.7 (CH), 124.4 (CH), 121.3 (CH), 115.8 (CH); HRMS (ESI⁺, ASAP) m/z calculated for C₂₂H₁₇ClN₃ [M+H]⁺ 358.1106, found 358.1109.

2.10. Synthesis of 8

To a stirred mixture of compound **11** (150 mg, 0.42 mmol), sodium hydrogenocarbonate (43 mg, 0.50 mmol) and Pd-C (5 mg, 0.04 mmol) in EtOH (5 mL), dihydrogen was introduced in bubbles at rt for 20 h. The mixture was then filtered with celite and the solvent removed. The crude product was purified by column chromatography (SiO₂, AcOEt/Petroleum Ether, 1:4); yellow solid; 124 mg; yield 92%; mp : 104-107°C; ¹H NMR

(300 MHz, CDCl₃) δ 7.08-7.17 (m, 8H), 7.25-7.33 (m, 4H), 7.61 (d, J = 5 Hz, 1H), 7.95 (d, J = 8.7 Hz, 2H), 8.67 (d, J = 5 Hz, 1H), 9.19 (s, 1H); ¹³C NMR and JMOD (75 MHz, CDCl₃) δ 163.3 (C), 158.9 (CH), 157.0 (C), 150.7 (CH), 146.9 (CH), 129.5 (CH), 128.9 (C), 128.1 (CH), 125.4 (CH), 124.0 (CH), 121.7 (CH), 116.1 (C); HRMS (ESI⁺, ASAP) m/z calculated for C₂₂H₁₇N₃ [M+H]⁺ 324.1501, found 324.1511

2.11. Synthesis of **12**

Compound **12** was synthesized according to general Suzuki Cross coupling reaction procedure and the crude product was purified by crystallization from of mixture of DCM/pentane; orange solid; 261 mg; yield 34%; mp : $125-130^{\circ}$ C; ¹H NMR (300 MHz, CDCl₃) δ 7.03-7. 05 (m, 6H), 7.12-7.16 (m, 12H), 7.26-7.32 (m, 9H), 7.51 (d, *J* = 8.7 Hz, 4H), 8.13 (d, *J* = 8.7 Hz, 4H), 8.15 (d, *J* = 8.7 Hz, 4H), 7.95 (d, *J* = 15.9 Hz, 2H), 9.12 (s, 1H); ¹³C NMR and JMOD (75 MHz, CDCl₃) δ 162.9 (C), 158.8 (C), 147.8 (C), 147.6 (C), 141.7 (CH), 136.8 (CH), 134.1 (C), 133.8 (C), 129.5 (CH), 128.3 (CH), 127.7 (CH), 127.0 (CH), 125.5 (CH), 124.7 (CH), 123.7 (CH), 123.3 (CH), 116.6(CH); HRMS (ESI⁺, ASAP) m/z calculated for C₅₆H₄₂N₄ [M+H]⁺ 771.3480, found 771.3488.

3. Results and discussion

The UV-Vis and photoluminescence (PL) spectroscopic data of compounds **1-9** measured in dichloromethane at 25 °C are presented in table 3. Analyses have been carried out using low concentration solutions in order to avoid self-absorption effect. As an example, the spectra for derivative **9** are shown in Figure 3. All compounds are photostable during photophysical experiments.

<Table 3>

<Figure 3>

The absorption and emission maxima have been studied according to Taguchi methodology and analysis of variance (ANOVA) was conducted to determine the significance of each factor. For absorption, the effects of each level of each factor are presented in Table 4 and the results of ANOVA calculations are given in Table 5. The average absorption wavelength is 367 nm. As shown in Table 4, the diphenylamino group (level 3 of factor A) is the substituent leading to the most red-shifted absorption. 2-substituted pyrimidine (level 1, factor B) are the most blue shifted whereas 4,6-disubstituted derivatives (level 3, factor B) are the most red shifted. Concerning the π -conjugated core, phenylenevinylene derivatives (level 2, factor C) are the most red shifted.

Similar tables for emission are given in supporting information, the average emission wavelength is 476 nm. It should be noted that the substitution position of the pyrimidine core affect differently the absorption and the emission: 4-monosubstituted derivatives (level 2, factor B) are more red shifted in emission than 4,6-disubstituted pyrimidines (level 3, factor B).

Either for λ_{abs} and λ_{em} , the influence of residue is low (below 3.3%) indicating that there is no interaction between the three factors studied. The F-ratio test was conducted to evaluate the effect of each factor on the signal/noise ratio. For absorption, the three factors A, B and C have F-ratios superior to the F-value for a 90% confidence level (F-value_{0.9} = 9) making them significant. For emission, only factors A and C are significant. As shown in table 4, the nature of the electron-donating group (factor A, 48.0%) and the substituted position of the pyrimidine ring (factor B, 28.4%) are the main factor influencing the absorption wavelength. The emission wavelength mainly depends on the nature of the π -conjugated core (factor C, 45.3%) and of the electron-donating group (factor A, 44.3%).

The absorption wavelengths calculated according to the average value and the effects of the corresponding level of each factor (Table 4) are in good agreement with experimental values

for the compounds described in the literature ($\Delta\lambda < 18$ nm, see supporting information). For emission, the differences between calculated and experimental values are a bit higher ($\Delta\lambda <$ 39 nm, see supporting information) in particular for the most red-shifted compounds.

<Table 4>

<Table 5>

In an effort to gain further insight into the photophysical process within this series of pyrimidine push-pull chromophores, we investigated the emission behavior of compounds **1-9** in different aprotic solvents. The results of these investigations are summarized in Table 6. As an example the emission spectra in various solvents for compound **6** are shown in Figure 3. For all compounds, the emission maxima were plotted *versus* the Dimroth-Reichardt polarity parameter ($\Delta E_T(30)$). In all case a good linearity was obtained (see supporting information). The slopes of the corresponding regression lines (SP) are good indications of the intensity of the ICT into the chromophores. The highest value of SP is observed for compound **5**.

<Table 6>

<Figure 4>

The ANOVA method was employed to study the influence of each factor on the value of SP in a similar manner than for absorption and emission maxima. The results are given in Tables 7 and 8. The average value of SP is 1.73. Once again, the influence of the residue is low (2.1%). Two factors significantly explain SP according to their F-ratios: the nature of the electron-donating group (factor A, 46.8 %) and of the π -conjugated core (factor C, 40.4%). The substituted position of the pyrimidine ring seems to be less important. According to Table 7, the optimal configuration consists in diphenylamino donor groups (level 3 factor A) linked to a 4-6-disubstituted pyrimidine (level 3 factor B) by a biphenylenevinylene bridge (level 3

factor C). The corresponding structure (compound **12**) exhibits a higher SP value than compounds **1-9** which validates the methodology.

<Table 7>

<Table 8>

4. Conclusion

To conclude, we have shown that Taguchi design of experiment methodology can be efficiently applied to predict the photophysical properties (absorption and emission maxima) of a family of pyrimidine chromophores. ICT in these compounds estimated by emission solvatochromy has been also optimized using the same approach. Taguchi design of experiment methodology appears useful to study the influence various parameters involved in the photophysical properties of a series of dyes and to identify quickly the optimized chromophore. This strategy could be applied to other family of chromophores.

Figure 1. General structure of the studied dyes.

Figure 2. Chromophores 1, 2, 3, 4, 5, 6, 7, 8, 9

Figure 3. Absorption (broken line) and emission spectra (solid line) of chromophores 9

Figure 4. Normalized emission spectra of chromophore 6 in various solvents.

Scheme 1. Synthesis of compound 4

Scheme 2. Synthesis of compound 7.

Scheme 3. Synthesis of compound 8.

Scheme 4. Synthesis of compound 12.

 Table 1. Parameters and their values corresponding to their levels to be studied in experiments.

Table 2. Experimental layout using the L₉ orthogonal array.

Table 3. UV/Vis and photoluminescence (PL) data

Table 4. Effect of each level of each factor on λ_{abs} in nm

Table 5. Analysis of variance (ANOVA) for on λ_{abs} in nm.

 Table 6. Emission solvatochromic range of pyrimidine chromophores 1-9 and 12 in various aprotic solvents

Table 7. Effect of each level of each factor on SP

Table 8. Analysis of variance of SP

[1]a) Gompper R, Wagner HU. Donor-Acceptor-Substituted π -Electron Systems–Probes for Theories and Building Blocks for New Materials. Angew Chem Int Ed Engl 1988;27:1437-55. b) Kivala M, Diederich F. Acetylene-derived strong organic acceptors for planar and nonplanar push-pull chromophores. Acc Chem Res;42:235-48.

[2]Kulhánek J, Bureš F, Opršal J, Kuznik W, Mikysek T, Růžička A. 1,4-Phenylene and 2,5thienylene π -linkers in charge-transfer chromophores. Asian J Org Chem 2013;2:422-31.

[3]See for example : a) Guliyev R, Coskun A, Akkaya EU, Design strategies for ratiometric chemosensors: modulation of excitation energy transfer at the energy donor site. J Am Chem Soc 2009;131:9007-13. b) Costero AM, Parra M, Gil S, Gotor R, Mancini PME, Martínez-Máñez R, Sancenón F, Royo S. Chromo-fluorogenic detection of nerve-agent mimics using triggered cyclization reactions in push-pull dyes. Chem Asian J 2010;5:1573-85. c) Hadad C, Achelle S, López-Solera I, García-Martínez JC. Metal cation complexation studies of 4-arylvinyl-2,6-di(pyridin-2-yl)pyrimidines: effect on the optical properties. Dyes Pigm 2013:97:230-7. d) Nepraš M, Almonasy N, Bureš F, Kulhánek J, Dvořák M, Michl M. Fluorescence and photophysical properties of D- π -A push-pull systems featuring a 4,5-dicyanoimidazole unit. Dyes Pigm 2011;91:466-73. e) Denneval C, Moldovan O, Baudequin C, Achelle S, Baldeck P, Plé N, Darabantu M, Ramondenc Y. Synthesis and photophysical properties of push-pull structures incorporating diazines as attracting part with a fluorene core. Eur J Org Chem 2013;5591-06.

[4] a) Lin Y, Li Y, Zhan X. Small molecule semiconductors for high-efficiency organic photovoltaics. Chem Soc Rev 2012;41:4245-72. b) Hagfeldt A, Boschloo G, Sun L, Kloo L, Pettersson H. Dye-sensitized solar cells. Chem Rev 2010;110:6595-663. c) Günes S, Neugebauer H, Sariciftci NS. Conjugated polymer-based organic solar cells. Chem Rev 2007;107:1324-38.

[5] a) Walzer K, Maennig B, Pfeiffer M, Leo K. Highly efficient organic devices based on electrically doped transport layers. Chem Rev 2007;107:1233-71. b) Yang Y, Zhou Y, He QG, He C, Yang CH, Bai FL, Li YF. Solution-processable red-emission organic materials containing triphenylamine and benzothiodiazole units: synthesis and applications in organic light-emitting diodes. J Phys Chem B 2009;113:7745-52.

[6] a) Barlow S, Marder SR. Nonlinear Optical Properties of Organic Materials. In: Functional Organic Materials; Müller, TJJ, Bunz UHF, Eds.; Wiley-VCH: Weinheim, Germany, 2007; p. 393-437. b) Szablewski M, Thomas PR, Thornton A, Bloor D, Cross GH, Cole JM, Howard JAK, Malagoli M, Meyers F, Brédas JL, Wenseleers W, Goovaerts E. Highly dipolar, optically nonlinear adducts of tetracyano-*p*-quinodimethane: synthesis, physical characterization, and theoretical aspects. J Am Chem Soc 1997:119;3144-54. c) Tykwinski RR, Gubler U, Martin RE, Diederich F, Bosshard C, Günter P. Structure–property relationships in third-order nonlinear optical chromophores. J Phys Chem B 1998;102:4451-65.

[7] a) Taniuchi T, Okada S, Nakanishi H. Widely tunable terahertz-wave generation in an organic crystal and its spectroscopic application. J Appl Phys 2004;95:5984-8. b) Taniuchi T, Ikeda S, Mineno Y, Okada S, Nakanishi H. Terahertz properties of a new organic crystal, 4'-dimethylamino-*N*-methyl-4-stilbazolium *p*-chlorobenzenesulfonate. Jpn J Appl Phys 2005;44:L932-4. c) Schneider A, Stillhart M, Günter P. High efficiency generation and detection of terahertz pulses using laser pulses at telecommunication wavelengths. Opt Express 2006;14:5376-84. d) Yang Z, Mutter L, Stillhart M, Ruiz B, Aravazhi S, Jazbinsek M, Schneider A, Gramlich V, Günter P. Large-size bulk and thin-film stilbazolium-salt single crystals for nonlinear optics and THz generation. Adv Funct Mater 2007;17:2018-23.

[8] a) He GS, Tan LS, Zheng Q, Prasad PN. Multiphoton absorbing materials: molecular designs, characterizations, and applications. Chem Rev 2008;108:1245-330. b) Malval JP, Jin

M, Morlet-Savary F, Chaumeil H, Defoin A, Soppera O, Scheul T, Bouriau M, Baldeck PL. Enhancement of the two-photon initiating efficiency of a thioxanthone derivative through a chevron-shaped architecture. Chem. Mater. 2011;23:3411-20. c) Kobayashi K, Kano SS. Multilayered optical storage with nonlinear read write. Opt Rev 1995;2:20-3.

[9] Paley MS, Harris JM, Looser H, Baumert JC, Bjorklund GC, Jundt D, Twieg RJ. A solvatochromic method for determining second-order polarizabilities of organic molecules. J Org Chem 1989;54:3774-8.

[10] a) Achelle S, Plé N. Pyrimidine ring as building block for the synthesis of functionalized Pi-conjugated materials. Curr Org Synth 2012;9:163-87. b) Achelle S, Baudequin C. Recent advances in pyrimidine derivatives as luminescent, photovoltaic and non-linear optical materials. Targets Heterocycl Systems 2013;17:under press.

[11] See for example: a) Achelle S, Ramondenc Y, Marsais F, Plé N. Star- and banana-shaped oligomers with a pyrimidine core: synthesis and light-emitting properties. Eur J Org Chem 2008;3129-40.b) Itami K , Yamazaki D, Yoshida Ji. Pyrimidine-core extended π -systems: general synthesis and interesting fluorescent properties. J Am Chem Soc 2004;126:15396-7. c) Bagley MC, Lin Z, Pope SJA. Barium manganate in microwave-assisted oxidation reactions: synthesis of solvatochromic 2,4,6-triarylpyrimidines. Tetrahedron Lett 2009;50:6818-22. d) Tumkevičius S, Voitechovičius A, Adoménas P, Synthesis of novel 2,4,6-triarylpyrimidines Chemija 2012;23:61-7.

[12] see for example a) Liu B, Hu XL, Liu J, Zhao YD, Huang ZL. Synthesis and photophysical properties of novel pyrimidine-based two-photon absorption chromophores. Tetrahedron Lett 2007;48:5958-62. b) Li L, Ge J, Wu H, Xu QH, Yao S Q. Organelle-specific detection of phosphatase activities with two-photon fluorogenic probes in cells and tissues. J Am Chem Soc 2012;134:12157-67. c) Achelle S, Malval JP, Aloïse S, Barsella A, Spangenberg A, Mager L, Akdas-Killig H, Fillaut JL, Caro B, Robin-le Guen F. Synthesis,

photophysics and nonlinear optical properties of stilbenoid pyrimidine-based dyes bearing methylenepyran donor groups. ChemphysChem 2013;14:2725-36.

[13] a) Achelle S, Barsella A, Baudequin C, Caro B, Robin-le Guen F. Synthesis and photophysical investigation of a series of push-pull arylvinyldiazine chromophores. J Org Chem 2012;77:4087-96. b) Castet F, Pic A, Champagne B, Linear and nonlinear optical properties of arylvinyldiazine dyes: a theoretical investigation. Dyes Pigm doi: 10.1016/j.dyepig.2014.03.021

[14] a) Bureš F, Pytela O, Diederich F. Solvent effects on electronic absorption spectra of donor-substituted 11,11,12,12-tetracyano-9,10-anthraquinodimethanes (TCAQs) J Phys Org Chem 2009;22:155-62. b) Bureš F, Pytela O, Kivala M, Diederich F. Solvatochromism as an efficient tool to study N,N-dimethylamino- and cyano-substituted π -conjugated molecules with an intramolecular charge-transfer absorption J Phys Org Chem 2011;24:274-81.

[15] a) Lartia R, Allain C, Bordeau G, Schmidt F, Fiorini-Debuisschert, Charra F, Teulade-Fichou MP. Synthetic strategies to derivatizable triphenylamines displaying high two-photon absorption. J Org Chem 2008;73:1732-44. b) Katan C, Terenziani F, Mongin O, Werts MHV, Porrès L, Pons T, Mertz J, Tretiak S, Blanchard-Desce M. Effects of (multi)branching of dipolar chromophores on photophysical properties and two-photon absorption. J Phys Chem A 2005;109:3024-3037. c) Detert H, Schmitt V. Quadrupolar donor-acceptor substituted oligo(phenylenevinylene)s-synthesis and solvatochromism of the fluorescence. J Phys Org Chem 2004;17:1051-6.

[16] a) Katritzky AR, Fara DC, Yang H, Tämm K. Quantitative measures of solvent polarity. Chem Rev 2004;104:175-98. b) Mancini PM, Adam CG, Fortunato GG, Vottero LR. A comparison of nonspecific solvent scales. Degree of agreement of microscopic polarity values obtained by different measurement methods. Arkivoc 2007;xvi:266-80.

[17] a) Reichardt C. Solvatochromic dyes as solvent polarity indicators. Chem Rev 1994;94:2319-58. b) Cerón-Carrasco JP, Jacquemin D, Laurence C, Planchat A, Reichardt C, Sraïdi K. Solvent polarity scales : determination of new $E_T(30)$ values for 84 organic solvents. J Phys Org Chem 2014;27:512-8.

[18] a) Taguchi G (ed.). System of experimental design. White Plains, NJ: Quality Resources Press;1987. b) Taguchi G (ed.). Taguchi methods: orthogonal arrays and linear graphs (softcover edn). Dearborn, Mich, USA: American Supplier Institute;1987. c) Ross PJ (ed.). Taguchi techniques for quality engineering (hardcover edn). City of New York, New York, USA: McGraw-Hill; 1996.

[19] See for example a) Pourjavadi A, Ayyari M, Amini-Fazl MS. Taguchi optimized synthesis of collagen-g-poly(acrylic acid)/kaolin composite superabsorbent hydrogel. Eur Polym J 2008;44:1209-16. b) Masoumi HRF, Kassim A, Basri M, Abdullah DK. Determining optimum conditions for lipase-catalyzed synthesis of triethanolamine (TEA)-based esterquat cationic surfactant by a Taguchi robust design method. Molecules 2011;16:4672-80. c) Drogat N, Barrière M, Granet R, Sol V, Krausz P. High yield preparation of purpurin-18 from *Spirulina maxima*. Dyes Pigm 2011;88:125-7.

[20] Kulhánek J, Bureš F, Pytela O, Mikysek T, Ludvík J, Růžička A. Puh-Pull molecules with a systematically extended π -conjugated system featuring 4,5-dicyanoimidazole. Dyes Pigm 2010;85:57-65.

[21] Chen L, Chu CI, Chen KJ, Chen PY, Hu SF, Liu RS. An intelligent approach to the discovery of luminescent materials using combinatorial approach combined with Taguchi methodology. Luminescence 2011;26:229-38.

[22] Zheng X, Song B, Xu B. Palladium-catalyzed regioselective C-H Bond orthoacetoxylation of arylpyridines 2010;4376-80.

[23] Pascal L, Vanden Eynde JJ, Van Haverbeke Y, Dubois P, Michel A, Rant U, Zojer E, Leising G, Van Dorn LO, Gruhn NE, Cornil J, Brédas JL. Synthesis and characterization of novel *para-* and *meta-*phenylenevinylene derivatives: fine tuning of the electronic and optical properties of conjugated materials. J Phys Chem B 2002;106:6442-50.

[24] Achelle S, Nouira I, Pfaffinger B, Ramondenc Y, Plé N, Rodríguez-López J. V-shaped 4,6-bis(arylvinyl)pyrimidine oligomers: synthesis and optical properties. J Org Chem 2009;74:3711-7.

Parameter	Levels		
	1	2	3
A: Nature of electron-	OMe	NMe ₂	NPh ₂
donating group			
B: Substituted positions of	2	4	4 and 6
the pyrimidine core.			
C: Nature of π -conjugated	₹-{\}-}	,	
linker		ş/	
			2
		A L	
L.			
\mathbf{C}			

Compound	A	В	С	D
1	1	1	1	1
2	1	2	2	2
3	1	3	3	3
4	2	1	2	3
5	2	2	3	1
6	2	3	1	2
7	3	1	3	2
8	3	2	1	3
9	3	3	2	1
				K
				Ĺ
)	

	UV/vis λ_{max} , nm	PL		Stokes shift
Compd ^a	$(\epsilon, M^{-1} \cdot cm^{-1})$	λ_{max} , nm	$\Phi_{ m F}{}^b$	cm^{-1}
1	283 (26600)	356	< 0.01	7245
2	338 (28500)	418	< 0.01	5662
3^b	373 (60500)	483	0.59	6106
4	251 (18900), 379 (31900)	472	0.01	5199
5 ^c	296 (24600), 386 (32400)	570	0.66	8362
6	299 (16000), 378 (38100)	429	0.72	3145
7	303 (38400), 373 (36500)	530	0.60	7942
8	<mark>300 (13000), 367 (31400)</mark>	482	<mark>0.55</mark>	<mark>6501</mark>
9 ^b	301 (38900), 427 (47600)	540	0.55	4901

^{*a*} All spectra were recorded in CH₂Cl₂ solutions at room temperature at $c = 1.0 \times 10^{-5}$ M to 3.0×10^{-5} M for absorption and $c = 1.0 \times 10^{-6}$ M to 3.0×10^{-6} M for emission. ^{*b*} Fluorescence quantum yield (± 10%) determined relative to quinine sulfate in 1 M H₂SO₄ ($\Phi_F = 0.54$) or harmane in 0.1 M H₂SO₄ as standard ($\Phi_F = 0.83$). ^{*b*} data from ref [24]. ^{*c*} data from ref [13a].

factor	\mathbf{S}^{a}	dof^b	\mathbf{V}^{c}	F-ratio ^d	Explained % ^e
A	<mark>5856</mark>	2	<mark>2928</mark>	<mark>35.6</mark>	<mark>48.0</mark>
В	<mark>3462</mark>	2	<mark>1731</mark>	<mark>21.1</mark>	<mark>28.4</mark>
С	<mark>2713</mark>	2	<mark>1356</mark>	<mark>16.5</mark>	<mark>22.2</mark>
Residue	<mark>164</mark>	2	<mark>82</mark>		1.3
Total	12281	8			100

^{*a*} Sum of squared deviations from the mean. ^{*b*} degree of freedom. ^{*c*} variance: mean of squares per dof. ^{*d*} ratio between squared deviation and the mean of square residue. ^{*e*} relative effects of the factors on λ_{abs} .

	A		В	}	C	2
	Average	Effect	Average	Effect	Average	Effect
1	<mark>331</mark>	<mark>-36</mark>	<mark>345</mark>	<mark>-22</mark>	<mark>343</mark>	<mark>-24</mark>
2	<mark>381</mark>	<mark>14</mark>	<mark>364</mark>	<mark>-3</mark>	<mark>381</mark>	<mark>14</mark>
3	<mark>389</mark>	<mark>22</mark>	<mark>393</mark>	<mark>26</mark>	<mark>377</mark>	10

	<i>n</i> -heptane	Toluene	THF	CH_2Cl_2	DMSO	
	$\Delta E_{\rm T}(30)^a = 0.0$	$\Delta E_{\rm T}(30)^a = 13.6$	$\Delta E_{\rm T}(30)^a = 27.2$	$\Delta E_{\rm T}(30)^a = 40.7$	$\Delta E_{\rm T}(30)^a = 59.0$	SP^b
	$\lambda_{em} (nm)$	$\lambda_{em} (nm)$	$\lambda_{em} (nm)$	$\lambda_{em} (nm)$	λ_{em} (nm)	
1	C		352	356	_ ^c	0.30
2	384	401	410	418	427	0.70
3	405	422	458	483	509	1.85
4	423	439	460	472	497	1.25
5	453	490	554	569	632	3.01
6	373	403	431	435	480	1.70
7	418	458	498	530	566	2.52
8	<mark>400</mark>	<mark>430</mark>	<mark>456</mark>	<mark>482</mark>	<mark>509</mark>	<mark>1.85</mark>
9	443	476	515	540	584	2.38
<mark>12</mark>	<mark>440</mark>	<mark>481</mark>	<mark>545</mark>	<mark>585</mark>	<mark>623</mark>	<mark>3.43</mark>

^{*a*} Dimroth–Reichardt polarity parameter, J·mol⁻¹. ^{*b*} SP: slope of the regression line λ_{em} (nm) versus $\Delta E_{T}(30)$ in nm mol J⁻¹. ^{*c*} due to very low emission intensity no emission maxima can be obtained.

	А		В	}	С	1
	Average	Effect	Average	Effect	Average	Effect
1	<mark>0.95</mark>	<mark>-0.78</mark>	<mark>1.36</mark>	<mark>-0.37</mark>	<mark>1.28</mark>	- <mark>0.45</mark>
2	<mark>1.99</mark>	<mark>0.26</mark>	<mark>1.85</mark>	<mark>0.12</mark>	<mark>1.44</mark>	<mark>-0.29</mark>
3	2.25	<mark>0.52</mark>	1.98	<mark>0.25</mark>	2.46	<mark>0.73</mark>

Factor	\mathbf{S}^{a}	dof^b	\mathbf{V}^{c}	F-ratio ^d	Explained % ^e
A	<mark>2.834</mark>	2	<mark>1.42</mark>	<mark>22.3</mark>	<mark>46.8</mark>
В	<mark>0.645</mark>	2	0.32	<mark>5.10</mark>	10.7
С	<mark>2.444</mark>	2	1.22	<mark>19.3</mark>	<mark>40.4</mark>
Residue	<mark>0.127</mark>	2	<mark>0.03</mark>		2.1
Total	<mark>6.051</mark>	8			100

^{*a*} Sum of squared deviations from the mean. ^{*b*} degree of freedom, ^{*c*} variance: mean of squares per dof. ^{*d*} ratio between squared deviation and the mean of square residue. ^{*e*} relative effects of the factors on SP.



R = electron-donating group















The photophysical properties of pyrimidine chromophores have been studied. The influence of three parameters has been studied by Taguchi methodology. Internal charge transfer estimated by emission solvatochromy has been optimized.

Prediction of photophysical properties of pyrimidine chromophores using Taguchi method.

Charline Denneval,¹ Sylvain Achelle,^{2*} Christine Baudequin,¹ and Françoise Robin-le Guen.²

¹ Normandie Univ. COBRA, UMR 6014 et FR 3038; Univ. Rouen; INSA Rouen; CNRS,

IRCOF, 1 rue Tesnière, 76821 Mont Saint Aignan Cedex, France

² Institut des Sciences Chimiques de Rennes UMR CNRS 6226, IUT de Lannion, rue Edouard Branly, BP 30219, F22302 Lannion Cedex, France

Corresponding author: sylvain.achelle@univ-rennes1.fr Tel: +33 (0)2 96 46 94 48 Fax : + 33 (0)2 96 46 93 54.

Electronic supporting informations

Effect of each level of each factor on λ_{em} in nm	S1
Analysis of variance (ANOVA) for on λ_{em} in nm.	S2
Calculated and experimental photophysical data (in CH ₂ Cl ₂) from literatu	re of some
pyrimidine chromophores.	S2
$\lambda_{em}~(nm)$ versus $\Delta E_T(30)~(J/mol)$ for compounds 1-9 and 12	S 3
Copy of ¹ H and ¹³ C NMR spectra of compounds 4, 7, 8 and 10-12	S6
References	S11

Table SI. EII	ect of each lev	vel of each f	actor on Λ_{em} in	nm		
¥.	А		E	8	С	
	Average	Effect	Average	Effect	Average	Effect
1	419	-57	453	-24	424	-52
2	492	16	490	14	477	0

3	517	41	486	10	528	51

Factor	S ^a	dof ^b	V ^c	F-ratio ^d	Explained % ^e
А	15672	2	7836	13.2	44.3
В	2520	2	1260	2.13	7.1
С	16017	2	8009	13.5	45.3
Residue	1183	2	591		3.3
Total	35394	8			100

Table S2. Analysis of variance (ANOVA) for on λ_{em} in nm.

^a Sum of squared deviations from the mean ^b degree of freedom ^c variance: mean of squares per dof ^dratio between squared deviation and the mean of square residue ^e relative effects of the factors on λ_{em} .

Table S3. Calculated and experimental photophysical data (in CH₂Cl₂) from literature of some pyrimidine chromophores.¹

	Calculated	Calculated
Compounds	λ_{abs} in nm	λ_{em} in nm
	(experimental	(experimental
	λ_{abs} in nm)	λ_{em} in nm)
Me ₂ N NMe ₂	421 (429)	502 (530)
MeO OMe	371 (353)	429 (406)
Ph ₂ N NPh ₂	429 (427)	527 (540)

Me ₂ N NMe ₂	417 (407)	553 (592)
OMe	342 (338)	433 (418)
N N N NMe ₂	392 (393)	506 (488)
N N N NPh ₂	400 (400)	531 (526)
	338 (346)	484 (447)
N N N NMe ₂	388 (386)	557 (570)
N N N NPh ₂	396 (384)	582 (563)





















¹ a) Achelle S, Barsella A, Baudequin C, Caro B, Robin-le Guen F. Synthesis and photophysical investigation of a series of push-pull arylvinyldiazine chromophores. J Org Chem 2012;77:4087-96. b) Achelle S, Nouira I, Pfaffinger B, Ramondenc Y, Plé N, Rodríguez-López J. V-shaped 4,6-bis(arylvinyl)pyrimidine oligomers: synthesis and optical properties. J Org Chem 2009;74:3711-7.