

High Oxidation State Alkyl and Carbene Complexes of Molybdenum Containing the *t*-Butylimido Ligand

Didier Ehrenfeld, Jacky Kress, Barry D. Moore, John A. Osborn,* and Guy Schoettel


UA au CNRS n° 424, Institut Le Bel, Université Louis Pasteur, 4, rue Blaise Pascal, 67000 Strasbourg, France

Alkylation reactions of imido-oxo complexes of molybdenum(vi) lead to the formation of various imido-alkyl and imido-carbene compounds.

High oxidation state alkyl and carbene chemistry of molybdenum¹ has been almost totally neglected when compared with that of tungsten. This is somewhat surprising since molybdenum to carbon bonds may be involved in several important catalytic oxidation processes (*e.g.* conversion of propene into acrolein or acrylonitrile²) as well as in olefin^{1b,3} metathesis. The presence of oxo and/or imido ligands on Mo is also of considerable interest both as possible reactive⁴ and stabilising or activating⁵ units. In recent years a number of interesting imido complexes of Mo with ancillary ligands such as dithiocarbamate,^{4a} phosphine, and cyclopentadienyl⁶ groups have thus been synthesised. We present now the convenient synthesis of new oxo-imido and imido complexes of molyb-

denum(vi), including several containing molybdenum to carbon single and double bonds (Scheme 1).

Compound (1)⁷ when treated with RNCO (1 equiv.; R = But^t) yields CO₂ and {MoO(NR)Cl₂(MeCN)}_n (2) as an orange powder in quantitative yield. We suggest an oxo-bridged structure for (2) ($\nu_{\text{Mo-O-Mo}}$ 720 cm⁻¹), probably trimeric as found for {W(μ -O)(NPh)Me₂(PMe₃)₃}₃,⁸ although a dimeric form cannot be excluded. Complex (2) reacts with 2 equiv. of Li(OCH₂R) to give LiCl and the yellow-green crystalline product MoO(NR)(OCH₂R)₂ (3). The $\nu_{\text{Mo-O}}$ frequency at 850 cm⁻¹ shows (3) to be possibly monomeric [*cf.* W(NR)₂(OR)₂]⁹, although bridging may occur *via* the imido ligands [see {Mo(NR)₂Me₂}₂]^{1c}].



Scheme 1.† ($\text{R} = \text{Bu}^t$). *Reagents* (25 °C): i, RNCO (1 equiv.), MeCN , 48 h (ca. 100%); ii, $\text{Li}(\text{OCH}_2\text{R})$ (2 equiv.), Et_2O , 1 h (80%); iii, $(\text{RCH}_2)_2\text{Mg}$ -dioxane (1.1 equiv.), Et_2O , 1 h (35%); iv, $(\text{RCH}_2)_2\text{Li}$ (1 equiv.), C_6H_6 , 10 min (75%); v, $(\text{RCH}_2)_2\text{Mg}$ -dioxane (1 equiv.), Et_2O , 2 h (75%); vi, $\text{C}_6\text{F}_5\text{OH}$ (1 equiv.), pentane, 10 min; vii, Ph_3SiOH (1 equiv.), pentane, 1 h.

The oxo ligand in (2) and (3) is surprisingly reactive. If alkylation of (2) is carried out with $(\text{RCH}_2)_2\text{Mg}$ (1.1 equiv.), $\text{Mo}(\text{NR})(\text{CH}_2\text{R})_3\text{Cl}$ (4) is obtained as colourless crystals.

† *Spectroscopic data*: ^1H (200 MHz) and ^{13}C (50 MHz) n.m.r. δ , C_6D_6 , J in Hz; i.r. in Nujol mull, ν in cm^{-1} ; eq. = equatorial, ax. = axial.

(2) $\{\text{MoO}(\text{NCMe}_3)\text{Cl}_2(\text{MeCN})\}_3$: ^1H n.m.r. 2.3 (s, 3H, MeCN), 1.6 (s, 9H, NCMe_3); i.r. 720 ($\nu_{\text{Mo}-\text{O}-\text{Mo}}$), 330 ($\nu_{\text{Mo}-\text{Cl}}$), 2280, 2305 ($\nu_{\text{C}=\text{N}}$). (3) $\text{MoO}(\text{NCMe}_3)(\text{OCH}_2\text{CMe}_3)_2$: ^1H n.m.r. 4.90 (s, 4H, OCH_2CMe_3), 1.50 (s, 9H, NCMe_3), 1.20 (s, 18H, OCH_2CMe_3); i.r. 850 ($\nu_{\text{Mo}-\text{O}}$), 670 ($\nu_{\text{Mo}-\text{O}}$).

(4) $\text{Mo}(\text{NCMe}_3)(\text{CH}_2\text{CMe}_3)_3\text{Cl}$: ^1H n.m.r. 2.97 (s, 6H, CH_2CMe_3), 1.41 (s, 9H, NCMe_3), 1.28 (s, 27H, CH_2CMe_3); $^{13}\text{C}\{^1\text{H}\}$ n.m.r. 77.8 (CH_2CMe_3), 34.0 (CH_2CMe_3), 28.0 (NCMe_3); i.r. 260 ($\nu_{\text{Mo}-\text{Cl}}$).

(5) $\text{Mo}(\text{NCMe}_3)(\text{CHCMe}_3)(\text{CH}_2\text{CMe}_3)_2$: ^1H n.m.r. 9.22 (s, 1H, CHCMe_3), 1.72 [d, 2H, $^2J(\text{H}_\text{A}\text{H}_\text{B})$ 11, $\text{CH}_\text{A}\text{H}_\text{B}\text{CMe}_3$], 1.56 (s, 9H, NCMe_3), 1.39 (s, 9H, CHCMe_3), 1.32 [d, 2H, $^2J(\text{H}_\text{A}\text{H}_\text{B})$ 11, $\text{CH}_\text{A}\text{H}_\text{B}\text{CMe}_3$], 1.26 (s, 18H, CH_2CMe_3); ^{13}C n.m.r. 249.3 [d, $J(\text{CH})$ 106, CHCMe_3], 73.9 (t , CH_2CMe_3), 70.7 (s, NCMe_3), 43.3 (s, CHCMe_3), 34.6 (q, CH_2CMe_3), 32.6 and 32.5 (q, NCMe_3 and CHCMe_3).

(6) $\text{Mo}(\text{NCMe}_3)(\text{CH}_2\text{CMe}_3)_2(\text{OCH}_2\text{CMe}_3)_2$: ^1H n.m.r. 4.49 (s, 2H, OCH_2CMe_3 ax.), 4.33 (s, 2H, OCH_2CMe_3 eq.), 2.88 [d, 2H, $^2J(\text{H}_\text{A}\text{H}_\text{X})$ 7, $\text{CH}_\text{A}\text{H}_\text{X}\text{CMe}_3$], 2.18 [d, 2H, $^2J(\text{H}_\text{A}\text{H}_\text{X})$ 7, $\text{CH}_\text{A}\text{H}_\text{X}\text{CMe}_3$], 1.55 (s, 9H, NCMe_3), 1.38 (s, 9H, OCH_2CMe_3 ax.), 1.36 (s, 18H, CH_2CMe_3), 1.06 (s, 9H, OCH_2CMe_3 eq.); $^{13}\text{C}\{^1\text{H}\}$ n.m.r. 94.5 (OCH_2CMe_3 ax.), 81.3 (OCH_2CMe_3 eq.), 74.2 (CH_2CMe_3), 35.8 (NCMe_3), 35.2 (OCH_2CMe_3 ax.), 34.3 (CH_2CMe_3), 32.6 (OCH_2CMe_3 eq.), 31.0 (NCMe_3), 28.0 (OCH_2CMe_3 ax.), 26.7 (OCH_2CMe_3 eq.).

(7) $\text{Mo}_2\text{O}(\text{NCMe}_3)_2(\text{CHCMe}_3)(\text{CH}_2\text{CMe}_3)_4$: ^1H n.m.r. 11.71 (s, 1H, CHCMe_3), 2.65 [d, 1H, $^2J(\text{H}_\text{A}\text{H}_\text{B})$ 9, $\text{CH}_\text{A}\text{H}_\text{B}\text{CMe}_3$], 2.57 (d, 3H, $^2J(\text{H}_\text{A}\text{H}_\text{B})$ 7, $\text{CH}_\text{A}\text{H}_\text{B}\text{CMe}_3$], 2.46 [d, 3H, $^2J(\text{H}_\text{A}\text{H}_\text{B})$ 7, $\text{CH}_\text{A}\text{H}_\text{B}\text{CMe}_3$], 2.17 [d, 1H, $^2J(\text{H}_\text{A}\text{H}_\text{B})$ 9, $\text{CH}_\text{A}\text{H}_\text{B}\text{CMe}_3$], 1.67 (s, 9H, NCMe_3), 1.60 (s, 9H, NCMe_3), 1.53 (s, 9H, CHCMe_3), 1.46 (s, 9H, CH_2CMe_3), 1.29 (s, 27H, CH_2CMe_3); $^{13}\text{C}\{^1\text{H}\}$ n.m.r. 269.2 (CHCMe_3), 73.7 (CH_2CMe_3), 57.0 (CH_2CMe_3), 34.5 (CH_2CMe_3), 33.1 (CMe_3), 29.0 (CMe_3); i.r. 770, 720 ($\nu_{\text{Mo}-\text{O}-\text{Mo}}$).

(8) $\text{Mo}(\text{NCMe}_3)(\text{CH}_2\text{CMe}_3)_3(\text{OC}_6\text{F}_5)$: ^1H n.m.r. 2.62 (s, 6H, CH_2CMe_3), 1.50 (s, 9H, NCMe_3), 1.17 (s, 27H, CH_2CMe_3).

(9) $\text{Mo}(\text{NCMe}_3)(\text{CHCMe}_3)(\text{CH}_2\text{CMe}_3)(\text{OSiPh}_3)$: ^1H n.m.r. 11.46 (s, 1H, CHCMe_3), 7.9, 7.3 (m, 15H, OSiPh_3), 2.42 [d, 2H, $^2J(\text{H}_\text{A}\text{H}_\text{B})$ 11, $\text{CH}_\text{A}\text{H}_\text{B}\text{CMe}_3$], 2.30 [d, 2H, $^2J(\text{H}_\text{A}\text{H}_\text{B})$ 11, $\text{CH}_\text{A}\text{H}_\text{B}\text{CMe}_3$], 1.36 (s, 9H, NCMe_3), 1.32 (s, 9H, CHCMe_3), 1.26 (s, 9H, CH_2CMe_3).

Spectroscopic data indicate (4) to be isostructural with $\text{MoO}(\text{CH}_2\text{R})_3\text{Cl}^{1b}$ and $\text{W}(\text{NR}')(\text{CH}_2\text{R})_3\text{Cl}^{10,11}$ ($\text{R}' = \text{Me}$ or Ph) previously described. Treatment of (4) with 1 equiv. of RCH_2Li yields $\text{Mo}(\text{NR})(\text{CHR})(\text{CH}_2\text{R})_2$ (5), the first simple alkylidene complex of molybdenum in a high oxidation state. Complex (5) is obtained as a brown oil (at 25 °C) after purification by short-path distillation (40 °C; 10^{-4} mm Hg) and is characterised in particular by the carbene n.m.r. signals at δ 9.22 (^1H) and 249.5 (^{13}C). The observation by ^1H n.m.r. spectroscopy of two equivalent neopentyl ligands possessing two diastereotopic methylenic protons allows us to ascribe to (5) the structure shown in Scheme 1. A single isomer is indicated, although the n.m.r. spectra would also be consistent with the presence of two isomers equilibrating rapidly by rotation of the carbene ligand about the molybdenum-carbon double bond. Such a rotation process is slow however for analogous tungsten carbene compounds which have recently been described.¹² Alkylation of (3) with $(\text{RCH}_2)_2\text{Mg}$ (1 equiv.) yields $\text{Mo}(\text{NR})(\text{CH}_2\text{R})_2(\text{OCH}_2\text{R})_2$ (6) as brown crystals which can be purified by recrystallisation from pentane or sublimation (70 °C; 10^{-4} mm Hg). ^1H and ^{13}C N.m.r. data clearly indicate the structure shown in Scheme 1. Using 1.5 equiv. of $(\text{RCH}_2)_2\text{Mg}$ leads to formation of $\text{Mo}(\text{NR})(\text{CH}_2\text{R})_3(\text{OCH}_2\text{R})$, an alkoxyated derivative of (4), as well as variable yields of an unusual carbene complex (7) of formula $\text{Mo}_2\text{O}(\text{NR})_2(\text{CHR})(\text{CH}_2\text{R})_4$. This latter yellow crystalline complex shows in particular n.m.r. signals of the carbene ligand at δ 11.71 (^1H) and 269.2 (^{13}C), as well as two non-equivalent NR ligands and two types of $-\text{CH}_2\text{R}$ groups of relative ratio 3:1. The i.r. spectrum shows the probable presence of an Mo-O-Mo linkage, and although certain dimeric structures can be proposed for (7), details must await an X-ray determination.

These carbene complexes undergo some interesting reactions. For example, $\text{C}_6\text{F}_5\text{OH}$ reacts rapidly with (5) to yield $\text{Mo}(\text{NR})(\text{CH}_2\text{R})_3(\text{OC}_6\text{F}_5)$ (8) resulting from addition of $\text{C}_6\text{F}_5\text{O}-\text{H}$ across the metal-carbon double bond. By contrast Ph_3SiOH reacts to form the new carbene complex $\text{Mo}(\text{NR})(\text{CHR})(\text{CH}_2\text{R})(\text{OSiPh}_3)$ (9), in which a neopentyl group in (5) is replaced by a siloxo ligand. Like their tungsten analogues,¹² certain of these complexes metathesise olefins

without a Lewis acid cocatalyst. These results will be the subject of a separate publication.

Received, 11th September 1986; Com. 1304

References

- 1 For molybdenum(vi) alkyl chemistry see (a) D. N. Clark and R. R. Schrock, *J. Am. Chem. Soc.*, 1978, **100**, 6774; (b) J. Kress, M. J. M. Russell, M. G. Wesolek, and J. A. Osborn, *J. Chem. Soc., Chem. Commun.*, 1980, 431; (c) W. A. Nugent and R. L. Harlow, *J. Am. Chem. Soc.*, 1980, **102**, 1759; (d) G. N. Schrauzer, L. A. Hughes, P. R. R. Strampach, and E. O. Schlemper, *Organometallics*, 1982, **1**, 44. For molybdenum carbene chemistry see (e) H. Arzoumanian, A. Baldy, R. Lai, J. Metzger, M. L. Nkeng Peh, and M. Pierrot, *J. Chem. Soc., Chem. Commun.*, 1985, 1151.
- 2 J. D. Burrington, C. T. Kartisek, and R. K. Grasselli, *J. Catal.*, 1983, **81**, 489, and references therein.
- 3 K. Seyferth and R. Taube, *J. Mol. Catal.*, 1985, **28**, 53.
- 4 (a) W. A. Nugent and B. L. Haymore, *Coord. Chem. Rev.*, 1980, **31**, 123, and references therein; (b) D. M. T. Chan and W. A. Nugent, *Inorg. Chem.*, 1985, **24**, 1422.
- 5 A. K. Rappe and W. A. Goddard, *J. Am. Chem. Soc.*, 1982, **104**, 448.
- 6 E.g. (a) C. Y. Chou, J. C. Huffman, and E. A. Maatta, *Inorg. Chem.*, 1986, **25**, 822; (b) M. L. H. Green and K. J. Moynihan, *Polyhedron*, 1986, **5**, 921.
- 7 R. J. Butcher, J. P. Guntz, R. G. A. R. MacLagan, H. K. J. Powell, C. J. Wilkins, and Y. S. Hian, *J. Chem. Soc., Dalton Trans.*, 1975, 1223.
- 8 D. C. Bradley, M. B. Hursthouse, K. M. A. Malik, and A. J. Nielson, *J. Chem. Soc., Chem. Commun.*, 1981, 103.
- 9 W. A. Nugent, *Inorg. Chem.*, 1983, **22**, 965.
- 10 S. F. Pedersen and R. R. Schrock, *J. Am. Chem. Soc.*, 1982, **104**, 7483.
- 11 J. Kress, M. Wesolek, J. P. Le Ny, and J. A. Osborn, *J. Chem. Soc., Chem. Commun.*, 1981, 1039.
- 12 C. J. Schaverien, J. C. Dewan, and R. R. Schrock, *J. Am. Chem. Soc.*, 1986, **108**, 2771.