



Communication

Subscriber access provided by UB + Fachbibliothek Chemie | (FU-Bibliothekssystem)

Total Synthesis and Structural Revision of (+)-Muironolide A

Qing Xiao, Kyle Young, and Armen Zakarian J. Am. Chem. Soc., Just Accepted Manuscript • Publication Date (Web): 30 Apr 2015 Downloaded from http://pubs.acs.org on May 1, 2015

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

1 2

Total Synthesis and Structural Revision of (+)-Muironolide A

Qing Xiao,^{†,‡} Kyle Young,[†] and Armen Zakarian*

Department of Chemistry and Biochemistry, University of California, Santa Barbara, California 93106, USA.

Supporting Information Placeholder

ABSTRACT: Muironolide A is a fascinating tetrachlorinated marine polyketide isolated from the sponge of *Phorbas* sp. Only 90 μ g had been isolated, and the structure was established by nanoscale NMR techniques. Herein we report the total synthesis of the substance with the assigned structure of muironolide A, propose a revised structure based on NMR data, and complete the enantioselective total synthesis of muironolide A.

In 2009, Molinski and co-workers reported the structure of a remarkable marine natural product, muironolide A (1),¹ isolated from the sponge of the Phorbas species, the same specimen that earlier provided phorboxazoles A and B.² With only 90 µg (152 nmol) available, the complete structure of muironolide A was determined using recently developed nanoscale NMR techniques³ as well as degradation chemistry with 30 µg of the material. The structure of muironolide A represents a new chemical entity, the main features of which include a 16membered diester lactone with an unprecedented hexahydro-1H-isoindolone, a trichlorocarbinol ester, and chlorocyclopropane subunits. Given the extremely small amounts of this "nearly extinct" natural product, only fragmented studies of its bioactivity were possible, which, nevertheless, identified cytotoxic activity against the HCT-116 solid colon tumor cell line (IC₅₀ 96.5 μ g/mL) and antifungal activity against Cryptococcus neoformans (MIC 16 μ g/mL).

Herein we report the enantioselective total synthesis of compound **1** with the originally assigned structure of muironolide A along with its C21,C22,C23-diastereomer **2** (Figure 1). Based on ¹H and ¹³C NMR data from these two compounds, none of which matched those of the natural substance, we made a projection for the corrected structure. This structure was also accessed by total synthesis, and characterization data for this compound were found to be fully consistent with those of the natural product, resulting in the reassignment of stereochemistry at C21 and establishing the absolute stereochemistry as the enantiomer of **3**. Although the initially assigned structure did not fully match the data¹ for the natural product, inconsistencies appear to be confined to the fragment produced by the degradation chemistry, thus underscoring the power of the nanoscale NMR methods for determining the structure of this exceedingly rare complex natural product.



Figure 1. Original and revised structures of muironolide A

The key features of our synthesis design are outlined in Scheme 1.⁴ The design calls for a late stage macrolactone formation; the specific order of esterification was intentionally left flexible for optimization. A central transformation in this plan is the *exo*-selective lanthanide-catalyzed intramolecular Diels-Alder reaction for the construction of the hexahydro-1*H*-isoindolone subunit.^{5,6} In this reaction, the enol form of the γ , δ -unsaturated β -keto amide precursor **3** serves as the dienophile, which is further activated by raising the HOMO of the diene after deprotonation in the presence of a lanthanide additive, giving rise to enolate chelate **i**.

Scheme 1.



To access the β -keto amide precursor of the IMDA reaction, (+)- β -citronellene was first converted to trichlorocarbinol 4 by selective ozonolysis⁷ and reaction of the resultant aldehyde with Me₃SiCCl₃ in the presence of sodium formate.⁸ To set the configuration at C17, an oxidation-reduction sequence was required. Oxidation of 4 with DMSO/(CF₃CO)₂O (80% yield) followed by reducthe trichloromethyl ketone tion of with $[Ru(cymene)Cl]_2$ and (R,R)-TsDPEN under transfer hydrogenation conditions delivered 5 in 97% yield and 10:1 diastereoselectivity.9 After benzyloxymethylation (BOMCl, *i*-Pr₂NEt, *n*-Bu₄NI, CH₂Cl₂, 45 °C, 12 h, 92% yield), cross-metathesis with methacrolein in the presence of 3 mol % of Hoveyda-Grubbs II catalysts (HGII) afforded aldehyde **6** in 67% yield as a 10:1 mixture of *E* and *Z* alkene isomers. The reaction was accompanied by the recovery of ~20% of the starting material. Olefination with dioxinone phosphonate **7** readily afforded **8**, which, upon thermolysis with amine **9**⁵ generated amide **10** in 93% yield. Cross-metathesis with methyl acrylate or functionalized acrylate **11**^{10,11,12} (7 mol % HGII, CH₂Cl₂, 45 °C, 12 h) completed the synthesis of IMDA precursors **3a** and **3b**.

Preliminary investigations demonstrated the feasibility of the planned IMDA reaction for the assembly of the hexahydro-1H-isoindolone ring system of muironolide A.⁵ When the substrate possesses the Z geometry at the C4–C5 double bond, as in 3, the requisite relative stereochemistry of all newly formed stereogenic centers is achieved. The IMDA reaction can be performed directly by thermal activation or under catalytic conditions.¹³ We found that the thermal cycloaddition can be performed simply by heating the substrate at 110 °C in toluene, with no base additive required. It was also characterized by a very high level of diastereocontrol favoring the exo-IMDA product (>30:1). Together, these observations are indicative of a polar concerted mechanism rather than a double Michael addition pathway.

The reactive hydroxy diene intermediate is accessed from the β -keto amide precursor through ketone-enol tautomerization (Scheme 3). We hypothesize that the exo-IMDA pathway is strongly favored due to hydrogen-bonding stabilization in the transition structure TS1 (X = H). In the transition structure for the disfavored endo-IMDA pathway the hydrogen bond is disrupted (TS2, X = H). Furthermore, replacing the hydrogen bond in **TS1** with a chelated metal enolate (TS1, X = M) was expected to increase electron density in the diene system, raising its HOMO and activating the diene for cycloaddition. As anticipated, such stabilized enolates could be generated catalytically via soft enolization, and a screen of Lewis acid additives revealed that lanthanide triflates or nitrates were effective catalysts.¹³

Scheme 3.

1

2

3

ACS Paragon Plus Environment

1

2

3

4



In the event, when **3b** was treated with 10 mol % of La(OTf)₃, 12 mol % of PYBOX ligand 12, and triethylamine in ethyl acetate at 45 °C, exo-IMDA product 13 was obtained as an inseparable 3:1 mixture with its C4,C5,C8,C11-diastereomer in 61% combined yield. Reduction of 13 with NaBH₄ afforded a mixture of two isomeric products which now could be readily separated by preparative HPLC. The major product (14) afforded high quality single crystals suitable for X-ray crystallographic analysis, which, critically, ascertained the configuration of all stereogenic centers of the target natural product. The functionalization of the hexahydro-1*H*-isoindolone ring system was concluded by dehydration with DCC and CuCl upon heating in toluene, giving **15** in high yield.¹⁴ Of note, other methods (MsCl, Et₃N, then DBU; Burgess or Martin reagents) afforded the deconjugated C9-C10 alkene as the sole product.

Scheme 4.



Simultaneous removal of the *tert*-butyl ester and benzyloxymethyl groups was achieved effectively by exposure of **15** to trifluoroacetic acid in CH₂Cl₂ at room temperature. Of various attempted methods for macrolactonization of *seco*-acid **16**, the Yamaguchi reagent¹⁵ appeared most promising. However, even under optimal conditions, hydroxyl acid **17** was isolated as a major product, apparently arising from β -elimination in the chlorocyclopropyl acid subunit. Other products resulted from higher order macrolactonizations, and the amount of the intended product was low at best. Therefore, the alternative order of ester installations for the assembly of the 16-membered macrolactone was tested.

Using the same condition for the IMDA reaction of **3a** (10 mol % La(OTf)₃, 12 mol % ligand **12**, Et₃N, EtOAc, 45 °C, 24 h), isoindolone **18** was isolated in 61% yield, again as a 3:1 mixture with its diastereomer (Scheme 5). After reduction with sodium borohydride and purification by preparative HPLC, dehydration of **20**¹⁶ followed by removal of the BOM group (trifluoroacetic acid, CH₂Cl₂, 0 °C, 1 h) readily afforded trichlorocarbinol **22**.

Esterification of **22** with carboxylic acid **23**¹⁰ to ester 24 was best achieved using the Yamaguchi conditions (97% yield) (Scheme 6). Both the methyl ester and the triethylsilyl ether in 24 could be cleaved in one step under nucleophilic conditions with microwave irradiation. In contrast to intermediate 16, the resulting seco acid 25 underwent a rather clean macrolactonization upon mild heating in toluene with 2,4,6-trichlorobenzoyl chloride, DMAP, and pyridine, affording 26 in 55% yield. After careful optimization, we found that the final amide deprotection is ideally performed under oxidative conditions upon heating at 100 °C with DDQ in dioxane with controlled water content of 0.09% (90% yield).17 The 1H and 13C NMR data of the resulting compound did not match those reported for the natural product.

Scheme 5.

5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

Journal of the American Chemical Society



Because carboxylic acid **23** has its origin in an enzymatic resolution process,10 its enantiomer ent- was also readily available. Replicating the same sequence of reactions that delivered **1** with *ent-23* afforded another isomer of muironolide A, compound 2. Its NMR data also did not match the natural substance; however, chemical shift analysis of ¹H and ¹³C NMR spectra revealed a markedly closer correlation (Figure 2). As can be seen from the analysis of ¹³C NMR spectra, the major differences are confined to the C21-C26 region, i.e. mostly to the *trans*-chlorocyclopropyl ketide (CCK) unit. Originally, the configuration of this section of the muironolide A could not be determined spectroscopically and required an audacious effort that combined degradation chemistry with 30 µg of the natural sample, chemical correlation, and Mosher ester analysis. In our work, we ascribed a much closer correlation of NMR data between the natural product and 2, vs. 1, to the inverted configuration at C21, which we presumed to have a larger impact on the macrocycle conformation vis-a-vis the exocvclic CCK unit. On the basis of these conjectures. as well as an alternative interpretation of HPLC correlation data in the original report, we hypothesized that muironolide A is the C21 epimer of 1.

This hypothesis was put to test when **22** had been converted to the macrolactone using CCK acid **24**¹⁰ and the same four-step sequence of reactions (Scheme 7). The physical data for the resulting compound were in full agreement with those reported for natural muironolide A; however, circular dichroism (CD) spectroscopy demonstrated that it is enantiomeric to the natural sample,¹⁰ thus establishing its absolute configuration as *opposite* to that of **3**.

Scheme 6.



Figure 2. Differences in the ¹³C NMR chemical shifts between (a) **1**, (b) **2**, and (c) **3** (synthetic) and natural muironolide A.

1

2



The enantioselective total synthesis of (+)muironolide A resulted in the adjustment of configuration at C21 and reassignment of the absolute configuration of the natural product. Delivering 25 mg of the natural product in the first unoptimized effort, over 250 times the amount isolated from the natural sources, it paves a way for a more systematic evaluation of the biological profile of muironolide A. Perhaps more importantly, this work is a highlight of the powerful combination of nanoscale NMR analysis with total synthesis for uncovering the full potential of "nearly extinct", exceedingly rare components of natural product extracts.

Corresponding Author

zakarian@chem.ucsb.edu

Present Addresses

*Boehringer Ingelheim Shanghai Pharmaceuticals Co., Ltd., No. 1010 Longdong Avenue, Zhangjiang, Shanghai 201203, China

Author Contributions

[†] authors contributed equally.

Supporting Information

Detailed experimental procedures, copies of NMR spectra, CD measurements. This material is available free of charge via the Internet at http://pubs.acs.org.

ACKNOWLEDGMENT

We thank Professor Molinski for insightful discussions and for performing the CD measurements of **3**. This work is supported by the NSF CHE-1463819, Amgen, and Eli Lilly. K.Y. is supported by an NSF Graduate Fellowship (DGE-1144085). Dr. Guan Wu is thanked for assistance with X-ray crystallography.

References

Dalisay, D. S.; Morinaka, B. I.; Skepper, C. K.; Molinski, T. F. J. Am. Chem. Soc. 2009, 131, 7552.

² (a) Searle, P. A.; Molinski, T. F. J. Am. Chem. Soc. 1995, 117, 8126. (b) Searle, P. A.: Molinski, T. F.: Brzezinski, L. J.: Leahy, J. W. J. Am. Chem. Soc. 1996, 118, 9422. (c) Molinski, T. F. Tetrahedron Lett. 1996, 37, 7879.

³ Molinski, T. F. Nat. Prod. Rep. 2010, 27, 321.

⁴ The protecting groups were defined after preliminary experimentation.

⁵ Xiao, Q.; Young, K.; Zakarian, A. Org. Lett. **2013**, *15*, 3314.

⁶ (a) Flores, B.; Molinski, T. F. Org. Lett. **2011**, 13, 3932. (b) Shaner, C. E.; Ferrence, G. M.; Mitchell, A. T. Synlett. 2013, 24, 1861.

Fürstner, A.; Feyen, F.; Prinz, H.; Waldmann, H.; Angew. Chem. Int. Ed. 2003, 42, 5361.

Kister, J.; Mioskowski, C. J. Org. Chem. 2007, 72, 3925.

⁹ Perryman, M. S.; Harris, M. E.; Foster, J. L.; Joshi, A.; Clarkson, G. J.; Fox. D. J. Chem. Commun. 2013, 49, 10022.

¹⁰ See supporting information for its synthesis.

¹¹ (a) Tan, C-H.; Holmes, A. B. Chem. Eur. J. 2001, 7, 1845. (b) Araoz, R.; Servent, D.; Molgó, J.; Iorga, B. I.; Fruchart-Gaillard, C.; Benoit, E.; Gu, Z.; Stivala, C.; Zakarian, A. J. Am. Chem. Soc. 2011, 133, 10499. (c) Lu, C.-D.; Zakarian, A. Org. Lett. 2007, 9, 3161.

¹² Beaulieu, L-P, B.; Zimmer, L. E.; Gagnon, A.; Charette, A. B. Chem. Eur. J. 2012, 18, 14784.

¹³ Young, K.; Xiao, Q.; Zakarian, A. Eur. J. Org. Chem. 2015,

2337. ¹⁴ Liu, Y.; Wang, Q.; Zhang, Q.; Huang, J.; Nie, L.; Chen, J.; Cao, W.; Wu, X. J. Org. Chem. 2013, 78, 12009.

¹⁵ Parenty, A.; Moreau, X.; Niel, G.; Campagne, J. M. Chem. Rev. 2013, 113, PR1-PR40.

¹⁶ The structure of **20** was confirmed by chemical correlation to 14

Palmer, C. F.; Parry, P. K.; Roberts, S. M.; Sik, V. J. Chem. Soc., Perkin Trans. 1, 1992, 8, 1021.

