


Electrochemistry Hot Paper


Electrooxidative Rhodium-Catalyzed [5+2] Annulations via C–H/O–H Activations

Yulei Wang, João C. A. Oliveira, Zhipeng Lin, and Lutz Ackermann*

Abstract: Electrooxidative annulations involving mild transition metal-catalyzed C–H activation have emerged as a transformative strategy for the rapid construction of five- and six-membered heterocycles. In contrast, we herein describe the first electrochemical metal-catalyzed [5+2] cycloadditions to assemble valuable seven-membered benzoxepine skeletons by C–H/O–H activation. The efficient alkyne annulation featured ample substrate scope, using electricity as the only oxidant. Mechanistic studies provided strong support for a rhodium(III/I) regime, involving a benzoxepine-coordinated rhodium(I) sandwich complex as the catalyst resting state, which was re-oxidized to rhodium(III) by anodic oxidation.

Based on major achievements in the C–H activation arena during the past two decades, transition metal-catalyzed annulations involving the activation of otherwise unreactive C–H bonds have revolutionized the art of preparing cyclic compounds.^[1–4] Despite indisputable advances, sacrificial chemical oxidants, such as Cu(OAc)₂ and AgOAc, are generally required to facilitate these processes, thus resulting in the generation of undesired byproducts and reducing the atom economy.

Electricity has been considered as a green and atom-economic redox equivalent.^[5,6] Significant recent momentum has been gained by the merger of electrocatalysis with organometallic C–H activation.^[7–11] These reactions have provided efficient routes for the assembly of a variety of heterocycles, normally five- and six-membered rings through formal [3+2]^[9] or [4+1]^[10] or [4+2]^[11] cycloadditions, respectively, with major contributions by the groups of Mei, Lei, Xu, and Ackermann, among others (Scheme 1 b). However, while seven-membered rings, such as benzoxepine derivatives, are the core structures of many natural products and pharmacologically relevant molecules (Scheme 1 a),^[12] the construction of these scaffolds by means of metallaelectro-catalyzed



Scheme 1. Electrochemical metal-catalyzed C–H annulation.

annulations has proven elusive. Moreover, while rhodium catalysts have been widely used in C–H activation, the key low-valent rhodium(I) intermediates could seldom be isolated and their redox-chemistry was rarely studied by electroanalysis. Within our program on sustainable C–H activation,^[13] we herein report on a uniquely efficient electrooxidative rhodium(III/I)-catalyzed annulation reaction to assemble the valuable seven-membered benzoxepine skeleton (Scheme 1 b). Salient features of our approach comprise a) the first electrooxidative [5+2] cycloaddition, b) annulations by resource economical O–H/C–H functionalization, c) electrons as catalysts in cathodic proton reduction, d) isolation of key rhodium(I) intermediates and e) detailed mechanistic insights into electrooxidative rhodium catalysis.

We initiated our studies by exploring reaction conditions for the envisioned electrochemical [5+2] cycloadditions using 2-vinylphenol (**1a**) and diphenylacetylene (**2a**) in an undivided cell setup equipped with graphite felt (GF) and platinum plate (Pt) as anode and cathode material, respectively (Table 1). After considerable experimentations, the desired product **3aa** was isolated in 88% yield with [Cp*₂RhCl₂]₂ (2.5 mol %), NaOPiv (2.0 equiv) as additive in *t*AmOH/H₂O (3:1) at 100 °C for 18 h (Table 1, entry 1). The [5+2] annulation was not viable in the absence of NaOPiv,^[14] while KOAc in lieu of NaOPiv afforded a sharp decrease of

[*] Dr. Y. Wang, Dr. J. C. A. Oliveira, Z. Lin, Prof. Dr. L. Ackermann
 Institut für Organische und Biomolekulare Chemie
 and Wöhler Research Institute for Sustainable Chemistry
 Georg-August-Universität Göttingen
 Tammannstrasse 2, 37077 Göttingen (Germany)
 E-mail: Lutz.Ackermann@chemie.uni-goettingen.de

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202016895>.

© 2021 The Authors. Angewandte Chemie International Edition published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

Table 1: Optimization of the rhodium-catalyzed [5+2] cycloadditions.^[a]


Entry	Variation from standard conditions	3 aa [%] ^[b]
1	no change	88
2	no NaOPiv	trace
3	KOAc in place of NaOPiv	44
4	H ₂ O	47
5	tAmOH	15
6	GF as cathode electrode	trace
7	8 mA, 9 h	65
8	no electricity	7
9	under N ₂	79
10	Pd(OAc) ₂ in place of [Cp*RhCl ₂] ₂	0
11	[Cp*CoI ₂] ₂ in place of [Cp*RhCl ₂] ₂	0
12	[Cp*IrCl ₂] ₂ in place of [Cp*RhCl ₂] ₂	0

[a] Undivided cell, GF anode, Pt cathode, constant current = 4 mA, **1a** (1.0 mmol), **2a** (0.5 mmol), NaOPiv (1.0 mmol), [Cp*RhCl₂]₂ (2.5 mol %), solvent (4 mL), under air, 18 h. [b] Yields of isolated product **3aa**.

the yield (Table 1, entries 2–3). With H₂O or tAmOH alone as the solvent the electrocatalysis proved to be inefficient (Table 1, entries 4–5). Replacing the platinum cathode by a GF electrode inhibited the electrocatalysis (Table 1, entry 6). Increasing the current to 8 mA reduced the yield of product **3aa** to 65 %, and a control experiment confirmed the essential role of the electricity for the electrooxidative annulation (Table 1, entries 7–9). Reactions with other transition-metal catalysts, including Pd(OAc)₂ as well as Cp*-ligated iridium and cobalt complexes, proved to be ineffective for the annulation process (Table 1, entries 10–12).

With the optimized reaction conditions for the electrochemical [5+2] cycloadditions in hand (Table 1, entry 1), its performance was first explored with a set of substituted alkynes **2** (Scheme 2). The annulation was amenable to diverse diaryl alkynes **2** featuring both electron-withdrawing as well as electron-donating substituents on the aryl group (**3ab–3an**). Functional groups, including chloro, bromo, and even alkyl chloride as well as unprotected primary alcohol, were well tolerated. The annulation with unsymmetrical diaryl alkynes **2o** and **2p** results in 3.6:1 and 4.7:1 regioselectivities, respectively. The rhodaelectrocatalysis was also effective for dialkyl alkyne **2q**. Notably, unsymmetrical arylalkyl alkynes **2r–2w** were also efficiently converted to the corresponding benzoxepines with high levels of regiocontrol placing the alkyl group distal to the heteroatom (**3ar–3aw**). This regioselectivity was assessed by means of computational studies at the PW6B95-D4/def2-QZVP + SMD (methanol)//PBE0-D3(BJ)/def2-SVP level of theory (Figure 1).^[15] The regioisomer **3aw** was shown to be favored, placing the aryl group proximal to the heteroatom. Non-covalent secondary interactions play a dominant role in the stabilization of the preferred transition state.

The scope of the electrorhoda-catalyzed annulative reaction was examined with diversely substituted 2-vinylphenols

**Scheme 2.** Rhodaelectro-catalyzed [5+2] annulations with alkynes **2**.

1 (Scheme 3). Various phenols **1** were thereby selectively converted into the desired products **3** in high to excellent yields (**3ba–3ma**). Remarkably, the power of the metal-laelectrocatalysis was embodied in the chemo-selective C–H functionalization/annulation with sensitive iodovinylphenol **1l** to afford the desired product (**3la**).

The fluorescent BODIPY motifs are widely used as effective biological labels, luminescent tags and laser dyes.^[17] To our delight, the BODIPY-containing alkynes are suitable substrates for the metallaelectro-catalyzed annulations, and corresponding fluorescence-labeled benzoxepines (**3ax–3ay**)

- *DFT analysis of migratory insertion*





Figure 1. Computed Gibbs free energy profiles in kcal mol⁻¹ for the regioselectivity of the migratory insertion step of alkyne **3w** calculated at the PW6B95-D4/def2-QZVP+SMD(methanol)//PBE0-D3(BJ)/def2-SVP level of theory and visualization of noncovalent interactions calculated from the NCI plot for the respective transition states. In the latter, strong attractive interactions are given in blue and weak attractive interactions are given in green, respectively, while red corresponds to strong repulsive interactions.


could be easily obtained in one step (Scheme 4a). This transition metal-catalyzed C–H activation/annulation to assemble various BODIPY-labeled heterocycles would have great potential applications in drug deliveries, dyes, and optoelectronics.

The scalability of the electrochemical rhodium-catalyzed [5+2] annulation was further demonstrated for the assembly of benzoxepines **3**. The gram-scale reaction of substrates **1a** and **2a** hence yielded 1.33 g of the desired product **3aa** (Scheme 4b).

To gain further insight into the reaction mechanism, we conducted a series of experiments. Monitoring the catalytic process by NMR spectroscopy revealed that a low-valent rhodium complex **4** was likely the catalyst resting state in the electrorhoda-catalyzed annulative reaction (Scheme 5a). Notably, the corresponding rhodium(I) complexes **4aa** and **4aj** could be prepared and isolated by the reaction of Cp*Rh(OAc)₂ with the substrates **1a** and **2a** or **2j**, respectively. X-ray diffraction analysis featured rhodium(I) sandwich complexes in which the benzoxepines **3** are coordinated to the metal center as four-electron ligands (Scheme 5b).^[16] Complex **4aa** proved to be competent in the catalytic reaction (Scheme 5c). Electrolysis of **4aa** released product **3aa**, and the rhodium(I) was oxidized to the rhodium(III) species




Scheme 3. Rhodaelectro-catalyzed [5+2] cycloadditions with 2-vinylphenols. [a] 5.0 mol % [RhCp*Cl₂]₂.



Scheme 4. a) Syntheses of BODIPY-labeled benzoxepines. b) Gram-scale synthesis of **3 aa**.

Cp*Rh(OPiv)₂ (Scheme 5d). Increasing the electric current resulted in a higher initial reaction rate, indicating the reoxidation of rhodium(I) to rhodium(III) to be the rate-determining step (RDS) (Scheme 6a). H/D exchange experiments were conducted using D₂O as the deuterium source, and deuterium incorporation was not observed in the recovered 2-vinylphenols **1a** (Scheme 6b). Kinetic studies suggested a facial vinylic C–H metalation with a KIE value of $k_H/k_D \approx 1.0$ (Scheme 6c). Competition experiments with alkynes **2a** and **2j** revealed the greater reactivity of the electron-deficient alkyne **2j** (Scheme 6d).


CV studies of rhodium(I) complexes **4aa** and **4aj** exhibited irreversible oxidation waves at -0.058 V versus $\text{Fc}^{0/+}$ and -0.029 V versus $\text{Fc}^{0/+}$, respectively (Figure 2). It is reasonable that the latter has a higher oxidation potential



Scheme 5. Isolation and electrolysis studies of rhodium(I) intermediates.

since **4aj** possesses a electron-deficient metal center compared to **4aa**. A constant potential was conducted at 1.0 V affording the desired product **3aa** in 61% yield. Additional CV studies showed that the rhodium(I) intermediates **4aa** and **4aj** had lower oxidation potential than the substrates and products (Figure S10). The addition of PivOH had no significant influence on the oxidation potential of complex **4aa** (Figure S11).

On the basis of our findings, a plausible catalytic cycle is presented that commences with a facile O–H/C–H activation to afford a rhodacycle **A** (Figure 3). Then, alkyne coordination–insertion occurs to produce intermediate **C**, which rapidly undergoes reductive elimination to deliver the rhodium(I) sandwich complex **4**. Finally, the rhodium(I) species is re-oxidized to rhodium(III) at the anode, releasing the desired product **3** and generating molecular hydrogen as the by-product at the cathode. An alternative oxidation–



Scheme 6. Summary of key mechanistic findings.




Figure 2. Cyclic voltammetry studies. Conditions: **4aa** (2.5 mM) or **4aj** (2.5 mM), $n\text{Bu}_4\text{NPF}_6$ (0.1 M), MeCN, 100 mVs^{-1} .




Figure 3. Proposed catalytic cycle.

induced reduction elimination pathway may also be viable as depicted in Figure S9.^[15]

In conclusion, we have reported on the first electrocatalyzed [5+2] annulations to assemble valuable seven-membered benzoxepine skeletons by C-H/O-H activation. The versatility of the rhodaelectrosynthesis was demonstrated by its broad substrate scope and excellent functional group tolerance. The C-H activation employed electrons as catalysts for cathodic proton reduction, generating hydrogen as the sole byproduct. Detailed mechanistic studies provided strong support for a fast C-H rhodation and a rhodium(III/I) regime involving an efficient electrooxidation of the key benzoxepine-ligated rhodium(I) intermediate.

Acknowledgements

Generous support by the DFG (Gottfried-Wilhelm-Leibniz award to L.A.), the Alexander von Humboldt Foundation (fellowship to Y.W.), and the CSC (fellowship to Z.L.) is gratefully acknowledged. We thank Dr. Christopher Golz (Göttingen University) for assistance with the X-ray diffraction analysis. Open access funding enabled and organized by Projekt DEAL.

Conflict of interest

The authors declare no conflict of interest.

Keywords: [5+2] cycloaddition · benzoxepine · C-H activation · electrochemistry · electrooxidative annulation

- [1] For selected reviews of transition metal-catalyzed cycloadditions involving C-H bond activation, see: a) S. Rej, Y. Ano, N. Chatani, *Chem. Rev.* **2020**, *120*, 1788–1887; b) P. Gandeepan, T. Müller, D. Zell, G. Cera, S. Warratz, L. Ackermann, *Chem. Rev.* **2019**, *119*, 2192–2452; c) Y. Park, Y. Kim, S. Chang, *Chem. Rev.* **2017**, *117*, 9247–9301.
- [2] a) M. P. Huestis, L. Chan, D. R. Stuart, K. Fagnou, *Angew. Chem. Int. Ed.* **2011**, *50*, 1338–1341; *Angew. Chem.* **2011**, *123*, 1374–1377; b) S. Rakshit, F. W. Patureau, F. Glorius, *J. Am.*

- Chem. Soc.* **2010**, *132*, 9585–9587; c) Z. Shi, C. Zhang, S. Li, D. Pan, S. Ding, Y. Cui, N. Jiao, *Angew. Chem. Int. Ed.* **2009**, *48*, 4572–4576; *Angew. Chem.* **2009**, *121*, 4642–4646; d) D. R. Stuart, M. Bertrand-Laperle, K. M. N. Burgess, K. Fagnou, *J. Am. Chem. Soc.* **2008**, *130*, 16474–16475.
- [3] a) G. Mihara, K. Ghosh, Y. Nishii, M. Miura, *Org. Lett.* **2020**, *22*, 5706–5711; b) N. Casanova, A. Seoane, J. Mascareñas, M. Gulías, *Angew. Chem. Int. Ed.* **2015**, *54*, 2374–2377; *Angew. Chem.* **2015**, *127*, 2404–2407; c) J. D. Dooley, S. Reddy Chidipudi, H. W. Lam, *J. Am. Chem. Soc.* **2013**, *135*, 10829–10836; d) J. R. Huckins, E. A. Bercot, O. R. Thiel, T.-L. Hwang, M. M. Bio, *J. Am. Chem. Soc.* **2013**, *135*, 14492–14495; e) J. M. Neely, T. Rovis, *J. Am. Chem. Soc.* **2013**, *135*, 66–69; f) B. Ye, N. Cramer, *Science* **2012**, *338*, 504–506; g) X. Xu, Y. Liu, C.-M. Park, *Angew. Chem. Int. Ed.* **2012**, *51*, 9372–9376; *Angew. Chem.* **2012**, *124*, 9506–9510; h) X. Tan, B. Liu, X. Li, B. Li, S. Xu, H. Song, B. Wang, *J. Am. Chem. Soc.* **2012**, *134*, 16163–16166; i) T. K. Hyster, L. Knorr, T. R. Ward, T. Rovis, *Science* **2012**, *338*, 500–503; j) L. Ackermann, A. V. Lygin, N. Hofmann, *Angew. Chem. Int. Ed.* **2011**, *50*, 6379–6382; *Angew. Chem.* **2011**, *123*, 6503–6506; k) H. Shiota, Y. Ano, Y. Aihara, Y. Fukumoto, N. Chatani, *J. Am. Chem. Soc.* **2011**, *133*, 14952–14955; l) K. Sakabe, H. Tsurugi, K. Hirano, T. Satoh, M. Miura, *Chem. Eur. J.* **2010**, *16*, 445–449.
- [4] a) A. Seoane, N. Casanova, N. Quiñones, J. L. Mascareñas, M. Gulías, *J. Am. Chem. Soc.* **2014**, *136*, 834–837; b) S. Cui, Y. Zhang, Q. Wu, *Chem. Sci.* **2013**, *4*, 3421–3426; c) Z. Shi, C. Grohmann, F. Glorius, *Angew. Chem. Int. Ed.* **2013**, *52*, 5393–5397; *Angew. Chem.* **2013**, *125*, 5503–5507.
- [5] a) T. H. Meyer, I. Choi, C. Tian, L. Ackermann, *Chem* **2020**, *6*, 2484–2496; b) J. C. Siu, N. Fu, S. Lin, *Acc. Chem. Res.* **2020**, *53*, 547–560; c) P. Xiong, H.-C. Xu, *Acc. Chem. Res.* **2019**, *52*, 3339–3350; d) Y. Yuan, A. Lei, *Acc. Chem. Res.* **2019**, *52*, 3309–3324; e) S. Möhle, M. Zirbes, E. Rodrigo, T. Gieshoff, A. Wiebe, S. R. Waldvogel, *Angew. Chem. Int. Ed.* **2018**, *57*, 6018–6041; *Angew. Chem.* **2018**, *130*, 6124–6149; f) M. Yan, Y. Kawamata, P. S. Baran, *Chem. Rev.* **2017**, *117*, 13230–13319; g) R. Feng, J. A. Smith, K. D. Moeller, *Acc. Chem. Res.* **2017**, *50*, 2346–2352; h) R. Francke, R. D. Little, *Chem. Soc. Rev.* **2014**, *43*, 2492–2521; i) A. Jutand, *Chem. Rev.* **2008**, *108*, 2300–2347; j) J.-i. Yoshida, K. Kataoka, R. Horcajada, A. Nagaki, *Chem. Rev.* **2008**, *108*, 2265–2299.
- [6] For selected recent examples on electrooxidative syntheses, see: a) H. Yan, Z.-W. Hou, H.-C. Xu, *Angew. Chem. Int. Ed.* **2019**, *58*, 4592–4595; *Angew. Chem.* **2019**, *131*, 4640–4643; b) Y. Liang, F. Lin, Y. Adeli, R. Jin, N. Jiao, *Angew. Chem. Int. Ed.* **2019**, *58*, 4566–4570; *Angew. Chem.* **2019**, *131*, 4614–4618; c) M. Rafiee, F. Wang, D. P. Hruszkewycz, S. S. Stahl, *J. Am. Chem. Soc.* **2018**, *140*, 22; d) P. Xiong, H.-H. Xu, J. Song, H.-C. Xu, *J. Am. Chem. Soc.* **2018**, *140*, 2460–2464; e) L. Schulz, M. Enders, B. Elsler, D. Schollmeyer, K. M. Dybala, R. Franke, S. R. Waldvogel, *Angew. Chem. Int. Ed.* **2017**, *56*, 4877–4881; *Angew. Chem.* **2017**, *129*, 4955–4959; f) E. J. Horn, B. R. Rosen, Y. Chen, J. Tang, K. Chen, M. D. Eastgate, P. S. Baran, *Nature* **2016**, *533*, 77–81.
- [7] For reviews on metallalectro-catalyzed C-H functionalization, see: a) Y. Qiu, C. Zhu, M. Stangier, J. Struwe, L. Ackermann, *CCS Chem.* **2020**, *2*, 1529–1552; b) L. Ackermann, *Acc. Chem. Res.* **2020**, *53*, 84–104; c) R. C. Samanta, T. H. Meyer, I. Siewert, L. Ackermann, *Chem. Sci.* **2020**, *11*, 8657–8670; d) K.-J. Jiao, Y.-K. Xing, Q.-L. Yang, H. Qiu, T.-S. Mei, *Acc. Chem. Res.* **2020**, *53*, 300–310; e) N. Sauermann, T. H. Meyer, Y. Qiu, L. Ackermann, *ACS Catal.* **2018**, *8*, 7086–7103.
- [8] For selected recent examples of metallalectro-catalyzed C-H functionalization, see: a) U. Dhawa, C. Tian, T. Wdowik, J. C. A. Oliveira, J. Hao, L. Ackermann, *Angew. Chem. Int. Ed.* **2020**, *59*, 13451–13457; *Angew. Chem.* **2020**, *132*, 13553–13559; b) T. H. Meyer, J. C. A. Oliveira, D. Ghorai, L. Ackermann, *Angew.*

- Chem. Int. Ed.* **2020**, *59*, 10955–10960; *Angew. Chem.* **2020**, *132*, 11048–11053; c) R. C. Samanta, J. Struwe, L. Ackermann, *Angew. Chem. Int. Ed.* **2020**, *59*, 14154–14159; *Angew. Chem.* **2020**, *132*, 14258–14263; d) S.-K. Zhang, J. Struwe, L. Hu, L. Ackermann, *Angew. Chem. Int. Ed.* **2020**, *59*, 3178–3183; *Angew. Chem.* **2020**, *132*, 3204–3209; e) Z.-J. Wu, F. Su, W. Lin, J. Song, W.-B. Wen, H.-J. Zhang, H.-C. Xu, *Angew. Chem. Int. Ed.* **2019**, *58*, 16770–16774; *Angew. Chem.* **2019**, *131*, 16926–16930; f) Q.-L. Yang, X.-Y. Wang, J.-Y. Lu, L.-P. Zhang, P. Fang, T.-S. Mei, *J. Am. Chem. Soc.* **2018**, *140*, 11487–11494; g) S. Tang, L. Zeng, A. Lei, *J. Am. Chem. Soc.* **2018**, *140*, 13128–13135; h) N. Sauermann, T. H. Meyer, C. Tian, L. Ackermann, *J. Am. Chem. Soc.* **2017**, *139*, 18452–18455; i) Q.-L. Yang, Y.-Q. Li, C. Ma, P. Fang, X.-J. Zhang, T.-S. Mei, *J. Am. Chem. Soc.* **2017**, *139*, 3293–3298.
- [9] a) H. Shen, T. Liu, D. Cheng, X. Yi, Z. Wang, L. Liu, D. Song, F. Ling, W. Zhong, *J. Org. Chem.* **2020**, *85*, 13735–13746; b) F. Xu, Y. J. Li, C. Huang, H. C. Xu, *ACS Catal.* **2018**, *8*, 3820–3824.
- [10] a) C. Tian, U. Dhawa, A. Scheremetjew, L. Ackermann, *ACS Catal.* **2019**, *9*, 7690–7696; b) Y. Qiu, M. Stangier, T. H. Meyer, J. C. A. Oliveira, L. Ackermann, *Angew. Chem. Int. Ed.* **2018**, *57*, 14179–14183; *Angew. Chem.* **2018**, *130*, 14375–14379; c) L. Zeng, H. Li, S. Tang, X. Gao, Y. Deng, G. Zhang, C.-W. Pao, J.-L. Chen, J.-F. Lee, A. Lei, *ACS Catal.* **2018**, *8*, 5448–5453; d) Y. Qiu, W.-J. Kong, J. Struwe, N. Sauermann, T. Rogge, A. Scheremetjew, L. Ackermann, *Angew. Chem. Int. Ed.* **2018**, *57*, 5828–5832; *Angew. Chem.* **2018**, *130*, 5930–5934.
- [11] a) L. Yang, R. Steinbock, A. Scheremetjew, R. Kuniyil, L. H. Finger, A. M. Messinis, L. Ackermann, *Angew. Chem. Int. Ed.* **2020**, *59*, 11130–11135; *Angew. Chem.* **2020**, *132*, 11223–11229; b) Q.-L. Yang, Y.-K. Xing, X.-Y. Wang, H.-X. Ma, X.-J. Weng, X. Yang, H.-M. Guo, T.-S. Mei, *J. Am. Chem. Soc.* **2019**, *141*, 18970–18976; c) W.-J. Kong, L. H. Finger, A. M. Messinis, R. Kuniyil, J. C. A. Oliveira, L. Ackermann, *J. Am. Chem. Soc.* **2019**, *141*, 17198–17206; d) T. H. Meyer, J. C. A. Oliveira, S. Chandra Sau, N. W. J. Ang, L. Ackermann, *ACS Catal.* **2018**, *8*, 9140–9147; e) R. Mei, N. Sauermann, J. C. A. Oliveira, L. Ackermann, *J. Am. Chem. Soc.* **2018**, *140*, 7913–7921; f) R. Mei, J. Koeller, L. Ackermann, *Chem. Commun.* **2018**, *54*, 12879–12882; g) S. Tang, D. Wang, Y. Liu, L. Zeng, A. Lei, *Nat. Commun.* **2018**, *9*, 798–803; h) P. Huang, P. Wang, S. Wang, S. Tang, A. Lei, *Green Chem.* **2018**, *20*, 4870–4874; i) M.-J. Luo, M. Hu, R.-J. Song, D.-L. He, J.-H. Li, *Chem. Commun.* **2019**, *55*, 1124–1127.
- [12] a) M. D. Delost, D. T. Smith, B. J. Anderson, J. T. Njardarson, *J. Med. Chem.* **2018**, *61*, 10996–11020; b) K. Narita, K. Nakamura, Y. Abe, T. Katoh, *Eur. J. Org. Chem.* **2011**, 4985–4988 and references therein.
- [13] a) P. Gandeepan, L. H. Finger, T. H. Meyer, L. Ackermann, *Chem. Soc. Rev.* **2020**, *49*, 4254–4272; b) L. Ackermann, *Acc. Chem. Res.* **2014**, *47*, 281–295; c) L. Ackermann, *Synlett* **2007**, 0507–0526.
- [14] L. Ackermann, *Chem. Rev.* **2011**, *111*, 1315–1345.
- [15] For detailed information, see the Supporting Information.
- [16] Deposition numbers 2042970 (**3ak**), 2042973 (**3as**), 2012969 (**3ka**), 2042968 (**4aa**) and 2042967 (**4aj**) contain the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service.
- [17] a) T. Kowada, H. Maeda, K. Kikuchi, *Chem. Soc. Rev.* **2015**, *44*, 4953–4972; b) A. Kamkaew, S. H. Lim, H. B. Lee, L. V. Kiew, L. Y. Chung, K. Burgess, *Chem. Soc. Rev.* **2013**, *42*, 77–88; c) A. Loudet, K. Burgess, *Chem. Rev.* **2007**, *107*, 4891–4932.

Manuscript received: December 20, 2020

Accepted manuscript online: January 20, 2021

Version of record online: February 8, 2021