

Synthesis of 25-Hydroxycholesterol from 3 β -Hydroxyandrost-5-en-17-one. A Method for Stereospecific Construction of a Sterol Side-chain

By JERZY WICHA* and KAROL BAL

(*Institute of Organic Chemistry of the Polish Academy of Sciences, 00-961 Warszawa, Kasprzaka 44, Poland*)

Summary The side-chain of 25-hydroxycholesterol was constructed stereospecifically on the androstane skeleton *via* addition of an ethyl acetate unit to C-17 and sub-

sequent alkylation of the resulting ester C-20 with a C₅ unit; the remaining carbon atom (C-27) was added by a Grignard reaction.


RECENTLY there has been growing interest in developing efficient methods for the preparation of 25-hydroxycholesterol¹ required as a key intermediate in the synthesis of 1 α ,25-dihydroxy-vitamin D₃. We report the first stereospecific synthesis of this compound from an easily accessible androstane derivative.

Our approach to the stereospecific construction of a side chain on the androstane skeleton was based on the assumption that C-20 alkylation of the ester (2) would give predominantly the needed product (3a) with the (20*R*) configuration. Also, the (20*S*) epimer, if formed, could be converted into the (20*R*) epimer upon equilibration (*cf.* refs. 2 and 3). The androstene acetate (1) was transformed into the ester (2)[†] by treatment with ethyl bromoacetate under Reformatsky conditions followed by dehydration, selective hydrogenation of the 17–20 double bond,⁴ and exchange of the protecting groups.⁵ The ester (2) was alkylated with the dioxolan⁶ (5) by the procedure of Cregge *et al.*⁷ A solution of (2) in tetrahydrofuran under argon at –78 °C was treated with di-isopropyl-lithium amide (3 equiv.) and the bromo-compound (5) (2 equiv.) in hexamethylphosphorotriamide was added stepwise to the resulting solution of the enolate anion. The product (3a) was isolated by extraction with ether; m.p. 159–165 °C (from MeOH); δ (100 MHz, CDCl₃) 5.36 br (s, 1H, 6-H), 4.72 br (s, 1H, THP-H), 3.92 (s, 4H, dioxolan-H), 3.87 (m, 1H, 3 α -H), 3.65 (s, 3H, OMe), 3.50 (m, 2H, THP-H), 1.28 (s, 3H, 26-H), 1.00 (s, 3H, 19-H), and 0.70 (s, 3H, 18-H) (66% yield).

Compound (3a) was reduced with LiAlH₄ to the alcohol (3b) which was tosylated to give (3c). Hydrogenolysis of the tosyloxy-group in (3c) by LiAlH₄ gave the 21-unsubstituted compound (4a). Hydrolysis of (4a) furnished the alcohol (4b) which was acetylated (Ac₂O in pyridine) to (4c).

The homogeneity of the acetate (4c) was confirmed by t.l.c. and high-pressure liquid chromatography. Its physical properties (m.p. 138–139.5 °C, [α]_D –44°) were in good agreement with those of an authentic sample. The 21-Me signal in its ¹H n.m.r. spectrum was in a position which is

typical of the (20*R*) epimer (δ 0.95, d, *J* 7 Hz); there was no signal at δ 0.86 corresponding to the (20*S*) epimer.



Reaction of the acetoxy ketone (4b) with MeMgI gave the 25-hydroxycholesterol (4d), in 42% overall yield from (1) and 53% from (2).

We thank Dr. Uskokovic for an authentic sample of (4c).

(Received, 1st October 1975; Com. 1121.)

[†] Satisfactory elemental analyses were obtained for all new compounds.

¹ J. J. Partridge, S. Faber, and M. R. Uskoković, *Helv. Chim. Acta*, 1974, **57**, 764; T. A. Narwid, K. E. Cooney, and M. R. Uskoković, *ibid.*, p. 771; J. P. Moreau, D. J. Aberhart, and E. Caspi, *J. Org. Chem.*, 1974, **39**, 2018; A. Rotman and Y. Mazur, *J.C.S. Chem. Comm.*, 1974, 15.

² D. H. R. Barton, T. Shioiri, and D. A. Widdowson, *Chem. Comm.*, 1970, 939; R. Hayatsu, *Pharm. Bull. (Japan)*, 1957, **5**, 452.

³ A. Bowers, T. G. Halsall, and G. C. Sayer, *J. Chem. Soc.*, 1954, 3070, and references therein.

⁴ T. Reichstein, H. Müller, C. Meystre, and M. Sutter, *Helv. Chim. Acta*, 1939, **22**, 741; Pl. A. Plattner and W. Schreck, *ibid.*, 1939, **22**, 1178; D. Magrath, D. S. Morris, V. Petrow, and R. Royer, *J. Chem. Soc.*, 1950, 2393.

⁵ J. Wicha, and K. Bal, *Roczniki Chem.*, in the press.

⁶ C. A. Grob and R. Mösch, *Helv. Chim. Acta*, 1959, **42**, 728.

⁷ R. J. Cregge, J. L. Herrmann, C. S. Lee, J. E. Richmann, and R. H. Schlessinger, *Tetrahedron Letters*, 1973, 2425.