

0020-1669/82/1321-2051\$01.25/0 © 1982 American Chemical Society



instances attempted one-electron oxidation led to two-electron oxidation of half the starting material.


When  $\text{CpCo}(\text{PPh}_3)_2$  (**1**) was oxidized with  $\text{AgCN}$ , the only characterized product was  $\text{CpCo}(\text{CN})_2(\text{PPh}_3)$  (**4**). This diamagnetic  $\text{Co(III)}$  product was obtained in high yield when 2 equiv of  $\text{AgCN}$  was used. It is an air-stable and thermally stable yellow, crystalline compound. The infrared spectrum ( $\text{KBr}$ ) displays a medium-intensity absorption at  $2130\text{ cm}^{-1}$  ( $\nu_{\text{CN}}$ ). The  $^1\text{H}$  NMR spectrum gives a phenyl multiplet at  $\tau$  2.5 and a sharp singlet at  $\tau$  4.9 (relative integration 15:5). A number of similar complexes (**4**), where  $\text{X} = \text{I}$  and  $\text{L} = \text{CO}$ ,  $\text{PR}_3$ ,  $\text{P(OR)}_3$ , or nitrogen donor ligands, have previously been reported.<sup>14</sup> They were derived from the oxidation of  $\text{CpCo}(\text{CO})_2$  with  $\text{I}_2$  to give  $\text{CpCoI}_2(\text{CO})$  with subsequent substitution of the carbonyl group.

Treating  $[\text{CpCo}(\text{PPh}_3)_2][\text{BF}_4]$  (**2**) with 1 equiv of  $\text{AgBF}_4$  affords the diamagnetic "16-electron" complex  $[\text{CpCo}(\text{PPh}_3)_2][\text{BF}_4]_2$  (**5**). The same product is isolated upon oxidation of  $\text{CpCo}(\text{CO})(\text{PPh}_3)$  with  $\text{AgBF}_4$ . The pale yellow, air-stable solid exhibits an infrared spectrum ( $\text{KBr}$ ), which reveals the presence of  $\text{Cp}$ ,  $\text{PPh}_3$ , and  $\text{BF}_4$ . The  $^1\text{H}$  NMR spectrum shows two  $\text{PPh}_3$  groups per  $\text{Cp}$  group, the latter giving a sharp singlet at  $\tau$  4.25. No evidence for a metal hydride resonance could be found in a relatively concentrated  $\text{CD}_2\text{Cl}_2$  solution.

In the  $\text{L} = \text{PEt}_3$  system different  $\text{Co(III)}$  complexes are isolated. When  $(\text{Et}_3\text{P})_2\text{CoCl}_2$  is treated sequentially with  $\text{TiCp}$  and  $\text{AgBF}_4$ , oxidation takes place instead of metathesis, and the diamagnetic crystalline complex  $[\text{CpCoCl}(\text{PEt}_3)_2][\text{BF}_4]$  (**6**) is isolated in good yield, presumably through the intermediacy of **3**. Besides absorptions typical for coordinated cyclopentadienyl,  $\text{PEt}_3$ , and  $\text{BF}_4$  counterion, the infrared spectrum ( $\text{KBr}$ ) shows a weak absorbance at  $310\text{ cm}^{-1}$  ( $\nu_{\text{Co-Cl}}$ ). The  $^1\text{H}$  NMR spectrum shows a  $\text{Cp}$  singlet at  $\tau$  4.65 and  $\text{PEt}_3$  multiplets at  $\tau$  8.0 and 8.9.

On the other hand, when  $[\text{CpCo}(\text{PEt}_3)_2][\text{BF}_4]$  (**2**) is treated with 1 equiv of  $\text{AgBF}_4$  in THF, deposition of silver metal occurs, and a dark solution similar to that of **6** is obtained from which only an uncharacterized oil could be separated. However, if this dark solution is treated dropwise with 1 equiv of  $\text{PEt}_3$ , a dramatic color change to yellow occurs, and a diamagnetic solid is isolated in good yield. The infrared spectrum shows bands for  $\text{Cp}$ ,  $\text{PEt}_3$ , and  $\text{BF}_4$  and is otherwise unremarkable. However, the  $^1\text{H}$  NMR spectrum reveals a complex pattern (see Experimental Section) revealing the presence of a substituted cyclopentadienyl ring (multiplets of intensity 2 at  $\tau$  3.7 and 4.4) and a metal hydride (triplet of intensity 1 at  $\tau$  25.4). These data and the elemental analysis are consistent with the formulation  $[(\text{Et}_3\text{PC}_5\text{H}_4)\text{CoH}(\text{PEt}_3)_2][\text{BF}_4]_2$  (**7**), which is an isomer of the expected product **8**. Werner and Hofmann<sup>19</sup> recently reported the reaction of  $\text{CpCo}(\text{PMe}_3)_2$  (**1**) with alkyl halides ( $\text{RX}$ ) to give the similar complexes  $[(\text{RC}_5\text{H}_4)\text{CoH}(\text{PMe}_3)_2][\text{X}]$ . They proposed and gave evidence for the intermediacy of  $[\text{CpCoRL}_2]\text{X}$ , followed by intramolecular migration of  $\text{R}$  to the cyclopentadienyl ring, followed by loss of a ring proton and reattack on the cobalt center. A similar reaction path using the neutral phosphine ligand ( $\text{L}$ ) instead of an alkyl group is represented by the counterclockwise path of Scheme II. It is interesting to note that whereas there is precedence for the migration of alkyl or aryl groups to cyclopentadienyl rings,<sup>20</sup> we are unaware of any reports of phosphine migration. An alternative is the more direct clockwise pathway of Scheme II, incorporating intermolecular attack of ( $\text{L}$ ) on the cyclopentadienyl ring in a position exo

Scheme II



to the metal with transfer of a ring proton to the metal. The attack of phosphines on organic ligands of cationic complexes has been reported for other systems,<sup>21</sup> and we favor it here on the basis of steric considerations. It should be noted that while we have found no evidence for **8** ( $\text{L} = \text{PEt}_3$ ), the less sterically crowded complex **8** ( $\text{L} = \text{PMe}_3$ ) has been isolated by Werner and Hofmann.<sup>10</sup> Also, we have been unable to convert  $[\text{CpCoCl}(\text{PEt}_3)_2][\text{BF}_4]$  (**6**) to either **7** or **8** by reaction of  $\text{PEt}_3$  and  $\text{TiBF}_4$  either sequentially or concurrently; it appears that instead the cyclopentadienyl group is displaced.

In the systems where  $\text{L} = \text{P(OR)}_3$  ( $\text{R} = \text{Me}$ ,  $\text{Et}$ ,  $i\text{-Pr}$ ) the reaction of either **1** or **2** with  $\text{AgBF}_4$  in the presence of free ligand afforded the diamagnetic complexes  $[\text{CpCo}(\text{P(OR)}_3)_2][\text{BF}_4]_2$  (**8**) in good yields. The same complexes can be obtained by sequential treatment of  $\text{CpCoI}_2(\text{CO})$  with 3 equiv of  $\text{P(OR)}_3$  and 2 equiv of  $\text{TiBF}_4$ . No evidence for complexes **3**, **6**, or **7** was detected with these ligands. However, treatment of  $\text{CpCoI}_2(\text{CO})$  with 1 equiv of  $\text{P(OR)}_3$  does give the corresponding complex **4**.

**Oxidation of  $\text{CpCo}(\text{CO})_2$ .** Oxidation of  $\text{CpCo}(\text{CO})_2$  with 1 equiv of  $\text{AgBF}_4$  causes evolution of gas and deposition of silver metal, leading to the isolation of an apparently air-stable, off-white solid. The infrared spectrum suggests the presence of  $\text{Cp}$ ,  $\text{BF}_4$ , and THF. We find no IR carbonyl bands. The product is highly paramagnetic with  $\chi = 1.62 \times 10^{-5}$ . Treatment of this product with phosphines or phosphites has

(19) Werner, H.; Hofmann, W. *Angew. Chem., Int. Ed. Engl.* **1977**, *16*, 794; **1978**, *17*, 464.

(20) Bart, J. C.; Knox, S. A. R.; McKinney, R. J.; Stone, F. G. A. *J. Chem. Soc., Dalton Trans.* **1977**, 1.

(21) (a) Sweigart, D. A.; Kane-Maguire, L. A. P. *J. Chem. Soc., Chem. Commun.* **1976**, 13. (b) Hackett, P.; Jaouen, G. *Inorg. Chim. Acta* **1975**, *12*, L19. (c) Fairhurst, G.; White, C. *J. Organomet. Chem.* **1979**, *166*, C23-C25.

given no recognized products, and so the structure of the complex remains unclear.

## Conclusions

It appears that stable paramagnetic organocobalt(II) complexes can be isolated, provided the ligands are sufficiently bulky to prevent dimerization of the Co(II) complex (eq 2) or disproportionation to Co(I) and Co(III) complexes. Accordingly, treatment of any of the isolated complexes **2** with  $P(OMe)_3$  results in immediate disproportionation to diamagnetic products. The disproportionation reaction will be discussed in a later paper. All of the paramagnetic Co(II) complexes can be further oxidized to Co(III) containing two or three ligands, depending on the steric requirements of the ligand. However, sterics alone do not control the geometry of the Co(III) complex as revealed by the formation of **7** when  $L = PEt_3$ . Also it is surprising that treating **5** ( $L = PPh_3$ ) with  $PEt_3$  does not convert it to **7** but merely substitutes one of the  $PPh_3$  groups. If the ligands are small enough, i.e.,  $PMe_3$  or  $P(OR)_3$ , it appears that **8** will be formed.

## Experimental Section

All reactions and manipulations were carried out under nitrogen unless otherwise noted. Tetrahydrofuran (THF) was freshly distilled from sodium/benzophenone under nitrogen. All other solvents were purged with nitrogen and stored over 4A molecular sieves.

Proton NMR spectra and Evans method magnetic susceptibility measurements<sup>15</sup> were obtained at 60 MHz unless otherwise noted. Infrared spectra were obtained from KBr pellets on a Perkin-Elmer 283B spectrophotometer. Melting points were taken under nitrogen.

**$(Ph_3P)_3CoCl$ .** An electrolytic H-cell (1 L) separated by a Nafion perfluorosulfonic acid membrane was connected to a Princeton Applied Research Model 173 Potentiostat/Galvanostat.  $LiCl$  (25 g) was dissolved in absolute EtOH (1 L), and half of the solution was placed in one half of the cell with a Pt-gauze anode. The remaining solution, in which  $PPh_3$  (17 g) was dissolved with warming, was placed in the other half of the cell over a mercury-pool cathode.  $CoCl_2 \cdot 6H_2O$  (5.0 g) was dissolved in a minimum amount of EtOH and added to the  $PPh_3$  solution (Hg cathode), and controlled-potential (1.25 V vs.  $Ag/AgCl$  reference) electrolysis was carried out until the theoretical amount of current had been passed (ca. 2024 faradays, ~24 h). A bright green solid slowly separated from the blue solution. The mercury was drained away and the green suspension filtered to give a green solid, which was washed with two 50-mL portions of  $Et_2O$  and then pentane and dried under vacuum (17.4 g, 94%).

**$(\eta-C_5H_5)Co(PPh_3)_2$ .** A green suspension of  $(Ph_3P)_3CoCl$  (4.90 g, 5.6 mmol) in THF (250 mL) was treated with  $TiCl_3H_5$  (1.51 g, 5.6 mmol), giving a deep red solution with a white precipitate over a 15-min period. Stirring was continued for 3 h and the mixture filtered. Heptane (100 mL) was added to the red filtrate and the solution volume reduced to about 80 mL by aspiration. Red microcrystals separated and were filtered off and washed with pentane (3.06 g, 85%): mp 159–160 °C;  $^1H$  NMR ( $C_6D_6$ )  $\tau$  2.2 (m, 12), 3.0 (m, 18), 5.5 (t, 5,  $J = 1.5$  Hz).

**$(\eta-C_5H_5)Co(PEt_3)_2$ .**  $(Et_3P)_2CoI_2$  (6.00 g, 10.9 mmol) in THF (250 mL) was treated with  $TiCl_3H_5$  (2.94 g, 10.9 mmol) and excess Zn powder for 4 h at room temperature. The mixture was filtered and stripped down to a brown oil, which was extracted with warm pentane. The extract was allowed to evaporate slowly, leaving a brown solid (2.2 g, 55%):  $^1H$  NMR ( $C_6D_6$ )  $\tau$  5.35 (s, 5), 8.85 (m, 30).

**$(\eta-C_5H_5)Co(P(OMe)_3)_2$ .**  $(\eta-C_5H_5)_2Co$  (3.00 g, 15.9 mmol) was treated with 2-butyne (0.41 g, 7.7 mmol) and  $P(OCH_3)_3$  (9.45 g, 76 mmol) in heptane (30 cm<sup>3</sup>) and heated under reflux for 3 h. The solvent was evaporated under vacuum, and the residue was distilled under vacuum at ~150 °C to give a red oil. Crystallization from pentane at –35 °C gave an orange, crystalline solid (4.36 g; 74% yield):  $^1H$  NMR (acetone- $d_6$ )  $\tau$  5.35 (s, 5), 6.5 (t, 18,  $J = 6$  Hz).

**$(\eta-C_5H_5)Co(P(OPh)_3)_2$ .**  $(\eta-C_5H_5)_2CoI_2(CO)$  (2.00 g, 4.6 mmol) in THF (100 mL) was treated with  $P(OPh)_3$  (2.42 mL, 9.2 mmol), which caused evolution of gas, and then with excess zinc powder (0.5 g) for 3 h. The mixture was filtered and stripped to an orange solid, which was extracted with warm pentane. The orange pentane solution was reduced in volume by aspiration, giving an orange microcrystalline

solid (1.55 g). The filtrate was allowed to evaporate slowly at –10 °C, giving well-formed orange crystals (0.75 g; mp 130 °C; total yield 67%):  $^1H$  NMR (acetone- $d_6$ )  $\tau$  2.7 (m, 30), 6.0 (s, 5).

**$[(\eta-C_5H_5)Co(PPh_3)_2]TlBF_4$ .** (a)  $(\eta-C_5H_5)Co(PPh_3)_2$  (2.00 g, 3.1 mmol) dissolved in THF (50 mL) was treated with  $AgBF_4$  (0.60 g, 3.1 mmol) and stirred at room temperature for 1 h. Filtration gave a black solid which was extracted with  $CH_2Cl_2$ . The combined filtrate and extract were treated dropwise with hexane. Filtration gave a golden microcrystalline solid (1.95 g, 86%), mp 250 °C dec.

(b)  $(Ph_3P)_3CoCl$  (2.0 g, 2.3 mmol) suspended in THF (50 mL) was treated with  $TiCl_3H_5$  (0.61 g, 2.3 mmol) for 0.5 h and filtered,  $AgBF_4$  (0.44 g, 2.3 mmol) was added, and stirring was continued for a further 0.5 h. Filtration gave a black solid (0.28 g). The brown filtrate was treated with hexane. Filtration and washing with pentane gave a golden solid (1.40 g, 83%): mp 250 °C; magnetic susceptibility ( $CH_2Cl_2$ )  $\chi = 1.38 \times 10^{-6}$ ,  $\mu_{eff} = 1.6 \mu_B$ . Anal. Calcd: C, 66.94; H, 4.76. Found: C, 66.89; H, 4.87.

(c)  $(Ph_3P)_2CoCl_2$  (10.00 g, 15.3 mmol) dissolved in THF (500 mL) was treated with  $TiCl_3H_5$  (4.11 g, 15.3 mmol) and  $TlBF_4$  (4.45 g, 15.3 mmol) and stirred overnight. Filtration gave a tan solid, which was extracted with  $CH_2Cl_2$ , and the extract was treated with hexane to give a golden microcrystalline solid (6.8 g, 60%), mp 250 °C. The filtrate from above was reduced in volume and treated with hexane to give a golden solid (3.7 g, 33%), which has a melting point of 230 °C, suggesting it is impure (combined yield of about 90%).

**$(\eta-C_5H_5)CoCl(PPh_3)$ .** (a)  $(Ph_3P)_2CoCl_2$  (6.80 g, 10.4 mmol) was suspended in THF (250 mL), and the mixture was treated with  $TiCl_3H_5$  (2.80 g, 10.4 mmol) and stirred overnight. The maroon suspension was filtered (2.9 g of green solid recovered), and the solvent level was reduced and treated with hexane, which caused a maroon solid to separate. Recrystallization of the maroon solid from methylene chloride/hexane gave a maroon, microcrystalline solid (3.06 g, 70%): mp 150 °C dec; magnetic susceptibility ( $CH_2Cl_2$ )  $\chi = 3.2 \times 10^{-6}$ ,  $\mu_{eff} = 1.8 \mu_B$ . Anal. Calcd: C, 65.50; H, 4.75; Cl, 8.40. Found: C, 65.11, 65.30; H, 4.73, 4.79; Cl, 9.01.

(b)  $(\eta-C_5H_5)Co(PPh_3)_2$  (2.00 g, 3.1 mmol) dissolved in THF (50 mL) was treated with  $AgCl$  (0.44 g, 3.1 mmol). After 1 h of stirring, the solid silver (0.36 g; theory 0.33 g) was separated by filtration and the level of solvent was reduced to about 10 mL. Heptane (25 mL) was added, and the solution allowed to stand for 0.5 h, resulting in the formation of deep maroon crystals (0.85 g, 65%), which were identified by IR, melting point, and magnetic susceptibility.

**$(\eta-C_5H_5)CoBr(PPh_3)$ .**  $(Ph_3P)_2CoBr_2$  (4.00 g, 5.4 mmol) was dissolved in THF (200 mL), and the solution was treated with  $TiCl_3H_5$  (1.45 g, 5.4 mmol) and stirred at room temperature for 4 h. A green-tinted, white solid was separated by filtration, leaving a deep red solution. Heptane (50 mL) was added, and the volume was reduced (to about 70 mL). Filtration gave a brown, microcrystalline solid (1.77 g, 71%; mp 125 °C dec), which was dried under high vacuum at 50 °C for 3 h: magnetic susceptibility ( $CH_2Cl_2$ )  $\chi = 2.63 \times 10^{-6}$ ,  $\mu_{eff} = 1.7 \mu_B$ . Anal. Calcd: C, 59.2; H, 5.1. Found: C, 58.4; H, 4.1.

**$(\eta-C_5H_5)CoI(PPh_3)$ .**  $(Ph_3P)_2CoI_2$  (4.00 g, 4.8 mmol) dissolved in THF (100 mL) was treated as above with  $TiCl_3H_5$  (1.29 g, 4.8 mmol). The product was isolated as above as red-black crystals (1.44 g, 59%) and dried under vacuum at 50 °C (mp 135 °C dec). Anal. Calcd: C, 53.8; H, 3.9. Found: C, 54.3; H, 3.9.

**$[(\eta-C_5H_5)Co(PEt_3)_2]Tl$ .**  $(Et_3P)_2CoI_2$  (3.0 g, 5.5 mmol) was suspended in THF (100 mL) and treated with  $TiCl_3H_5$  for 4 h. Separation of the yellow solid by filtration gave a deep red filtrate, which was treated with heptane (50 mL) and reduced in volume. Filtration gave red-brown crystals (1.66 g, 82%): mp 149 °C; magnetic susceptibility ( $CH_2Cl_2$ )  $\chi = 2.48 \times 10^{-6}$ ,  $\mu_{eff} = 1.7 \mu_B$ . Anal. Calcd: C, 41.90; H, 7.19. Found: C, 41.85; H, 7.24.

**$[(\eta-C_5H_5)Co(PEt_3)_2]TlBF_4$ .** (a)  $[(\eta-C_5H_5)Co(PEt_3)_2]I$  (1.65 g, 3.4 mmol) dissolved in THF (100 mL) was treated with  $TlBF_4$  (1.00 g, 3.4 mmol) for 1.5 h. A brown solid was separated by filtration and extracted with  $CH_2Cl_2$ , leaving a yellow solid (TII). Slow treatment of the extract with hexane resulted in a light brown precipitate (1.08 g, 72%): mp 175 °C, discoloring at 289 °C dec; magnetic susceptibility ( $CH_2Cl_2$ )  $\chi = 2.55 \times 10^{-6}$ ,  $\mu_{eff} = 1.7 \mu_B$ . Anal. Calcd: C, 45.63; H, 7.83. Found: C, 45.61; H, 7.80.

(b)  $(Et_3P)_2CoCl_2$  (10.00 g, 27.3 mmol) dissolved in THF (500 mL) was treated with  $TiCl_3H_5$  (7.35 g, 27.3 mmol) for 0.5 h,  $TlBF_4$  (7.95 g, 27.3 mmol) was added, and stirring was continued for a further 3 h. Hexane (100 mL) was added, and filtration gave a light brown

solid, which was extracted with  $\text{CH}_2\text{Cl}_2$  until no further coloration of the solvent occurred. Treatment of the extract with hexane (300 mL, dropwise with stirring) caused a solid to separate. Filtration gave a brown microcrystalline solid (9.53 g, 78%).

(c)  $(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{PEt}_3)_2$  (1.00 g, 2.8 mmol) dissolved in THF (80 mL) was treated with  $\text{AgBF}_4$  (0.54 g, 2.8 mmol) for 1 h. Filtration gave a black solid, which was extracted with  $\text{CH}_2\text{Cl}_2$ , giving a brown solution and leaving a black solid (Ag). The brown extract was treated with hexane to give a brown-black crystalline product (0.79 g). The above filtrate gave additional product (0.12 g) when treated with hexane; total yield 0.91 g, 74%.

$(\eta\text{-C}_5\text{H}_5)\text{CoCl}(\text{PEt}_3)_2$ .  $(\text{Et}_3\text{P})_2\text{CoCl}_2$  (3.0 g, 8.2 mmol) dissolved in THF (100 mL) was treated with  $\text{TiCl}_3\text{H}_5$  (2.21 g, 8.2 mmol) as above for 3 h. Separation of a white solid by filtration gives a burgundy-colored filtrate. The filtrate was reduced in volume, treated with heptane, and refiltered, giving a small amount of green solid. The volume was reduced further and filtered, giving maroon-black crystals (0.15 g), mp 65 °C. Anal. Calcd: C, 47.57; H, 7.20. Found: C, 47.40; H, 7.34.

$(\eta\text{-C}_5\text{H}_5)\text{CoBr}(\text{PEt}_3)_2$ .  $(\text{Et}_3\text{P})_2\text{CoBr}$  (4.0 g, 8.8 mmol) dissolved in THF (100 mL) was treated as above with  $\text{TiCl}_3\text{H}_5$  (2.36 g, 8.8 mmol). Filtration and isolation as above gave maroon crystals (0.21 g), which may be a mixture of  $[\text{CpCoL}_2]\text{Br}$  and  $\text{CpCoBrL}$ . Anal. Calcd for  $\text{CpCoL}_2\text{Br}$ : C, 46.36; H, 7.9. Calcd for  $\text{CpCoBrL}$ : C, 41.0; H, 6.2 (for  $\text{CpCoBr}(\text{PEt}_3)_2$ ). Found: C, 44.92; H, 7.33.

$[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{PEtPh}_2)_2][\text{BF}_4]$ .  $(\text{Ph}_2\text{EtP})_2\text{CoCl}_2$  (3.00 g, 5.4 mmol) dissolved in THF (200 mL) was treated with  $\text{TiCl}_3\text{H}_5$  (1.45 g, 5.4 mmol) and  $\text{TIBF}_4$  (1.56 g, 5.4 mmol) as above. Isolation as above gave a brown microcrystalline solid (2.23 g, 65%); mp 176–177 °C; magnetic susceptibility ( $\text{CH}_2\text{Cl}_2$ )  $\chi = 1.70 \times 10^{-6}$ ,  $\mu_{\text{eff}} = 1.65 \mu_{\text{B}}$ . Anal. Calcd: C, 61.97; H, 5.48. Found: C, 61.81; H, 5.49.

$(\text{Ph}_2\text{EtP})_2\text{CoX}_2$  with  $\text{TiCl}_3\text{H}_5$  (X = Cl, Br, I).  $(\text{Ph}_2\text{EtP})_2\text{CoX}_2$  was treated with an equimolar amount of  $\text{TiCl}_3\text{H}_5$  in THF at room temperature for 3 h. Filtration yielded a stoichiometric amount of  $\text{TiX}$ . Addition of heptane and reduction in volume followed by standing at –30 °C overnight led to the isolation of brown crystals, which melted in the absence of the mother liquor.

$(\eta\text{-C}_5\text{H}_5)\text{CoBr}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)$ .  $(\text{DPPE})\text{CoBr}_2$  (4.00 g, 6.5 mmol) suspended in THF (100 mL) was treated with  $\text{TiCl}_3\text{H}_5$  (1.74 g, 6.5 mmol) overnight. The mixture was filtered, treated with heptane (25 mL), and reduced in volume. Filtration gave a maroon solid (1.09 g, 28%); mp 145–147 °C; magnetic susceptibility ( $\text{CH}_2\text{Cl}_2$ )  $\chi = 1.18 \times 10^{-6}$ ,  $\mu_{\text{eff}} = 1.3 \mu_{\text{B}}$ . Anal. Calcd: C, 61.79; H, 4.82. Found: C, 61.05; H, 4.70.

$(\eta\text{-C}_5\text{H}_5)\text{CoI}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)$ .  $(\text{DPPE})\text{CoI}_2$  (4.00 g, 5.6 mmol) suspended in THF (100 mL) was treated with  $\text{TiCl}_3\text{H}_5$  (1.51 g, 5.6 mmol) for 3 h. The mixture was filtered, treated with heptane, and stripped of solvent. The resulting violet solid was washed with THF and hexane to give a violet solid (3.7 g, >95%), mp 183–185 °C. Anal. Calcd: C, 57.32; H, 4.47. Found: C, 57.28; H, 4.92.

$[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{P}(\text{O}-i\text{-Pr})_3)_2][\text{BF}_4]$ . Anhydrous  $\text{CoI}_2$  (1.00 g, 3.2 mmol) and  $\text{P}(\text{O}-i\text{-Pr})_3$  (1.33 g, 1.53 mL, 6.4 mmol) dissolved in THF (200 mL) were treated with  $\text{TiCl}_3\text{H}_5$  (0.86 g, 3.2 mmol) and stirred for 0.5 h. Addition of  $\text{TIBF}_4$  (0.93 g, 3.2 mmol) with continued stirring for 3 h resulted in a brown solution with a yellow precipitate. Filtration gave a brown solution, which was evaporated to a brown oil and extracted with  $\text{CH}_2\text{Cl}_2$ . The extract was treated with  $\text{Et}_2\text{O}$  and allowed to evaporate slowly under nitrogen. Filtration gave a yellow-green crystalline solid, which was washed with pentane (1.00 g, 50%); mp 95 °C; magnetic susceptibility ( $\text{CH}_2\text{Cl}_2$ )  $\chi = 1.45 \times 10^{-5}$ ,  $\mu_{\text{eff}} = 1.5 \mu_{\text{B}}$ . Anal. Calcd: C, 44.0; H, 7.50. Found: C, 43.8; H, 7.76.

$[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{P}(\text{OPh})_3)_2][\text{BF}_4]$ .  $(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{P}(\text{OPh})_3)_2$  (1.00 g, 1.3 mmol) dissolved in THF (25 mL) was treated with a solution of  $\text{AgBF}_4$  (0.30 g, 1.5 mmol) in THF (20 mL). After addition, the mixture was immediately filtered. The gray solid was extracted with  $\text{CH}_2\text{Cl}_2$  and treated with hexane, giving a green solid (0.20 g), mp 163 °C. The filtrate was treated with hexane to give green crystals (0.35 g), mp 155 °C. An analytical sample was recrystallized from  $\text{CH}_2\text{Cl}_2$ /hexane; magnetic susceptibility ( $\text{CH}_2\text{Cl}_2$ )  $1.6 \mu_{\text{B}}$ . Anal. Calcd: C, 59.2; H, 4.2. Found: C, 57.4; H, 4.4.

$[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{PPh}_3)_2][\text{BF}_4]$ . (a)  $(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{CO})(\text{PPh}_3)_2$  (2.0 g, 4.8 mmol) dissolved in THF (100 mL) was treated with  $\text{AgBF}_4$  (0.94 g, 4.8 mmol) for 1 h. Filtration gave a gray solid, which was extracted with  $\text{CH}_2\text{Cl}_2$ , giving a yellow solution and leaving silver metal (0.4 g). Treatment of the extract with hexane gave a light yellow solid (1.29 g, 65% based on  $\text{AgBF}_4$ ); mp 260 °C;  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ )  $\tau$

2.8 (m, 30), 4.25 (s, 5). Anal. Calcd: C, 59.85; H, 4.26; P, 7.54. Found: C, 59.87; H, 4.49; P, 7.16.

(b)  $[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{PPh}_3)_2][\text{BF}_4]$  (1.00 g, 1.4 mmol) suspended in THF (50 mL) was treated dropwise with a solution of  $\text{AgBF}_4$  (0.28 g, 1.4 mmol) in THF (10 mL). After 5 min the mixture was filtered. The gray solid was extracted with  $\text{CH}_2\text{Cl}_2$ , and the yellow extract was treated with hexane to give a yellow solid (0.12 g). The filtrate above was treated with hexane to give a pale yellow solid (0.70 g; mp 260 °C).

$(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{CN})_2\text{PPh}_3$ . (a)  $(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{PPh}_3)_2$  (2.00 g, 3.1 mmol) dissolved in THF (150 mL) was treated with  $\text{AgCN}$  (0.83 g, 6.2 mmol) for 2 h with stirring. The mixture was evaporated to dryness and extracted with  $\text{CH}_2\text{Cl}_2$ , giving a yellow solution, which was treated with hexane, resulting in a yellow crystalline solid (1.23 g, 91%; mp 250 °C dec). A sample was dried in vacuo at 50 °C for 24 h to remove  $\text{CH}_2\text{Cl}_2$  of crystallization. Anal. Calcd: C, 68.49; H, 4.57; N, 6.39. Found: C, 68.43; H, 4.86; N, 6.34.

$[(\eta\text{-C}_5\text{H}_5)\text{CoCl}(\text{PEt}_3)_2][\text{BF}_4]$ .  $(\text{Et}_3\text{P})_2\text{CoCl}_2$  (2.00 g, 5.5 mmol) dissolved in THF (150 mL) was treated with  $\text{TiCl}_3\text{H}_5$  (1.47 g, 5.5 mmol), resulting in a color change from blue to deep red and precipitation of a white solid. After 1 h the mixture was filtered (1.30 g of white solid; theory for  $\text{TiCl}$  1.31 g) and the red filtrate treated with  $\text{AgBF}_4$ . A rapid color change to red-brown occurred with precipitation of a light-colored solid, which darkened over a period of about 1 h. The mixture was filtered and the dark precipitate extracted repeatedly with warm THF until little color was washed away (ca. 1 L). The extracts were combined, and the volume was reduced to about 100 mL. Filtration gave a black, microcrystalline solid, which was washed with pentane (1.85 g, 70%);  $^1\text{H}$  NMR ( $\text{CD}_2\text{Cl}_2$ )  $\tau$  4.65 (s, 5), 8.0 (m, 14), 8.9 (m, 21). Anal. Calcd: C, 42.28; H, 7.25. Found: C, 42.42; H, 7.29.

$[(\eta\text{-C}_5\text{H}_5)\text{CoH}(\text{PEt}_3)_2][\text{BF}_4]$ .  $[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{PEt}_3)_2][\text{BF}_4]$  (0.21 g, 0.5 mmol) suspended in THF (80 mL) was treated with  $\text{AgBF}_4$  (0.09 g, 0.5 mmol), resulting in a deep brown solution with precipitated silver metal. The brown solution was separated by filtration and treated with  $\text{PEt}_3$  (0.1 mL), causing an immediate color change to yellow and precipitation of a yellow solid, which was filtered off and washed with pentane (0.27 g, 88%); mp 153–154 °C;  $^1\text{H}$  NMR (acetone- $d_6$ , 220 MHz)  $\tau$  3.7 (m, 2), 4.4 (m, 2), 7.2 (dq, 6,  $J = 13.5$ , 7.5 Hz), 8.0 (dq, 12,  $J = 7.5$ , 7.5 Hz), 8.6 (dt, 9,  $J = 20$ , 7.5 Hz), 8.75 (dt, 18,  $J = 15$ , 7.5 Hz), 25.4 (t, 1,  $J = 77$  Hz). Anal. Calcd: C, 42.33; H, 7.67. Found: C, 42.49; H, 7.77.

$[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{P}(\text{OCH}_3)_3)_2][\text{BF}_4]$ .  $(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{CO})\text{I}_2$  (0.50 g, 1.2 mmol) dissolved in THF (100 mL) was treated with  $\text{P}(\text{OCH}_3)_3$  (0.5 mL), causing a color change to red-brown with precipitation of a solid. Addition of  $\text{TIBF}_4$  (1.0 g, 3.4 mmol) with vigorous stirring for 0.5 h caused the mixture to turn bright yellow. The mixture was filtered, the solid extracted with  $\text{CH}_2\text{Cl}_2$  again, and the extract treated with hexane, resulting in a yellow solid (0.30 g); mp 245–246 °C dec;  $^1\text{H}$  NMR  $\tau$  4.1 (s, 5), 5.9 (virtual quartet, 27).

$[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{P}(\text{OEt})_3)_2][\text{BF}_4]$ .  $(\text{EtO})_3\text{P}_3\text{CoCl}$  (2.42 g, 4.1 mmol) dissolved in THF (100 mL) was treated with  $\text{TiCl}_3\text{H}_5$  (1.10 g, 4.1 mmol) for 1 h. The mixture was filtered and filtrate treated with  $\text{AgBF}_4$  (0.80 g, 4.1 mmol) for 1.5 h with stirring. Filtration gave a black solid, which was extracted with  $\text{CH}_2\text{Cl}_2$ , giving a green solution. Addition of  $\text{Et}_2\text{O}$  caused precipitation of a yellow solid (1.30 g, 40% based on  $\text{L}_3\text{CoCl}$ , 80% based on  $\text{AgBF}_4$ ); mp 220–221 °C dec;  $^1\text{H}$  NMR  $\tau$  3.9 (s, 5), 5.4 (m, 18), 8.5 (t, 27). Anal. Calcd: C, 36.7; H, 6.3. Found: C, 35.0; H, 6.3.

$[(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{P}(\text{O}-i\text{-Pr})_3)_2][\text{BF}_4]$ .  $(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{P}(\text{O}-i\text{-Pr})_3)_2$  (10.0 g, 18.5 mmol) dissolved in THF (100 mL) with excess  $\text{P}(\text{O}-i\text{-Pr})_3$  (10 mL) was treated with  $\text{AgBF}_4$  (7.1 g, 36.5 mmol) for 1 h with stirring. Filtration gave a yellow-gray solid, which was extracted with  $\text{CH}_2\text{Cl}_2$ . THF was added to the yellow extract and the volume reduced, giving a yellow, microcrystalline solid (10.3 g, 60%); mp 130 °C dec;  $^1\text{H}$  NMR  $\tau$  4.2 (s, 5), 4.85 (m, 9), 8.4 (d, 54). Anal. Calcd: C, 41.6; H, 7.4. Found: C, 41.5; H, 7.2.

$(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{CO})_2$  with  $\text{AgBF}_4$ .  $(\eta\text{-C}_5\text{H}_5)\text{Co}(\text{CO})_2$  (1.80 g, 10.0 mmol) dissolved in THF (100 mL) was treated with  $\text{AgBF}_4$  (1.95 g, 10.0 mmol) for 1 h. Filtration gave a gray solid, which was extracted with  $\text{CH}_2\text{Cl}_2$ . The extract was treated with hexane, giving a yellow solid (1.44 g); mp 164 °C dec; magnetic susceptibility ( $\text{CH}_2\text{Cl}_2$ )  $\chi = 1.65 \times 10^{-5}$ .

Registry No. 1a, 32993-07-0; 1b, 79639-49-9; 1c, 80719-04-6; 1e, 32677-72-8; 1f, 32611-34-0; 1, L = CO and  $\text{PPh}_3$ , 12203-85-9; 1, L

