Organic Letters

# Rhodium-Catalyzed Annulative Coupling of Isothiazoles with Alkynes through N–S Bond Cleavage

Gen Mihara,<sup>†</sup> Teppei Noguchi,<sup>†</sup> Yuji Nishii,<sup>\*,‡</sup><sup>®</sup> Yoshihiro Hayashi,<sup>§</sup><sup>®</sup> Susumu Kawauchi,<sup>§</sup> and Masahiro Miura<sup>\*,†</sup><sup>®</sup>

<sup>†</sup>Department of Applied Chemistry and <sup>‡</sup>Frontier Research Base for Global Young Researchers, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

<sup>§</sup>Department of Chemical Science and Engineering, School of Materials and Chemical Technology, Tokyo Institute of Technology, Ookayama, Meguro-ku, Tokyo 152-8552, Japan

**Supporting Information** 

**ABSTRACT:** A Rh(III)-catalyzed annulative coupling of 3,5diarylisothiazoles and alkynes is reported. The N–S bond in the isothiazole ring acts as an internal oxidant to regenerate the Rh(III) species in combination with an external Cu(II) oxidant, and the corresponding 1:2 coupling products are



obtained. The remarkable difference in the reaction outcome between isothiazoles and the relevant isoxazoles has been investigated by DFT calculations, revealing that the relative stability of the enolate intermediates dictates the product selectivity.

he polycyclic heteroaromatic scaffold is ubiquitous in I many natural compounds and has been a key motif in a wide range of manufactured functional molecules. The past decades witnessed a tremendous achievement in the field of transition-metal-catalyzed C-H activation<sup>1</sup> and subsequent oxidative annulation (annulative coupling) with alkynes or their equivalents,<sup>2</sup> allowing us to assemble various fused-ring systems with operational simplicity. In this reaction system, particularly using group 9 metal complexes (Co, Rh, Ir), catalytically active high-valent species are regenerated by external oxidants to ensure the catalyst turnover. Alternatively, some coordinating functionalities may act as two-electron oxidants through heterolytic bond cleavage within them.<sup>3,4</sup> Fagnou et al. reported a seminal Rh(III)-catalyzed isoquinolone synthesis utilizing the N-O bond of hydroxamic acids as internal oxidants.<sup>4b</sup> Afterward, a series of catalytic coupling protocols based on such an N-O oxidizing directing group strategy has been developed by many research groups involving Glorius, Hartwig, Yu, and Chiba.<sup>4,5</sup> Furthermore, N-N,<sup>6</sup> N- $S_{1}^{7}$  and  $C-N^{8}$  bonds have also been adopted to establish redox-neutral coupling protocols.

Most of these reactions employed the oxidizing groups (-OR, -NR, -SR) as leaving counterparts. Our group,<sup>9</sup> Zhu,<sup>10a,b</sup> and Dong<sup>10c</sup> recently developed Rh(III)-catalyzed 1-substututed isoquinoline synthesis via the cleavage of N–O bonds embedded in the aromatic ring systems of isoxazoles<sup>11</sup> and oxadiazoles. In each of the reactions, the whole of the directing groups is incorporated into the product skeleton. Upon our continuous investigation into such ring-to-ring transformations, we turned our interest into developing an annulative coupling of isothiazoles whose N–S bond may act as an internal oxidant. The use of an oxidizing N–S bond in catalysis has rarely been successful to this date after the first

report by Dong et al., who applied N-sulfiny limines to the synthesis of substituted pyridines.  $^{7\rm a}$ 

Herein, we report that 3,5-diarylisothiazoles selectively undergo cascade oxidative annulation with the incorporation of 2 equiv of alkynes in the presence of an additional Cu(II) oxidant to construct isoquinoline and isothiochromene rings conjugated by a methine moiety in a one-pot manner (Scheme 1a). This is in contrast to the reaction of 3,5-diarylisoxazoles, which gives the 1:3 coupling products exclusively (Scheme 1b).<sup>9c</sup> In order to rationalize the outcome, preliminary DFT calculations have been carried out for the relative stability and reactivity of the corresponding enolate intermediates.

At the outset, we carried out an optimization study for the model reaction of 3,5-diphenylisothiazole (1a) with diphenylacetylene (2a) (Table 1). The corresponding 1:2 coupling product 3aa was obtained in 93% yield using a [Cp\*Rh- $(MeCN)_3$  [SbF<sub>6</sub>]<sub>2</sub> catalyst (4.0 mol %) and a Cu(OAc)<sub>2</sub> oxidant (2.0 equiv) at 100 °C in dioxane solvent (entry 1). The structure of 3aa was unambiguously determined by X-ray crystallographic analysis, where the isoquinoline and isothiochromene fragments are lying in almost the same plane with a small torsion angle of  $11.6^{\circ}$ . Both the rhodium catalyst and the copper oxidant were essential for the reaction (entries 2 and 3). Analogous neutral Rh(III) complexes (entries 4 and 5), Cp\*CoI<sub>2</sub>(CO) (entry 6), and [(p-cymene)RuCl<sub>2</sub>]<sub>2</sub> (entry 7) were totally inactive for the present transformation. Among the tested solvents, DCE afforded the highest 93% isolated yield at a lower temperature of 60 °C (entries 8-11). Notably, the reaction proceeded efficiently even at 40 °C (entry 12). The reaction could be conducted in 1.0 mmol scale (Scheme 2).

Received: December 11, 2019

#### Scheme 1. Rh-Catalyzed Annulative Coupling of Isoxazole/ Isothiazole with Alkyne

(a) This work: annulative coupling of isothiazole



(b)  $\ensuremath{\text{Previous work:}}$  annulative coupling of isoxazole [ref. 9c]



## Table 1. Optimization Study<sup>a</sup>

	N-S	catalyst (4.0 mol%) Cu(OAc) <sub>2</sub> (2.0 equiv)		S Ph Ph
Ph 🦳	Ph Ph	solvent, temp., 18 h		
	1a 2a (2.0 equiv)			3aa
entry	catalyst	solvent	temp	yield <sup>b</sup>
1	[Cp*Rh(MeCN) <sub>3</sub> ][SbF <sub>6</sub>	] <sub>2</sub> dioxane	100 °C	93%
2 <sup><i>c</i></sup>	[Cp*Rh(MeCN) <sub>3</sub> ][SbF <sub>6</sub>	] <sub>2</sub> dioxane	100 °C	trace
3		dioxane	100 °C	n.d.
4	$[Cp*RhCl_2]_2$	dioxane	100 °C	n.d.
5	$[Cp^{E}RhCl_{2}]_{2}$	dioxane	100 °C	n.d.
6	Cp*CoI <sub>2</sub> (CO) with AgS (8.0 mol %)	bF <sub>6</sub> dioxane	100 °C	n.d.
7	[( <i>p</i> -cymene)RuCl <sub>2</sub> ] <sub>2</sub> wit AgSbF <sub>6</sub> (8.0 mol %)	h dioxane	100 °C	n.d.
8	[Cp*Rh(MeCN) <sub>3</sub> ][SbF <sub>6</sub>	] <sub>2</sub> dioxane	60 °C	93%
9	[Cp*Rh(MeCN) <sub>3</sub> ][SbF <sub>6</sub>	] <sub>2</sub> toluene	60 °C	76%
10	[Cp*Rh(MeCN) <sub>3</sub> ][SbF <sub>6</sub>	] <sub>2</sub> <sup>t</sup> AmOH	60 °C	65%
11	[Cp*Rh(MeCN) <sub>3</sub> ][SbF <sub>6</sub>	] <sub>2</sub> DCE	60 °C	98% (93%)
12	[Cp*Rh(MeCN) <sub>3</sub> ][SbF <sub>6</sub>	] <sub>2</sub> DCE	40 °C	77%

<sup>*a*</sup>Reaction conditions: **1a** (0.1 mmol), **2a** (0.2 mmol), catalyst (4.0 mol % metal),  $Cu(OAc)_2$  (0.2 mmol), solvent (1.5 mL). <sup>*b*</sup>Determined by NMR analysis. Isolated yield in parentheses. <sup>*c*</sup>Without  $Cu(OAc)_2$ . n.d. = not detected.

#### Scheme 2. Scale-Up Reaction



With the optimized reaction conditions, we evaluated the substrate scope with respect to alkynes 2 (Scheme 3). For aromatic alkynes 2b-2h, the desired 1:2 coupling products 3ab-3ah were obtained in high to excellent yields. The structures of 3ab, 3ac, and 3af were confirmed by the X-ray analysis. Bromo (3af) and ester (3ag) functionalities remained intact during the reaction. An aliphatic alkyne (4-octyne, 2i) was also applicable to the present reaction to give the corresponding product 3ai in 59% yield.

We next examined the annulative coupling of various 3,5diarylisothiazoles 1b-1l with the alkyne 2a (Scheme 4). Apparently, substituents on the 3-aryl ring (1b-1g) and on the 5-aryl ring (1h-1l) did not significantly affect the reaction outcome, and they respectively fall into the isoquinoline and the isothiochromene subunits of the products. Both the

#### Scheme 3. Substrate Scope for Alkynes



## Scheme 4. Substrate Scope for Isothiazoles



#### Scheme 5. Stoichiometric Reaction of 1a with Cp\*Rh(III)Complex<sup>*a*</sup>



<sup>*a*</sup>The structure of **4** is drawn with a 40% thermal ellipsoid; hydrogen atoms and the counter anion are omitted for clarity.



electron-donating (1c, 1d, 1i) and -withdrawing (1e, 1j) groups were tolerated. It is noteworthy that, for the reaction of 1f and 1k, the C-H activation took place only at the sterically



#### Scheme 7. A Proposed Reaction Mechanism

Scheme 8. Mechanistic Consideration for the Selectivity



more accessible positions to produce **3fa** and **3ka** exclusively. The positions of the methyl protrusions in the coupling products (**3ba**, **3fa**, **3ga**, **3ha**, **3ka**, **3la**) were all determined by the X-ray analysis.

The isothiazole ring inherently offers two coordination sites, i.e., nitrogen and sulfur atoms. In order to elucidate which atom participates in the initial C–H activation event, we carried out a stoichiometric reaction of **1a** with the Rh(III) complex (Scheme 5). According to the literature procedure,<sup>12</sup> **1a** was treated with  $[Cp*Rh(MeCN)_3][SbF_6]_2$  in the presence of NaOAc as an acetate source. The corresponding metallacycle complex 4 was isolated in 17% yield, and the X-ray measurement revealed its structure as a nitrogen-coordinated species. This result indicated that the annulative coupling first proceeded at the nitrogen side to furnish the isoquinoline ring; therefore, the second alkyne molecule constructed the sulfur heterocycle. The complex **4** served as an active catalyst to give **3aa** in 90% yield under the standard conditions (Scheme 6).

A proposed reaction mechanism for the 1:2 annulative coupling of 1a and 2a is illustrated in Scheme 7. The isothiazole ring coordinates to a cationic Cp\*Rh(III) species to form a rhodacycle complex A through the proximal C–H bond activation. After the alkyne insertion into the Rh–C bond, sequential C–N reductive elimination and N–S oxidative addition generate an S-thioenolate (thiolate) complex C.<sup>13</sup> Following the second C–H activation, alkyne insertion and reductive elimination assemble the isothiochromene ring to give the product 3aa. The liberated Rh(I) species is oxidized by Cu(OAc)<sub>2</sub> to regenerate the catalytically active Rh(III) species, closing the catalytic cycle.

The difference in the product selectivity between the isoxazoles and the relevant isothiazoles would be dictated by the relative stabilities of the enolate intermediates. We



Figure 1. Partial enthalpy profile of the annulative coupling of (a) isoxazole and (b) isothiazole. Values in parentheses are relative Gibbs energies.

DOI: 10.1021/acs.orglett.9b04437 Org. Lett. XXXX, XXX, XXX–XXX previously found that the reaction of the isoxazole with alkyne proceeded via tautomerization from the *O*-enolate intermediate, which is structurally relevant to the complex **C** in Scheme 7, to the corresponding *C*-enolate (Scheme 8, top). Therefore, the isoxazoles selectively produced the 1:3 coupling products. In sharp contrast, this is not the case for the isothiazoles, since the formation of *C*-thioenolate (thioketone) species is thermodynamically quite unfavorable (Scheme 8, bottom).<sup>14</sup> Accordingly, the 1:2 coupling products are considered to be dominant for the reaction of the isothiazoles.

In order to gain some insight into this speculation, we conducted a computational study (Figure 1).<sup>15</sup> The phenyl groups on the isoquinoline ring were omitted to simplify the calculation. In both cases of isoxazole (Figure 1a) and isothiazole (Figure 1b), the most stable configurations were isoquinoline-coordinated six-membered enol complexes E-1. Without the coordination of the nitrogen atom, two metastable states K-2 (for C-enolate and C-thio-enolate) and E-2 (for Oenolate and S-thio-enolate) were obtained. For the isoxazole, the C-enolate (K-2) exhibited a lower  $\Delta H$  (+9.6 kcal/mol) as compared to that of *O*-enolate (E-2,  $\Delta H = +15.8 \text{ kcal/mol}$ ). In sharp contrast, for the isothiazole, the C-enolate (K-2,  $\Delta H =$ +33.4 kcal/mol) was of much higher energy content than the S-thioenolate species (E-2,  $\Delta H = +23.1$  kcal/mol). Additionally, we calculated the activation enthalpies of the following C-H bond cleavage reactions via the concerted metalation deprotonation (CMD) mechanism from the intermediates K-2 and E-2. The activation barriers for the isoxazole were, respectively, 20.1 kcal/mol in the keto pathway (K-2  $\rightarrow$  K-TS  $\rightarrow$  K-3) and 20.8 kcal/mol in the enol pathway (E-2  $\rightarrow$  E-TS  $\rightarrow$  E-3). Since there was no significant difference between the activation energies, it is reasonable to conclude that the reaction outcome is mainly directed by the relative stability of the intermediates (E-2 vs K-2), being consistent with the explanation in Scheme 8. In a similar manner, the barriers for the isothiazole were estimated as 9.0 kcal/mol in the keto pathway and 9.4 kcal/mol in the enol pathway. The enthalpies obtained herein were high considering the fact that the reaction proceeded even at 40 °C (Table 1, entry 12). This is probably because the stabilization effect of isoquinoline coordination is overestimated due to the omission of the phenyl substituents.

In summary, we have developed a Rh(III)-catalyzed annulative coupling of 3,5-diarylisothiazoles and alkynes that occurs selectively in a 1:2 manner to furnish structurally intriguing isoquinoline—isothiochromene conjugates. DFT calculations revealed that the differences in the reaction outcomes between the isoxazoles and the isothiazoles are dominantly dictated by the relative stabilities of the corresponding keto and enol intermediates.

#### ASSOCIATED CONTENT

#### **Supporting Information**

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acs.orglett.9b04437.

Detailed experimental procedures, spectroscopic data, and copy of NMR spectra (PDF)

Atomic coordinates of all calculated molecules (XYZ)

# **Accession Codes**

CCDC 1969592 and 1969598–1969607 contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data\_request/ cif, or by emailing data\_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

# AUTHOR INFORMATION

## **Corresponding Authors**

\*E-mail: miura@chem.eng.osaka-u.ac.jp. \*E-mail: y nishii@chem.eng.osaka-u.ac.jp.

#### ORCID 💿

Yuji Nishii: 0000-0002-6824-0639 Yoshihiro Hayashi: 0000-0002-7650-4083 Masahiro Miura: 0000-0001-8288-6439

# Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

This work was supported by JSPS KAKENHI Grant No. JP 19K15586 (Grant-in-Aid for Young Scientists) to Y.N. and JP 17H06092 (Grant-in-Aid for Specially Promoted Research) to M.M. The numerical calculations were carried out on the TSUBAME3.0 supercomputer at the Tokyo Institute of Technology, Tokyo, Japan, and on the supercomputer at the Research Center for Computational Science, Okazaki, Japan. This computational work was supported by JSPS KAKENHI Grant No. JP 17K17720 (Grant-in-Aid for Young Scientists (B)) to Y.H. and a JST CREST Grant No. JPMJCR1522 to S.K.

# REFERENCES

(1) For a recent review on C-H functionalization, see: Sambiagio, C.; Schönbauer, D.; Blieck, R.; Dao-Huy, T.; Pototschnig, G.; Schaaf, P.; Wiesinger, T.; Zia, M. F.; Wencel-Delord, J.; Besset, T.; Maes, B. U. W.; Schnürch, M. *Chem. Soc. Rev.* **2018**, *47*, 6603.

(2) For selected reviews, see: (a) Satoh, T.; Miura, M. Chem. - Eur. J.
2010, 16, 11212. (b) Colby, D. A.; Bergman, R. G.; Ellman, J. A. Chem. Rev. 2010, 110, 624. (c) Boyarskiy, V. P.; Ryabukhin, D. S.; Bokach, N. A.; Vasilyev, A. V. Chem. Rev. 2016, 116, 5894. (d) Santhoshkumar, R.; Cheng, C.-H. Chem. - Eur. J. 2019, 25, 9366. (3) For a recent review, see: Huang, H.; Ji, X.; Wu, W.; Jiang, H. Chem. Soc. Rev. 2015, 44, 1155.

(4) For seminal work, see: (a) Wu, J.; Cui, X.; Chen, L.; Jiang, G.;
Wu, Y. J. Am. Chem. Soc. 2009, 131, 13888. (b) Guimond, N.;
Gouliaras, C.; Fagnou, K. J. Am. Chem. Soc. 2010, 132, 6908. (c) Too,
P. C.; Wang, Y. F.; Chiba, S. Org. Lett. 2010, 12, 5688. (d) Ng, K.-H.;
Chan, A. S. C.; Yu, W.-Y. J. Am. Chem. Soc. 2010, 132, 12862.
(e) Tan, Y.; Hartwig, J. F. J. Am. Chem. Soc. 2010, 132, 3676.
(f) Rakshit, S.; Grohmann, C.; Besset, T.; Glorius, F. J. Am. Chem. Soc.
2011, 133, 2350.

(5) For selected recent examples, see: (a) Bian, M.; Ma, K.; Mawjuda, H.; Yu, X.; Li, X.; Gao, H.; Zhou, Z.; Yi, W. Org. Biomol. Chem. **2019**, *17*, 6114. (b) Ramesh, B.; Tamizmani, M.; Jeganmohan, M. J. Org. Chem. **2019**, *84*, 4058. (c) Li, X.-C.; Du, C.; Zhang, H.; Niu, J.-L.; Song, M.-P. Org. Lett. **2019**, *21*, 2863. (d) Zhu, H.; Zhuang, R.; Zheng, W.; Fu, L.; Zhao, Y.; Tu, L.; Chai, Y.; Zeng, L.; Zhang, C.; Zhang, J. Tetrahedron **2019**, *75*, 3108. (e) Zhou, C.; Jiang, J.; Wang, J. Org. Lett. **2019**, *21*, 4971. (f) Sarathkumar, S.; Kavala, V.; Yao, C.-F. Asian J. Org. Chem. **2019**, *8*, 1830. (g) Guntreddi, T.; Shankar, M.; Kommu, N.; Sahoo, A. K. J. Org. Chem. **2019**, *84*, 13033. (h) Saiegh, T. J.; Chédotal, H.; Meyer, C.; Cossy, J. Org. Lett. **2019**, *21*, 8364.

(6) (a) Wang, Y.-F.; Toh, K. K.; Lee, J.-Y.; Chiba, S. Angew. Chem., Int. Ed. **2011**, 50, 5927. (b) Chuang, S.-C.; Gandeepan, P.; Cheng, C.-H. Org. Lett. **2013**, 15, 5750. (c) Wang, C.; Sun, H.; Fang, Y.; Huang, Y. Angew. Chem., Int. Ed. **2013**, 52, 5795. (d) Huang, X.-C.; Yang, X.-

## **Organic Letters**

H.; Song, R.-J.; Li, J.-H. J. Org. Chem. 2014, 79, 1025. (e) Deshmukh, D. S.; Gangwar, N.; Bhanage, B. M. Eur. J. Org. Chem. 2019, 2019, 2919.

(7) (a) Zhang, Q.-R.; Huang, j.-R.; Zhang, W.; Dong, L. Org. Lett.
2014, 16, 1684. (b) Kong, L.; Xie, F.; Yu, S.; Qi, Z.; Li, X. Chin. J. Catal. 2015, 36, 925. (c) Wang, F.; Wang, H.; Wang, Q.; Yu, S.; Li, X. Org. Lett. 2016, 18, 1306. (d) Wang, F.; Wang, Q.; Bao, M.; Li, X. Chin. J. Catal. 2016, 37, 1423. (e) Wang, Q.; Li, Y.; Qi, Z.; Xie, F.; Lan, Y.; Li, X. ACS Catal. 2016, 6, 1971. (f) Shankar, M.; Ghosh, K.; Mukherjee, K.; Rit, R. K.; Sahoo, A. K. Org. Lett. 2018, 20, 5144.

(8) Yu, S.; Liu, S.; Lan, Y.; Wan, B.; Li, X. J. Am. Chem. Soc. 2015, 137, 1623.

(9) (a) Nishii, Y.; Bachon, A.-K.; Moon, S.; Bolm, C.; Miura, M. Chem. Lett. 2017, 46, 1347. (b) Noguchi, T.; Nishii, Y.; Miura, M. Chem. Lett. 2017, 46, 1512. (c) Noguchi, T.; Nishii, Y.; Miura, M. Synthesis 2019, 51, 258.

(10) (a) Yang, F.; Yu, J.; Liu, Y.; Zhu, J. Org. Lett. 2017, 19, 2885.
(b) Yang, F.; Yu, J.; Liu, Y.; Zhu, J. J. Org. Chem. 2017, 82, 9978.
(c) Zhang, J.; Sun, J.-S.; Xia, Y.-Q.; Dong, L. Adv. Synth. Catal. 2019, 361, 2037.

(11) Anthranil (2,1-benzisoxazole) has been utilized as an aminating reagent under Rh catalysis through N-O bond cleavage: (a) Yu, S.; Tang, G.; Li, Y.; Zhou, X.; Lan, Y.; Li, Z. Angew. Chem., Int. Ed. 2016, 55, 8696. (b) Yu, S.; Li, Y.; Zhou, X.; Wang, H.; Kong, L.; Li, X. Org. Lett. 2016, 18, 2812. (c) Shi, L.; Wang, B. Org. Lett. 2016, 18, 2820. (d) Zou, M.; Liu, J.; Tang, C.; Jiao, N. Org. Lett. 2016, 18, 3030. (e) Tang, C.; Zou, M.; Liu, J.; Wen, X.; Sun, X.; Zhang, Y.; Jiao, N. Chem. - Eur. J. 2016, 22, 11165. (f) Debbarma, S.; Maji, M. S. Eur. J. Org. Chem. 2017, 2017, 3699. (g) Jeon, M.; Park, J.; Dey, P.; Oh, Y.; Oh, H.; Han, S.; Um, S. H.; Kim, S. H.; Mishra, N. K.; Kim, I. S. Adv. Synth. Catal. 2017, 359, 3471. (h) Wang, M.; Kong, L.; Wang, F.; Li, X. Adv. Synth. Catal. 2017, 359, 4411. (i) Wu, X.; Xiao, Y.; Sun, S.; Yu, J.-T.; Cheng, J. Org. Lett. 2019, 21, 6653. (j) Biswas, A.; Sarkar, S.; Samanta, R. Chem. - Eur. J. 2019, 25, 3000.

(12) Tindall, D. J.; Werlé, C.; Goddard, R.; Philipps, P.; Farès, C.; Fürstner, A. J. Am. Chem. Soc. 2018, 140, 1884.

(13) Guimond, N.; Gorelsky, S. I.; Fagnou, K. J. Am. Chem. Soc. 2011, 133, 6449.

(14) Kuhn, N.; Verani, G. Chalcogenone C=E compounds. In Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium; Devillanova, F. A., du Mont, W.-W., Eds.; RSC Publishing: Cambridge, U.K., 2013; Vol. 1.

(15) For details, see the Supporting Information.