

## A Journal of the Gesellschaft Deutscher Chemiker A Deutscher Chemiker GDCh International Edition www.angewandte.org

## **Accepted Article**

- Title: Photocatalytic α-Tertiary Amine Synthesis via C–H Alkylation of Unmasked Primary Amines
- Authors: Alexander James Cresswell, Alison Ryder, William Cunningham, George Ballantyne, Tom Mules, Anna Kinsella, Jacob Turner-Dore, Catherine Alder, Lee Edwards, Blandine McKay, and Matthew Grayson

This manuscript has been accepted after peer review and appears as an Accepted Article online prior to editing, proofing, and formal publication of the final Version of Record (VoR). This work is currently citable by using the Digital Object Identifier (DOI) given below. The VoR will be published online in Early View as soon as possible and may be different to this Accepted Article as a result of editing. Readers should obtain the VoR from the journal website shown below when it is published to ensure accuracy of information. The authors are responsible for the content of this Accepted Article.

To be cited as: Angew. Chem. Int. Ed. 10.1002/anie.202005294

Link to VoR: https://doi.org/10.1002/anie.202005294

## WILEY-VCH

# Photocatalytic α-Tertiary Amine Synthesis *via* C–H Alkylation of Unmasked Primary Amines

Alison S. H. Ryder,<sup>[a]</sup> William B. Cunningham,<sup>[b]</sup> George Ballantyne,<sup>[b]</sup> Tom Mules,<sup>[b]</sup> Anna G. Kinsella,<sup>[b]</sup> Jacob Turner-Dore,<sup>[b]</sup> Catherine M. Alder,<sup>[c]</sup> Lee J. Edwards,<sup>[c]</sup> Blandine S. J. McKay,<sup>[c]</sup> Matthew N. Grayson,<sup>[b]</sup> and Alexander J. Cresswell\*<sup>[b]</sup>

Dedicated to the memory of Professor Jonathan M. J. Williams (deceased on 03/08/2019)

**ABSTRACT:** A practical, catalytic entry to  $\alpha, \alpha, \alpha$ -trisubstituted ( $\alpha$ -tertiary) primary amines by C–H functionalisation has long been recognised as a critical gap in the synthetic toolbox. We report a simple and scalable solution to this problem that does not require any *in situ* protection of the amino group and proceeds with 100% atomeconomy. Our strategy, which uses an organic photocatalyst in combination with azide ion as a hydrogen atom transfer (HAT) catalyst, provides a direct synthesis of  $\alpha$ -tertiary amines, or their corresponding  $\gamma$ -lactams. We anticipate that this methodology will inspire new retrosynthetic disconnections for substituted amine derivatives in organic synthesis, and particularly for challenging  $\alpha$ -tertiary primary amines.

#### Introduction

Aliphatic amines and their simple derivatives are pervasive in bioactive molecules, and their centrality in medicinal chemistry is evidenced by their occurrence in over 40% of drug candidates.<sup>[1]</sup> Saturated azacyclic motifs such as piperidines and pyrrolidines<sup>[2]</sup> - as well as more conformationally-constrained analogues such as azaspirocycles<sup>[3]</sup> - are now a mainstay in drug discovery programmes, where their high fraction of saturated carbon (Fsp<sup>3</sup>) can markedly reduce compound attrition rates. Unsurprisingly, the growing demand for functionalised aliphatic amines and saturated azacycles in drug design has continued to spur the development of practical, catalytic methods for their synthesis.<sup>[4]</sup> The alkylation of  $\alpha$ -C-H bonds in aliphatic amines provides a powerful alternative to established C-N bond-forming strategies,[5] but novel methods for the synthesis of a-tertiary amines via C-H functionalisation are still urgently needed.<sup>[1]</sup> Given that primary alkyl amines offer the greatest potential for diversification, and are prevalent in pharmaceutical compound libraries (e.g., >2,700 nonbenzylic primary amines in GSK's internal chemical inventory), methods for the α-C-H alkylation of this particular amine class

[a] A. S. H. Ryder

- Centre for Sustainable Chemical Technologies, 1 South, University of Bath, Claverton Down, Bath, BA2 7AY (UK)
- [b] Dr W. B. Cunningham, G. Ballantyne, T. Mules, A. G. Kinsella, J. Turner-Dore, Dr A. J. Cresswell, Dr M. N. Grayson Department of Chemistry, 1 South, University of Bath, Claverton Down, Bath, BA2 7AY (UK)
- a.j.cresswell@bath.ac.uk [c] Dr C. M. Alder, L. J. Edwards, Dr B. McKay
- Medicines Design, GSK Medicines Research Centre, Gunnels Wood Rd, Stevenage, SG1 2NY (UK)

Supporting information for this article is given via a link at the end of the document

could be transformative. Despite isolated reports of non-catalytic  $\alpha$ -C–H alkylations of primary aliphatic amines,<sup>[6]</sup> including an amine dehydrogenation sequence with stoichiometric quinones,<sup>[7]</sup> catalytic protocols to *directly* access *C*-alkylated primary amines have proven elusive, especially for non-benzylic amines.<sup>[8]</sup> Indirect methods reliant on *N*-protection, catalytic C–H alkylation<sup>[9a,b]</sup> or arylation,<sup>[9b]</sup> and subsequent *N*-deprotection have been realised, but the triflamide or benzamide groups that are required suffer from harsh deprotection protocols. Notably, a recent study from Rovis, Schoenebeck and co-workers showed that *in situ N*-protection of aliphatic primary amines with CO<sub>2</sub> enables a catalytic  $\alpha$ -C–H alkylation process, leading to  $\gamma$ -lactam products (Figure 1A).<sup>[10]</sup>

A. State-of-the-art using *in situ* amine masking (Rovis *et al.*, 2018)



Figure 1. [A] Prior art for catalytic  $\alpha$ -C–H alkylation of primary amines using *in situ N*-protection; [B] This work. EWG = electron-withdrawing group.

The lack of a practical and scalable catalytic entry to  $\alpha$ -tertiary primary amines by C–H functionalisation was recently stressed as a key unsolved problem for synthetic chemistry.<sup>[1a,7]</sup> To address this challenge, our lab has initiated a research programme on the use of unprotected aliphatic primary amines as formal *C*-nucleophiles in catalytic C–C bond-formation, seeking to avoid *N*-protection strategies altogether. With carbon electrophiles as reactants, a critical issue is to outcompete the innate background reactivity of the free amines, leading to *N*-alkylation, and to steer the reactions towards an unconventional ('umpolung') *C*-alkylation. In this work, we report a practical and scalable solution to this problem, based on photoredox catalysis,<sup>[11]</sup> for the catalytic generation of unprotected  $\alpha$ -amino radicals<sup>[12]</sup> **2** from primary amines **1**, and their interception with electrophilic Michael

acceptors **3** to give  $\alpha$ -tertiary amines **4** (Figure 1B). We also demonstrate that these products can be readily elaborated – in a telescoped process – to *N*-functionalised  $\alpha$ -tertiary amine derivatives<sup>[13]</sup> or pharmacologically-valuable  $\gamma$ -lactams.<sup>[3]</sup>

#### **Results and Discussion**

We began our investigations by using cyclohexylamine 5  $(E_{D/2} = +1.53 \text{ V vs SCE}, \text{ in MeCN})$  and butyl acrylate **6** as model substrates (Figure 2A). By irradiating 5 and 6 in MeCN with 34 W Kessil blue LEDs, in the presence of various photoredox catalysts (PCs) and hydrogen atom transfer (HAT) catalysts (Figure 2B-C),<sup>[14]</sup> we found that  $\alpha$ -C–H alkylation could be cleanly effected in 85% NMR yield using 4CzIPN<sup>[15]</sup> as the photocatalyst (PC) and tetrabutylammonium azide (Bu<sub>4</sub>N<sup>+</sup>N<sub>3</sub><sup>-</sup>) as the HAT catalyst ( $E_{p/2}$  of  $N_3^-$  = +0.87 V vs SCE in MeCN<sup>[16]</sup>). Surprisingly, azide ion **11**<sup>[10]</sup> outperformed both quinuclidine (42%) and tri(isopropyl)silanethiolate 10 (70%), despite the well-established pedigree of these species as HAT catalysts. Control experiments verified that 4CzIPN, visible light, and azide catalyst are all necessary components for reactivity. To determine the optimal wavelength of light to use in the reaction, we tested a series of pseudo-monochromatic LED light sources (405-475 nm), and used flow NMR spectroscopy<sup>[17]</sup> to monitor the reaction (Figure 2D). A 425 nm light source proved to be optimal, consistent with the absorption maximum of 4CzIPN at  $\lambda_{max}$  = 435 nm. These NMR experiments also revealed that lactamisation proceeds to a negligible extent under the reaction conditions (at 25 °C), such that α-tertiary amines are the primary products.

With optimised conditions in hand, we next investigated the scope of the reaction with respect to the amine partner 1 (Figure 3A). Methyl acrylate 14 [or in some cases 2-methoxyethyl acrylate 15<sup>[18]</sup>] was used as the alkylating agent, and a 1:1 stoichiometry of amine:acrylate was employed in all cases (except 1f and 1p). As the initially-formed y-amino esters 16 are prone to lactamisation on heating, we chose to purposefully convert them to their y-lactams 17 in a subsequent lactamisation step, though other manipulations of the amino group are possible (Figure 4A). Under these generalised conditions, an array of primary amines 1 were assessed, providing C-H alkylated products 17 in yields ranging from 32-84%. Cyclic primary amines with 5-7-membered ring sizes delivered the corresponding azaspirocycles (17a-c) in 60-68% yield. Unsurprisingly, cyclobutyl amine 1d proved more challenging – a consequence of the stronger  $\alpha$ -C–H bond (see Supporting Information) - although the alkylated product 17d could still be isolated in 49% yield. A range of acyclic amines (17e-j) were also surveyed, and we found that tertiary (1f,q) or quaternary carbons (**1h**)  $\beta$ - to the amino group are tolerated. For  $\alpha$ -monosubstituted amine **1f**, a mixture of  $\alpha$ -mono and  $\alpha$ -dialkylated products was evident on using a 1:1 amine:acrylate stoichiometry, due to overalkylation of the intermediate y-amino ester 16f (see Supporting Information). However, after some reoptimisation (i.e., photocatalyst, amine equiv), α-monosubstituted y-lactam 17f could be isolated in 49% yield. One notable limitation is the lack of reactivity of benzylamine 1i, which may arise from the increased stability of the  $\alpha$ -amino radical, rendering the addition step to the acrylate reversible. That said, the presence

#### WILEY-VCH



Figure 2. [A] Reaction optimisation. [a] Measured by GC (gas chromatography) against dodecane as an internal standard; note that lactamisation occurs under the analysis conditions. [b] Measured by <sup>1</sup>H NMR against mesitylene as an internal standard. [c] <1% of the aza-Michael adduct was formed. [d] 4% of the aza-Michael adduct was formed. [B] Selected photocatalysts. [C] HAT catalysts. [D] Concentration-time plots, at different irradiation wavelengths, for formation of the α-tertiary amine product 8 (using methyl acrylate).

of benzylic hydrogens elsewhere in the molecule poses no issue, as evidenced by the successful  $\alpha$ -alkylation of amine 1j. Diastereoselective  $\alpha$ -alkylation of norbornyl amine 1k also proved possible, with the radical intercepting the acrylate species on the expected exo face. We also tested functionalised amines containing acetal (11,m), hydroxyl (1n,v), ether (1o,p,w), carbamate (1q-s), thioether (1t), sulfone (1u), and ester moieties (1x). In several cases, competitive HAT at other 'hydridic' C-H bonds is thought to be operative, including tertiary C-H (17f-h) and C-H bonds  $\alpha$ - to hydroxyl (17n,v), ether (17m,o,p,w), and carbamate moieties (17q-s). Though all of these C-H bonds are

#### WILEY-VCH

## **RESEARCH ARTICLE**



Figure 3. [A] Scope for  $\alpha$ -C-H alkylation of primary amines. [a] Used methyl acrylate 14 as the acceptor. [b] Used 2-methoxyethyl acrylate 15 as the acceptor, and performed the lactamisation step with Et<sub>3</sub>N (4.0 equiv) in MeCN at 100 °C for 24 h. [c] ~20% unreacted amine 1d. [d] Used 2.0 equiv of amine 1f, 1.0 equiv of acrylate 15, and Ir[(dF(CF<sub>3</sub>)ppy)<sub>2</sub>(dtbbpy)]PF<sub>6</sub> (1 mol%) as the photocatalyst. [e] A 59:42 ratio of 17n to unreacted amine 1n was observed. [f] 3.0 equiv of amine 1p was used (N.B. 26% yield with only 1.0 equiv of 1p, with >50% unreacted 1p). [g] ~25% unreacted amine 1s. [h] The amine hydrochloride salt was used and Cs<sub>2</sub>CO<sub>3</sub> (1.0 equiv) was added. [B] Transition states and experimental rate ratio for  $\alpha$ -NH<sub>2</sub> versus  $\alpha$ -OH C-H abstraction by azidyl radical. [C] Scope for  $\alpha$ -C-H dialkylation of primary amines. [i] Used 4CzIPN (1 mol%) as the photocatalyst. [g] Used acrylate 15. Boc = tert-butoxycarbonyl; Tol = 4-tolyl. [D] Scope with respect to the alkene acceptor.

#### WILEY-VCH

## **RESEARCH ARTICLE**

generally stronger or less 'hydridic' than those a- to the unprotected amine, statistical effects (i.e., relative number of C-H bonds) and the impact of reaction conversion on kinetic partitioning will influence this competition. Isolation of the minor by-products in these cases proved fruitless, but the ~3.6 ppm region in the crude NMR spectra did evidence complex mixtures of minor (<5%) singlets (i.e., methyl ester-containing by-products), supportive of acrylate addition to sites other than the amine  $\alpha$ -C-H bond. To further our understanding, we attempted to quantify the relative ease of α-C-H abstraction by azidyl radical (N<sub>3</sub>) from primary amines versus alcohols, as a representative example. DFT calculations using the M06-2X functional indicate that the barrier to HAT is approximately doubled from cyclohexylamine 5 to cyclohexanol 18, and an intermolecular competition experiment between 5 and 18 gave a relative rate ratio of >20:1 (Figure 3B). In a standalone experiment, cyclohexanol 18 itself was α-C-H alkylated in only 12% yield under our conditions, with the remaining mass balance being mainly unreacted 18 (see Supporting Information). Despite this low reactivity of alcohols, we observed incomplete conversion in the case of 4-hydroxycyclohexylamine 1n (i.e., 59:42 ratio of 17n to unreacted **1n**), which may be due to intramolecular hydrogenbonding<sup>[19]</sup> in the syn-diastereomer, raising the barrier to HAT. In some other cases, such as the 4-membered ring substrates **1d**,**p**,**s** (bearing  $\alpha$ -C–H bonds strengthened by ring strain), incomplete conversion of the amine was also primarily responsible for the lower (<50%) isolated yields, as opposed to extensive by-product formation. Finally, a robustness screen<sup>[20]</sup> with 15 different functional group additives was performed, including ketones, alkynes, bromoarenes, nitriles, amides, benzofurans, imidazoles, and pyridines (see Supporting Information), though it should be qualified that these results are only strictly diagnostic of intermolecular competitions.

We next sought to access the prospect of an unprecedented α-C-H dialkylation of primary amines 1f,y-ab bearing two α-C-H bonds, to generate α-tertiary amines 19 directly (Figure 3C). This transformation proved more challenging - possibly due to sidereactions (e.g., telomerisation, aza-Michael) promoted by the higher acrylate concentration - but dialkylated y-lactams 20 could still be isolated in yields of 27–55%.<sup>[21]</sup> The scope of the  $\alpha$ -C–H monoalkylation process with respect to the alkene acceptor (21) was also evaluated (Figure 3D). Acrylates 21a-e bearing α- or β-substituents, including methyl, phenyl, and -N(Boc)<sub>2</sub> were all tolerated, as was an unsaturated y-lactone acceptor (giving 23f). The use of tert-butyl acrylate (21g) led to a y-amino ester product (22g) that proved substantially more resistant to lactamisation. Preliminary examination of non-acrylate partners was also carried out, with vinyl pyridines 21h,i successfully delivering α-alkylated products. No reactivity was observed with unactivated alkenes (i.e., 1-hexene) as the acceptor.

To validate the fact that our chemistry is useful for the rapid and modular assembly of a-tertiary amine derivatives, we next effected a series of N-functionalisations of the y-amino ester products. in lieu of lactamisation (Figure 4A). In order to reduce reaction times and facilitate scale-up, we conducted the reactions in continuous flow.<sup>[22]</sup> using a Vapourtec UV-150 reactor equipped with a 54 W output 420 nm LED lamp. With cyclohexylamine 5 as a representative amine, this led to a straightforward preparation of a series of  $\alpha$ -tertiary amine derivatives **24a**-f. To assess the scalability of our α-C-H alkylation process, we ran the reaction of cyclohexylamine 5 with methyl acrylate 14 on a decagram-scale in continuous flow, using a prototype of the recently commercialised Uniqsis PhotoSyn reactor (Figure 4B). Although the centre wavelength of 455 nm was sub-optimal for our purposes, a space-time yield (STY) of 0.74 mol L<sup>-1</sup> h<sup>-1</sup> for γ-amino ester 8 still proved possible, and we were able to isolate 10.1 g of



Figure 4. [A] Continuous flow synthesis of α-tertiary primary amine derivatives 24. Yields reported for products 24 are w.r.t. the γ-amino ester 8/22g. Fmoc, fluorenylmethoxycarbonyl. [B] Decagram-scale reaction in continuous flow.

product **17b** from a single 5.3 h run, post-lactamisation. Notably, this chemistry provides a scalable access to spirocyclic pyrrolidine building blocks for drug discovery that outcompetes the current most practical synthetic routes for process-scale work (5–7 steps).<sup>[3a]</sup>

Our proposed catalytic cycle for the  $\alpha$ -C–H alkylation process is outlined in Figure 5A. Initial oxidation of azide ion ( $E_{p/2}$  of  $N_3^-$  = +0.87 V vs SCE, in MeCN) by the photoexcited 4CzIPN [ $E_{\frac{1}{2}}$  $(PC^*/PC^{-}) = +1.35 \text{ V vs SCE}$  generates the azidyl radical, N<sub>3</sub><sup>•</sup> a potent oxidant that is capable of hydrogen atom abstraction even from unactivated alkanes.<sup>[23]</sup> This reductive quenching step supported by cyclic voltammetry and Stern-Volmer is luminescence quenching experiments (Figure 5B).<sup>[24]</sup> Subsequent HAT from the relatively weak  $\alpha$ -C-H bond of the primary amine (BDE =  $89-91 \pm 2$  kcal mol<sup>-1</sup>)<sup>[25,26]</sup> is thought to occur selectively, possibly augmented by a polarity-matching effect between the electrophilic N<sub>3</sub><sup>+</sup> radical and the 'hydridic' C-H bond.<sup>[4d]</sup> The resultant  $\alpha$ -amino radical **25** can undergo a rapid and polarity-matched addition to the acrylate acceptor to give an  $\alpha$ -carboxy stabilised radical **26** [ $E_{\gamma_2} \sim -0.63$  V vs SCE<sup>[27]</sup>]. Reduction of this radical to the corresponding enolate 27 ( $pK_{aH}$ ~24 in H<sub>2</sub>O) by the [4CzIPN]<sup>-</sup> radical anion ( $E_{\frac{1}{2}} = -1.21$  V vs SCE, in MeCN) is presumably followed by proton transfer from HN<sub>3</sub>  $(pK_a = 4.72 \text{ in } H_2O)$  to give the y-amino ester product and regenerate the azide ion. Alternatively, a chain process involving



Figure 5. [A] Proposed catalytic mechanism. [B] Evidence for azide ion as reductive quencher.

direct and polarity-matched HAT from the primary amine ( $\alpha$ -C–H BDE = ~90 kcal mol<sup>-1</sup>)<sup>[26]</sup> to the  $\alpha$ -carboxy radical **26** (BDE of  $\alpha$ -CO<sub>2</sub>Me C–H ~96 kcal mol<sup>-1</sup>]<sup>[26]</sup> is plausible.<sup>[6c,23a]</sup> However, the low quantum yield for product formation ( $\Phi$  = 0.04) rules out the presence of efficient chain processes, although it does not exclude the possibility of an inefficient photoredox process

followed by a short radical chain, where quantum yields can be <1.0.

#### Conclusion

In summary, we have developed the first visible-light photocatalysed  $\alpha$ -C–H alkylation of primary aliphatic amines with electrophilic alkenes that does not rely on *in situ* masking of the amino group's intrinsic reactivity. Our dual catalytic approach, which uses azide ion as a HAT catalyst, is amenable to the decagram-scale preparation of hitherto difficult-to-access  $\alpha$ -tertiary amines and aza(spiro)cyclic building blocks.<sup>[3a]</sup> We anticipate that this technology will open up new retrosynthetic strategies for the disconnection of substituted amine derivatives in organic synthesis, and find immediate application in conventional and fragment-based lead identification programmes.

#### Acknowledgements

This work was supported by the Engineering and Physical Sciences Research Council EP/L016354/1. A.J.C. thanks the Royal Society for a University Research Fellowship (UF150533) and a PhD studentship (G.B.), the University of Bath for generous financial support and a PhD studentship (A.G.K.), the Centre for Sustainable Chemical Technologies (CSCT) for an EPSRC CDT PhD studentship (A.S.H.R.), the EPSRC and AstraZeneca for an iCASE PhD studentship (T.M.), and the Dial-a-Molecule network for proof-of-concept funding. We thank Dr Catherine Lyall for assistance with flow NMR, Dr Adam Noble for assistance with quantum yield measurements, and Dr Josh Tibbetts for assisting with experiments. We also thank Duncan Guthrie at Vapourtec for the loan of a high-powered LED lamp unit, and Mark Ladlow from Uniqsis Ltd for the loan of a PhotoSyn reactor.

**Keywords:** Amines • Photocatalysis • C-H activation • Radicals • Spiro compounds

- (a) K. R. Campos, P. J. Coleman, J. C. Alvarez, S. D. Dreher, R. M. Garbaccio, N. K. Terrett, R. D. Tillyer, M. D. Truppo, E. R. Parmee, *Science* 2019, *363*, eaat0805. (b) D. C. Blakemore, L. Castro, I. Churcher, C. D. Rees, A. W. Thomas, D. M. Wilson, A. Wood, *Nat. Chem.* 2018, *10*, 383–394. (c) S. D. Roughley, A. M. Jordan, *J. Med. Chem.* 2011, *54*, 3451–3479.
- [2] E. Vitaku, D. T. Smith, J. T. Njardarson, J. Med. Chem. 2014, 57, 10257– 10274.
- (a) K. P. Melnykov, A. N. Artemenko, B. O. Ivanenko, Y. M. Sokolenko,
   P. S. Nosik, E. N. Ostapchuk, O. O. Grygorenko, D. M. Volochnyuk, S.
   V. Ryabukhin, ACS Omega. 2019, 4, 7498–7515. (b) N. J. Flodén, A.
   Trowbridge, D. Willcox, S. M. Walton, Y. Kim, M. J. Gaunt, J. Am. Chem.
   Soc. 2019, 141, 8426–8430. (c) E. M. Carreira, T. C. Fessard, Chem.
   Rev. 2014, 114, 8257–8322. (d) Y.-J. Zheng, C. M. Tice, S. B. Singh,
   Bioorg. Med. Chem. Lett. 2014, 24, 3673–3682.
- Selected examples: (a) A. Ruffoni, F. Juliá, T. D. Svejstrup, A. McMillan, J. J. Douglas, D. Leonori, *Nat. Chem.* 2019, *11*, 426–433. (b) A. Paul, D. Seidel, *J. Am. Chem. Soc.* 2019, *141*, 8778–8782. (c) A. J. Musacchio, B. C. Lainhart, X. Zhang, S. G. Naguib, T. C. Sherwood, R. R. Knowles, *Science* 2017, *355*, 727–730. (d) C. Le, Y. Liang, R. W. Evans, X. Li, D.

W. C. MacMillan, *Nature* **2017**, *547*, 79–83. (e) Y. Yang, S.-L. Shi, D. Niu, P. Liu, S. L. Buchwald, *Science* **2015**, *349*, 62–66.

- (a) A. Trowbridge, S. M. Walton, M. J. Gaunt, *Chem. Rev.* 2020, DOI: 10.1021/acs.chemrev.9b00462. (b) C. He, W. G. Whitehurst, M. J. Gaunt, *Chem.* 2019, *5*, 1031–1058.
- [6] (a) H. G. Richey, Jr., W. F. Erickson, J. Org. Chem. 1983, 48, 4349–4357.
  (b) D. H. R. Barton, G. Lamotte, W. B. Motherwell, S. C. Narang, J. Chem. Soc. Perkin Trans. 1 1978, 2030–2033. (c) W. H. Urry, O. O. Juveland, J. Am. Chem. Soc. 1958, 80, 3322–3328.
- [7] D. Vasu, A. L. Fuentes de Arriba, J. Leitch, A. de Gombert, D. J. Dixon, *Chem. Sci.* 2019, *10*, 3401–3407.
- [8] With Ti/Zr catalysis (intramolecular): (a) E. Chong, L. L. Schafer, *Org. Lett.* 2013, *15*, 6002–6005. (b) J. A. Bexrud, P. Eisenberger, D. C. Leitch, P. R. Payne, L. L. Schafer, *J. Am. Chem. Soc.* 2009, *131*, 2116–2118. (c) R. Kubiak, I. Prochnow, S. Doye, *Angew. Chem. Int. Ed.* 2009, *48*, 1153–1156. With UV-photocatalysis (intermolecular, and low yields of γ-lactams only): (d) S. Das, J. S. Dileep Kumar, K. George Thomas, K. Shivaramayya, M. V. George, *J. Org. Chem.* 1994, *59*, 628–634. (e) M. Pfau, R. Dulou, *Bull. Soc. Chim. Fr.* 1967, 3336–3342.
- [9] (a) M. A. Ashley, C. Yamauchi, J. C. K. Chu, S. Otsuka, H. Yorimitsu, T. Rovis, *Angew. Chem. Int. Ed.* 2019, *58*, 4002–4006. (b) A. W. Rand, H. Yin, L. Xu, J. Giacoboni, R. Martin-Montero, C. Romano, J. Montgomery, R. Martin, *ACS Catal.* 2020, *10*, 4671–4676.
- [10] J. Ye, I. Kalvet, F. Schoenebeck, T. Rovis, *Nature Chem.* 2018, 10, 1037–1041.
- [11] (a) L. Marzo, S. K. Pagire, O. Reiser, B. König, *Angew. Chem. Int. Ed.* 2018, 57, 10034–10072. (b) *Visible light photocatalysis in organic chemistry* (Eds.: C. R. J. Stephenson, T. Yoon, D. W. C. MacMillan), Wiley-VCH, Berlin, 2018. (c) N. A. Romero, D. A. Nicewicz, *Chem. Rev.* 2016, *116*, 10075–10166. (d) C. Prier, D. Rankic, D. W. C. MacMillan, *Chem. Rev.* 2013, *113*, 5322–5363.
- [12] (a) For reviews, see: Y.-Q. Zou, W.-J. Xiao in Visible light photocatalysis in organic chemistry (Eds.: C. R. J. Stephenson, T. Yoon, D. W. C. MacMillan), Wiley-VCH, Berlin, 2018, Chap. 4, pp 93–128. (b) J. W. Beatty, C. R. J. Stephenson, Acc. Chem. Res. 2015, 48, 1474–1484. (c) K. Nakajima, Y. Miyake, Y. Nishibayashi, Acc. Chem. Res. 2016, 49, 1946–1956. For selected examples, see: (d) A. McNally, C. Prier, D. W. C. MacMillan, Science 2011, 334, 1114–1117. (e) Y. Miyake, K. Nakajima, Y. Nishibayashi, J. Am. Chem. Soc. 2012, 134, 3338–3341. (f) Z. Zuo, D. T. Ahneman, L. Chu, J. A. Terrett, A. G. Doyle, D. W. C.

MacMillan, *Science* **2014**, *345*, 437–440. (g) M. Shaw, J. Shurtleff, J. A. Terrett, J. D. Cuthbertson, D. W. C. MacMillan, *Science* **2016**, *352*, 1304–1308. (h) J. B. McManus, N. P. R. Onuska, D. A. Nicewicz, *J. Am. Chem. Soc.* **2018**, *140*, 9056–9060.

- [13] A. Trowbridge, D. Reich, M. J. Gaunt, Nature 2018, 561, 522–527.
- [14] Selected examples of photoredox-HAT catalysis: (a) T. Ide, J. P. Barham, M. Fujita, K. Kawato, H. Egami, Y. Hamashima, *Chem. Sci.* 2018, *9*, 8453–8460. (b) J. L. Jeffrey, J. A. Terrett, D. W. C. MacMillan, *Science* 2015, 349, 1532–1536.
- [15] (a) S. M. Engle, T. R. Kirkner, C. B. Kelly, Org. Synth. 2019, 96, 455– 473. (b) J. Luo, J. Zhang, ACS Catal. 2016, 6, 873–877.
- [16] The half-wave redox potential for the N<sub>3</sub>/N<sub>3</sub><sup>-</sup> couple in neutral aqueous solution is reported as +1.32 V vs NHE (+1.08 V vs SCE); see: Z. B. Alfassi, A. Harriman, R. E. Huie, S. Mosseri, P. Neta, *J. Phys. Chem.* **1987**, *91*, 2120–2122.
- [17] A. M. R. Hall, J. C. Chouler, A. Codina, P. T. Gierth, J. P. Lowe, U. Hintermair, *Catal. Sci. Technol.* 2016, *6*, 8406–8417.
- [18] Use of 15 gave easier chromatographic separation of ester-containing by-products in some cases.
- [19] Stolow, R. D. J. Am. Chem. Soc. 1961, 83, 2592–2593.
- [20] K. D. Collins, F. Glorius, Nature Chem. 2013, 5, 597–601.
- [21] The Ir[(dF(CF<sub>3</sub>)ppy)<sub>2</sub>(dtbbpy)]PF<sub>6</sub> photocatalyst was used in place of 4CzIPN for amines **1y–ab**, informed by earlier studies that attempted to optimise for monoalkylation (see pg S53).
- [22] (a) C. Sambiagio, T. Noël, *Trends Chem.* 2020, *2*, 92–106. (b) Y. Su, N. J. W. Straathof, V. Hessel, T. Noël, *Chem. Eur. J.* 2014, *20*, 10562–10589.
- [23] (a) Y. Wang, X. Hu, C. A. Morales-Rivera, G.-X. Li, X. Huang, G. He, P. Liu, G. Chen, *J. Am. Chem. Soc.* 2018, *140*, 9678–9684. (b) X. Huang, J. T. Groves, *ACS Catal.* 2016, *6*, 751–759.
- [24] L. Buzzetti, G. E. M. Crisenza, P. Melchiorre, Angew. Chem. Int. Ed. 2019, 58, 3730–3747.
- [25] I. MacInnes, J. C. Walton, D. C. Nonhebel, J. Chem. Soc. Perkin Trans. 2 1987, 1789–1794.
- [26] Y.-R Luo, Comprehensive Handbook of Chemical Bond Energies, CRC Press: Boca Raton, FL, 2007.
- [27] N. Bortolamei, A. A. Isse, A. Gennaro, *Electrochim. Acta* 2010, 55, 8312– 8318.

#### WILEY-VCH

## **RESEARCH ARTICLE**

#### **Entry for the Table of Contents**

Insert graphic for Table of Contents here.



Catalytic strategies for the  $\alpha$ -C–H functionalisation of primary amines are a major challenge in organic synthesis. We report a photocatalytic protocol for the  $\alpha$ -C–H alkylation of unprotected primary amines that is amenable to the direct synthesis of  $\alpha$ -tertiary primary amines. This process is readily scalable in continuous flow to provide access to decagram quantities of valuable  $\gamma$ -lactams and azaspirocycles, for application in drug discovery.

Institute and/or researcher Twitter usernames: @UniofBath, @UniofBathSci, @BathChem, @csctbath