

Ni(0)-Catalyzed Formation of Azaaluminacyclopentenes via Azanickelacyclopentenes: A Unique Nickel/Aluminum Double Transmetalation Reaction

Masato Ohashi, Osamu Kishizaki, Haruo Ikeda, and Sensuke Ogoshi*


Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Suita, Osaka 565-0871, Japan

Received April 21, 2009; E-mail: ogoshi@chem.eng.osaka-u.ac.jp

Oxidative cyclization of two π components with low-valent transition metals has been the focus of a great deal of attention because of its promise as a method for constructing C–C bonds between a variety of functional groups.¹ Among these methods, the formation of heteronickelacycles is assumed to be a key step in multicomponent nickel-catalyzed coupling reactions.^{2–6} We recently developed the nickel-catalyzed [2 + 2 + 2] cycloaddition reaction of two alkynes and an imine to afford 1,2-dihydropyridines, in which the oxidative cyclization of an alkyne and an imine was proposed as a key step.^{6c} The formation of a key intermediate (**1a**) by oxidative cyclization of diphenylacetylene and *N*-benzylidene benzenesulfonamide (*N*-BBSA) with Ni(0) was observed.^{6c} Aza-nickelacycles were also assumed to be an important intermediate in the three-component reaction of an alkyne, an imine, and an alkylmetal reagent,⁴ and therefore, investigation of the reactivity of **1a** toward alkylmetal reagents was warranted. Herein, we report the stoichiometric reactions of **1a** with ZnMe₂ and AlMe₃. Treatment with ZnMe₂ gave a three-component coupling product, while the reaction with AlMe₃ resulted in the unexpected formation of an azaaluminacyclopentene by nickel/aluminum double transmetalation. We also achieved the construction of a novel nickel-catalyzed three-component cyclocondensation of an alkyne, an imine, and AlMe₃.

The reaction of **1a** with ZnMe₂ in toluene at room temperature in the presence of 2 equiv of TMSCH=CH₂ afforded the expected methylzincamido product **2a** in 74% isolated yield together with the formation of (PCy₃)Ni(CH₂=CHTMS)₂ (**3**) (Scheme 1).⁷ In addition, the three-component coupling reaction of diphenylacetylene, *N*-BBSA, and ZnMe₂, which afforded **2a**, also proceeded in the presence of Ni(cod)₂ and PCy₃ (10 and 20 mol %, respectively). Although **1a** was analogous to the reaction intermediate proposed in Jamison's work,⁴ **1a** did not react with BEt₃ even when heated at 60 °C for 2 h. This is consistent with the fact that *N*-tosyl imines do not react in the Ni-catalyzed three-component coupling of alkyne, imine, and triethylborane.⁴

Scheme 1. Reaction of **1a** with Alkylmetal Reagents



In contrast, treatment of **1a** with an equimolar amount of AlMe₃ under identical reaction conditions unpredictably gave the five-membered azaaluminacycle **4a** in 69% isolated yield (Scheme 1). Monitoring of the reaction by means of ¹H NMR spectroscopy indicated a concomitant generation of ethane (δ_{H} 0.80 in C₆D₆) and **3**. X-ray crystallography of **4a** showed that the aluminum atom





Figure 1. Molecular structure of **4a**, with thermal ellipsoids at the 30% probability level. H atoms and the solvated molecule (toluene) have been omitted for clarity. Symmetry transformation used to generate equivalent atoms S*: 2 – X, Y, 0.5 – Z.

was covalently bonded to both the carbon and nitrogen atoms to form a five-membered ring and that one methyl group also was bound to the aluminum center (Figure 1).

In the crystal lattice, the five-membered azaaluminacycle unit formed a dimer in which one of the oxygen atoms in the benzenesulfonyl group of **4a** is coordinated to the other aluminum atom. Unlike the three-component coupling product **2a**, azaaluminacycle **4a** is an organometallic reagent in which the Al–C bond can react with electrophiles. Indeed, **4a** can be treated with electrophiles such as proton and halogenonium to afford allylamine derivatives **5a–7a** (Scheme 2). To the best of our knowledge, this is the first example of the catalytic formation of azaaluminacyclopentenes, although cycloaluminumation of either olefins or acetylenes mediated by Cp₂Zr derivatives has been used in the preparation of organoaluminum compounds.^{8,9}

Scheme 2. Transformation of **4a** into Allylamine Derivatives^a



^a Reaction conditions: for **5a**, MeOH/toluene; for **6a**, NBS (2 equiv) in MeCN, 0.5 h, then MeOH; for **7a**, I₂ (excess) in CH₃CN, 2 h, then HCl.

As anticipated from the regeneration of the Ni(0) complex **3**, the transformation of diphenylacetylene, *N*-BBSA, and AlMe₃ into **4a**, in which the oxidative cyclization of diphenylacetylene and *N*-BBSA is a key step, proceeded catalytically. However, as shown by the equation in Table 1, the addition reaction of AlMe₃ and *N*-BBSA, which gave the corresponding amide **8**, also took place


Table 1. Nickel-Mediated Cyclocondensation of Alkynes, *N*-BBSA, and AlMe₃^a

run	alkyne		product	yield (%) ^b	yield of 8 (%) ^c
	R ¹	R ²			
1	Ph	Ph	4a	86 (71)	—
2	<i>p</i> -MeC ₆ H ₄	<i>p</i> -MeC ₆ H ₄	4b	85 (82)	—
3	<i>p</i> -CF ₃ C ₆ H ₄	<i>p</i> -CF ₃ C ₆ H ₄	4c	90 (99)	—
4 ^d	Ph	TMS	4d	85 (73)	7
5 ^e	Ph	Me	4e^f	65 (44)	13
6	Et	Et	4f^g	27	59

^a General conditions: alkyne and *N*-BBSA (0.3 mmol each), solvent (10 mL). After dropwise addition of AlMe₃, the reaction mixture was stirred until the color derived from **1** (typically purple) disappeared (1.0–6.0 h). ^b Isolated yields as allylamines **5** after protolysis. Cited yields in parentheses are of isolated **4**. ^c The cited yields, determined by ¹H NMR, were of the corresponding protonated products. ^d The reaction was carried out using 1.5 mmol of TMS-C≡Ph. ^e The reactions were carried out with concomitant addition of AlMe₃ and the alkyne. ^f The minor regioisomer (11%) was also obtained. ^g Formation of a 1,2-dihydropyridine derivative was also observed (also see ref 6e).

and was found to accelerate with the addition of 10 mol % Ni(cod)₂ and PCy₃ (Scheme 3, right circle).¹⁰ Therefore, the slow addition of AlMe₃ improved the yield of **4a** as a result of suppression of the competitive reaction. Finally, the three-component cyclocondensation of *N*-BBSA, diphenylacetylene, and AlMe₃ (via slow addition over 30 min) in the presence of 10 mol % Ni(cod)₂ and PCy₃ afforded **4a** in 71% isolated yield (Table 1, run 1).¹¹ While the isolated yield of **4a** was somewhat decreased because of losses during the purification process, NMR analysis indicated that this catalytic reaction proceeded quantitatively. In fact, protolysis of the crude product gave the corresponding allylamine **5a** in 86% isolated yield (run 1). The same reaction conditions were applied successfully to diphenylacetylene derivatives, which led to clean formation of **4b** and **4c** (runs 2 and 3, respectively).

Nonsymmetric alkynes were used as coupling components in the cyclocondensation with *N*-BBSA and AlMe₃; **4d** was formed in 85% yield as a single regioisomer, but use of 5 equiv of 1-phenyl-2-trimethylsilylacetylene was essential for the promotion of a smooth reaction, probably as a result of its diminished ability to coordinate to Ni(0) (run 4). In contrast, the reaction with 1-phenyl-1-propyne gave **4e** with 86:14 regioselectivity only when slow addition of both AlMe₃ and the alkyne was conducted in order to circumvent the insertion of the second alkyne into a five-membered

Scheme 3. Plausible Mechanism for Catalytic Formation of **4**

azanickelacycle (run 5).^{6c} 1,2-Dialkyl-substituted alkynes such as 3-hexyne did not react efficiently because of the rapid formation of the undesired seven-membered azanickelacycle even with slow addition of a mixture of the alkyne and AlMe₃ (run 6).^{6c} For several cases cited in runs 4, 5, and 6, formation of **8** was observed in the ¹H NMR spectra of the crude products. These results are consistent with the mechanism proposed in Scheme 3.

In summary, we have demonstrated the nickel-catalyzed three-component cyclocondensation of imines, alkynes, and AlMe₃ to yield unique azaaluminacyclopentenenes. Nickelacycle **1**, generated by the oxidative cyclization of an alkyne and an imine, is a key intermediate in the cyclocondensation reaction as well as in the three-component coupling reaction with ZnMe₂. Further study will focus on the related formation of oxaaluminacyclopentenenes.

Acknowledgment. The authors are grateful for financial support through Grants-in-Aid for Scientific Research (21245028 and 21750102) and a Grant-in-Aid for Scientific Research on Priority Areas (19028038, Chemistry of Concerto Catalysis) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

Supporting Information Available: Detailed experimental procedures, analytical and spectral data for all new compounds, and crystallographic data (CIF) for **2a** and **4a**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Recent reviews: (a) Ikeda, S.-I. *Angew. Chem., Int. Ed.* **2003**, *42*, 5120. (b) Montgomery, J. *Angew. Chem., Int. Ed.* **2004**, *43*, 3890. (c) Moslin, R. M.; Miller-Moslin, K.; Jamison, T. F. *Chem. Commun.* **2007**, 4441. (d) Skucas, E.; Ngai, M.-Y.; Komanduri, V.; Kricheldorf, M. J. *Acc. Chem. Res.* **2007**, *40*, 1394. (e) Rayabarapu, D. K.; Cheng, C.-H. *Acc. Chem. Res.* **2007**, *40*, 971.
- (2) Alkyne with an aldehyde/ketone: (a) Tsuda, T.; Kiyoi, T.; Miyane, T.; Saegusa, T. *J. Am. Chem. Soc.* **1988**, *110*, 8570. (b) Oblinger, E.; Montgomery, J. *J. Am. Chem. Soc.* **1997**, *119*, 9065. (c) Huang, W.-S.; Chan, J.; Jamison, T. F. *Org. Lett.* **2000**, *2*, 4221. (d) Ni, Y.; Amarasinghe, K. K. D.; Montgomery, J. *Org. Lett.* **2002**, *4*, 1743. (e) Miller, K. M.; Huang, W.-S.; Jamison, T. F. *J. Am. Chem. Soc.* **2003**, *125*, 3442. (f) Mahandru, G. M.; Liu, G.; Montgomery, J. *J. Am. Chem. Soc.* **2004**, *126*, 3698. (g) Miller, K. M.; Jamison, T. F. *J. Am. Chem. Soc.* **2004**, *126*, 15342. (h) Murakami, M.; Ashida, S.; Matsuda, T. *J. Am. Chem. Soc.* **2005**, *127*, 6932. (i) Luanphaisarnont, T.; Ndubaku, C. O.; Jamison, T. F. *Org. Lett.* **2005**, *7*, 2937. (j) Miller, K. M.; Jamison, T. F. *Org. Lett.* **2005**, *7*, 3077. (k) Tekevac, T. N.; Louie, J. *Org. Lett.* **2005**, *7*, 4037. (l) Knapp-Reed, B.; Mahandru, G. M.; Montgomery, J. *J. Am. Chem. Soc.* **2005**, *127*, 13156. (m) Chaulagain, M. R.; Mahandru, G. M.; Montgomery, J. *Tetrahedron* **2006**, *62*, 7560. (n) Murakami, M.; Ashida, S.; Matsuda, T. *J. Am. Chem. Soc.* **2006**, *128*, 2166. (o) Chaulagain, M. R.; Sormunen, G. J.; Montgomery, J. *J. Am. Chem. Soc.* **2007**, *129*, 9568. (p) Baxter, R. D.; Montgomery, J. *J. Am. Chem. Soc.* **2008**, *130*, 9662.
- (3) Alkene with an aldehyde/ketone: (a) Ng, S.-S.; Jamison, T. F. *J. Am. Chem. Soc.* **2005**, *127*, 14194. (b) Ho, C.-Y.; Ng, S.-S.; Jamison, T. F. *J. Am. Chem. Soc.* **2006**, *128*, 5362. (c) Ng, S.-S.; Ho, C.-Y.; Jamison, T. F. *J. Am. Chem. Soc.* **2006**, *128*, 11513. (d) Ho, C.-Y.; Jamison, T. F. *Angew. Chem., Int. Ed.* **2007**, *46*, 782.
- (4) Alkyne with an imine: (a) Patel, S. J.; Jamison, T. F. *Angew. Chem., Int. Ed.* **2003**, *42*, 1364. (b) Patel, S. J.; Jamison, T. F. *Angew. Chem., Int. Ed.* **2004**, *43*, 3941.
- (5) Alkene with an imine: Yeh, C.-H.; Korivi, R. P.; Cheng, C.-H. *Angew. Chem., Int. Ed.* **2008**, *47*, 4892.
- (6) Isolated examples of heteronickelacycles proposed as reaction intermediates: (a) Ogoshi, S.; Oka, M.-a.; Kurosawa, H. *J. Am. Chem. Soc.* **2004**, *126*, 11082. (b) Ogoshi, S.; Ueta, M.; Arai, T.; Kurosawa, H. *J. Am. Chem. Soc.* **2005**, *127*, 12810. (c) Ogoshi, S.; Nagata, M.; Kurosawa, H. *J. Am. Chem. Soc.* **2006**, *128*, 5350. (d) Ogoshi, S.; Tonomori, K.-i.; Oka, M.-a.; Kurosawa, H. *J. Am. Chem. Soc.* **2006**, *128*, 7077. (e) Ogoshi, S.; Ikeda, H.; Kurosawa, H. *Angew. Chem., Int. Ed.* **2007**, *46*, 4930. (f) Ogoshi, S.; Arai, T.; Ohashi, M.; Kurosawa, H. *Chem. Commun.* **2008**, 1347.
- (7) An ORTEP drawing of **2a** is given in the Supporting Information.
- (8) Selected reviews: (a) Dzhemilev, U. M.; Ibragimov, A. G. *Russ. Chem. Rev.* **2000**, *69*, 121. (b) Dzhemilev, U. M.; Ibragimov, A. G. *J. Organomet. Chem.* **1994**, *466*, 1.
- (9) Oxasilacyclopentenenes were very recently prepared via Ni-catalyzed cyclocondensation of aldehydes, alkynes, and dialkylsilanes (see ref 2p).
- (10) Biswas, K.; Prieto, O.; Goldsmith, P. J.; Woodward, S. *Angew. Chem., Int. Ed.* **2005**, *44*, 2232.
- (11) In contrast, slow addition of ZnMe₂ was unnecessary because the corresponding direct addition in the Ni-catalyzed three-component coupling reaction did not occur.

JA903175G