

Lewis Acid Catalyzed Rearrangement of Vinylcyclopropenes for the Construction of Naphthalene and Indene Skeletons


Li-Xiong Shao,[†] Yun-Peng Zhang,[†] Ming-Hui Qi,[†] and Min Shi^{*,†,‡}

School of Chemistry and Molecular Engineering, East China University of Science and Technology, 130 Meilong Road, Shanghai, China 200237, and State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Fenglin Road, Shanghai, China 200032

mshi@mail.sioc.ac.cn

Received November 2, 2006

ABSTRACT



The choice of Lewis acid catalyst can result in dramatic differences in the chemoselectivity of the rearrangement reactions of vinylcyclopropenes. When $\text{BF}_3 \cdot \text{OEt}_2$ was used as the catalyst, naphthalenes were formed. However, when $\text{Cu}(\text{OTf})_2$ was used as the catalyst, indenes were obtained.

Thermal and photochemical skeleton rearrangements of strained small rings with multiple bonds and functional groups have attracted much attention from both synthetic and mechanistic viewpoints. However, only a few examples of such thermal¹ and photochemical² skeletal conversions of vinylidenecyclopropanes **1** have been reported. Previously, we observed that naphthalene, 6*a*H-benzo[*c*]fluorine, and indene derivatives can be obtained by Lewis acid catalyzed rearrangement of arylvinylidenecyclopropanes **1** depending on the substitution pattern on the cyclopropyl ring and the electronic nature of aryl groups.³ In this paper, we wish to report the Lewis acid catalyzed rearrangement of vinylcyclopropenes **2**, which are derived from the corresponding vinylidenecyclopropanes **1** under basic conditions, to afford naphthalene and indene skeletons in good to high yields under mild conditions.

As shown in Table 1, vinylcyclopropenes **2** are readily accessible from simple isomerization of the corresponding vinylidenecyclopropanes **1** under basic conditions. For all

As shown in Table 1, vinylcyclopropenes **2** are readily accessible from simple isomerization of the corresponding vinylidenecyclopropanes **1** under basic conditions. For all

(2) (a) Akasaka, T.; Misawa, Y.; Ando, W. *Tetrahedron Lett.* **1990**, *31*, 1173. (b) Mizuno, K.; Sugita, H.; Isagawa, K.; Goto, M.; Otsuji, Y. *Tetrahedron Lett.* **1993**, *34*, 5737. (c) Mizuno, K.; Nire, K.; Sugita, H.; Otsuji, Y. *Tetrahedron Lett.* **1993**, *34*, 6563. (d) Mizuno, K.; Sugita, H.; Hirai, T.; Maeda, H. *Chem. Lett.* **2000**, 1144. (e) Mizuno, K.; Nire, K.; Sugita, H.; Maeda, H. *Tetrahedron Lett.* **2001**, *42*, 2689. (f) Mizuno, K.; Maeda, H.; Sugita, H.; Nishioka, S.; Hirai, T.; Sugimoto, A. *Org. Lett.* **2001**, *3*, 581. (g) Mizuno, K.; Sugita, H.; Hirai, T.; Maeda, H.; Otsuji, Y.; Yasuda, M.; Hashiguchi, M.; Shima, K. *Tetrahedron Lett.* **2001**, *42*, 3363. (h) Maeda, H.; Zhen, L.; Hirai, T.; Mizuno, K. *ITE Lett. Batteries, New Technol. Med.* **2002**, *3*, 485.


(3) (a) Shi, M.; Lu, J.-M. *J. Org. Chem.* **2006**, *71*, 1920. (b) Shi, M.; Lu, J.-M. *Synlett* **2005**, 2352. (c) Shi, M.; Ma, M.; Shao, L.-X. *Tetrahedron Lett.* **2005**, *46*, 7609. (d) Shi, M.; Lu, J.-M.; Xu, G.-C. *Tetrahedron Lett.* **2005**, *46*, 4745. (e) Xu, G.-C.; Ma, M.; Liu, L.-P.; Shi, M. *Synlett* **2005**, 1869. (f) Xu, G.-C.; Liu, L.-P.; Lu, J.-M.; Shi, M. *J. Am. Chem. Soc.* **2005**, *127*, 14552.





[†] East China University of Science and Technology.

[‡] Shanghai Institute of Organic Chemistry.

(1) (a) Paulson, D. R.; Crandall, J. K.; Bunnell, C. A. *J. Org. Chem.* **1970**, *35*, 3708. (b) Jones, M., Jr.; Hendrick, M. E.; Hardie, J. A. *J. Org. Chem.* **1971**, *36*, 3061. (c) Patrick, T. B.; Haynie, E. C.; Probst, W. J. *J. Org. Chem.* **1972**, *37*, 1553. (d) Sadler, I. H.; Stewart, J. A. G. *J. Chem. Soc., Perkin Trans. 2* **1973**, 278. (e) Sugita, H.; Mizuno, K.; Saito, T.; Isagawa, K.; Otsuji, Y. *Tetrahedron Lett.* **1992**, *33*, 2539. (f) Mizuno, K.; Sugita, H.; Kamada, T.; Otsuji, Y. *Chem. Lett.* **1994**, 449. (g) Maeda, H.; Hirai, T.; Sugimoto, A.; Mizuno, K. *J. Org. Chem.* **2003**, *68*, 7700.

Table 1. Isomerization of Vinylidenecyclopropanes **1** to Vinylcyclopropanes **2**



entry ^a	1 (R ¹ /R ² /R ³ /R ⁴)	yield (%) ^b
1	1a (C ₆ H ₅ /C ₆ H ₅ /Me/Me)	2a , 92
2	1b (4-FC ₆ H ₄ /C ₆ H ₅ /Me/Me)	2b , 63
3	1c (4-ClC ₆ H ₄ /C ₆ H ₅ /Me/Me)	2c , 81
4	1d (4-MeC ₆ H ₄ /C ₆ H ₅ /Me/Me)	2d , 79
5	1e (C ₆ H ₅ /4-ClC ₆ H ₄ /Me/Me)	2e , 93
6	1f (C ₆ H ₅ /4-BrC ₆ H ₄ /Me/Me)	2f , 93
7	1g (C ₆ H ₅ /4-MeC ₆ H ₄ /Me/Me)	2g , 85
8	1h (C ₆ H ₅ /C ₆ H ₅ /Me/H)	2h , 61
9	1i (C ₆ H ₅ /C ₆ H ₅ /H/H)	2i , 68
10		
11		

^a All reactions were carried out using **1** (0.2 mmol), NaOH (20.0 equiv), and Bu₄NHSO₄ (2.0 equiv) in toluene (5.0 mL) at 60 °C for 5 h. ^b Isolated yields.

vinylidenecyclopropanes **1** examined, the isomerization reaction proceeded smoothly to give vinylcyclopropanes **2** in good to high yields within 5 h under mild conditions (Table 1). Compound **2b** was unambiguously determined by an X-ray diffraction (Figure 1).⁴





Figure 1. ORTEP drawing of compound **2b**.

Initial examinations using vinylcyclopropene **2a** (R¹ = R² = C₆H₅; R³ = R⁴ = Me) as the substrate in the presence of various Brønsted acids, such as trifluoromethanesulfonic acid (TfOH), CF₃CO₂H, CH₃CO₂H, and the Lewis acid BF₃·OEt₂,

were aimed at determining the optimal conditions for rearrangement to **3a**. The results are summarized in the Supporting Information. It was found that BF₃·OEt₂ was the best catalyst and that 1,2-dichloroethane (DCE) was the solvent of choice, and **3a** was produced in 60% yield at 40 °C (Scheme 1) (see the Supporting Information for details). The

Scheme 1. Optimal Conditions for the Formation of Naphthalene **3a**



generality of this transformation was examined, and we found that for all of the employed substrates (**2**) the corresponding rearranged product **3** could be obtained in good yield (Table 2).

A plausible mechanism for the BF₃·OEt₂-catalyzed rearrangement of vinylcyclopropanes **2** to naphthalenes **3** is outlined in Scheme 1. Initially, reaction of BF₃·OEt₂ with trace amounts of water in the system generates the Brønsted acid catalyst **A**,⁵ which is similar to the Brønsted acid TfOH. The reaction of catalyst **A** with **2** produces cationic intermediate **B** (the corresponding counteranion in all of the intermediates has been omitted for convenience), which will rearrange to either ring-opened cationic intermediate **C** or resonance-stabilized intermediate **C'** via σ -bond *b* cleavage.⁶ Subsequent intramolecular Friedel–Crafts reaction of **C'** with the aromatic group R¹ gives the cyclized intermediate **D**, which affords intermediate **E** via deprotonation. Finally, aromatization of intermediate **E** furnishes the corresponding naphthalene **3** (Scheme 2).

Interestingly, we found that vinylcyclopropanes **2** can rearrange to the corresponding indene derivatives **4** in the

(4) The crystal data of **2b** have been deposited in the CCDC with number 602862. Empirical formula: C₂₅H₂₀F₂. Formula weight: 358.41. Crystal color, habit: colorless, prismatic. Crystal system: monoclinic. Lattice type: primitive. Lattice parameters: *a* = 12.5896(12) Å, *b* = 9.1294(8) Å, *c* = 17.6833(16) Å, α = 90°, β = 102.192(2)°, γ = 90°, *V* = 1986.6(3) Å³. Space group: *P*2(1)/*n*. *Z* = 4; *D*_c = 1.198 g/cm³; *F*₀₀₀ = 752. Diffractometer: Rigaku AFC7R. Residuals: *R*, *R*_w: 0.0431, 0.1098.

(5) For related papers, which report that trace amounts of water are required to the formation of cation **B** with BF₃ (BF₃ + H₂O + Me₂CH = CH₂ → Me₃CH⁺BF₃OH⁻), see: (a) Evans, A. G.; Holden, D.; Plesch, P.; Polanyi, M.; Skinner, H. A.; Weinberger, M. A. *Nature* **1946**, *157*, 102. (b) Evans, A. G.; Meadows, G. W.; Polanyi, M.; Skinner, H. A.; Weinberger, M. A. *Nature* **1946**, *158*, 94. (c) Evans, A. G.; Polanyi, M. *J. Chem. Soc.* **1947**, 252. (d) Prakash, G. K. S.; Mathew, T.; Hoole, D.; Esteves, P. M.; Wang, Q.; Rasul, G.; Olah, G. A. *J. Am. Chem. Soc.* **2004**, *126*, 15770. The rearrangement of **2a** in anhydrous DCE was also studied by bubbling BF₃ gas into the reaction solution by means of a Schlenk tube, under an argon atmosphere. The reaction proceeded slowly under these conditions. However, when the solution was exposed to the ambient atmosphere for several minutes, the reaction proceeded smoothly to give product **3a** in good yield.

(6) Reaction of electrophilic reagents with strained bonds has been extensively studied. See: (a) Bishop, K. C. *Chem. Rev.* **1976**, *76*, 461. (b) Leftin, J. H.; Gil-Av, G. *Tetrahedron Lett.* **1972**, *13*, 3368. (c) Padwa, A.; Rieker, W. F.; Rosenthal, R. J. *J. Am. Chem. Soc.* **1985**, *107*, 1710. (d) Padwa, A.; Blacklock, T. J.; Loza, R. *J. Am. Chem. Soc.* **1981**, *103*, 2404.

Table 2. $\text{BF}_3 \cdot \text{OEt}_2$ -Catalyzed Rearrangement of **2** for the Formation of Naphthalenes **3**




entry ^a	2 ($\text{R}^1/\text{R}^2/\text{R}^3/\text{R}^4$)	yield (%) ^b
1	2a ($\text{C}_6\text{H}_5/\text{C}_6\text{H}_5/\text{Me}/\text{Me}$)	3a , 60
2	2b ($4\text{-FC}_6\text{H}_4/\text{C}_6\text{H}_5/\text{Me}/\text{Me}$)	3b , 55
3	2c ($4\text{-ClC}_6\text{H}_4/\text{C}_6\text{H}_5/\text{Me}/\text{Me}$)	3c , 65
4	2d ($4\text{-MeC}_6\text{H}_4/\text{C}_6\text{H}_5/\text{Me}/\text{Me}$)	3d , 70
5	2e ($\text{C}_6\text{H}_5/4\text{-ClC}_6\text{H}_4/\text{Me}/\text{Me}$)	3e , 63
6	2f ($\text{C}_6\text{H}_5/4\text{-BrC}_6\text{H}_4/\text{Me}/\text{Me}$)	3f , 65
7	2g ($\text{C}_6\text{H}_5/4\text{-MeC}_6\text{H}_4/\text{Me}/\text{Me}$)	3g , 68
8	2h ($\text{C}_6\text{H}_5/\text{C}_6\text{H}_5/\text{Me}/\text{H}$)	3h , 63
9	2i ($\text{C}_6\text{H}_5/\text{C}_6\text{H}_5/\text{H}/\text{H}$)	3i , 60
10	2j (Cyclopentyl/Cyclopentyl/Cyclopentyl/Cyclopentyl)	3j , 66
11	2k (Cyclohexyl/Cyclohexyl/Cyclohexyl/Cyclohexyl)	3k , 65

^a All reactions were carried out using **2** (0.2 mmol) in the presence of $\text{BF}_3 \cdot \text{OEt}_2$ (10 mol %) and DCE (5.0 mL) at 40 °C for 5 h. ^b Isolated yields.


presence of the Lewis acid $\text{Cu}(\text{OTf})_2$ in DCE at 50 °C. The optimization of the reaction conditions for this process is summarized in the Supporting Information in the presence of various Lewis acids, such as $\text{Sc}(\text{OTf})_3$, $\text{Sn}(\text{OTf})_2$, $\text{Eu}(\text{OTf})_3$, and $\text{Zr}(\text{OTf})_4$. It was found that $\text{Cu}(\text{OTf})_2$ was the best catalyst and that 1,2-dichloroethane (DCE) was the

Scheme 2. Plausible Mechanism for the Formation of Naphthalenes **3**



solvent of choice, and **4a** was produced in 98% yield at 50 °C (Scheme 3) (see the Supporting Information for details).

Scheme 3. Optimal Conditions for the Formation of Indene **4a**



It should be emphasized here that, using Brønsted acid $\text{CF}_3\text{-CO}_2\text{H}$ as a catalyst, both indene and naphthalene products can be obtained as a mixture. Compound **4a** was unambiguously determined by an X-ray diffraction (Figure 2).⁷ The




Figure 2. ORTEP drawing of compound **4a**.


generality of this $\text{Cu}(\text{OTf})_2$ -catalyzed rearrangement reaction is satisfactory, and the reaction can tolerate various substituents, whether they are electron-rich or -poor aryl groups (R^1 and R^2) or methyl groups or hydrogen atoms (R^3 and R^4). All reactions proceeded smoothly to give the corresponding indene derivative **4** in good to high yields under the optimal conditions (Table 3).

A plausible mechanism for the formation of indenenes **4** is illustrated in Scheme 4. The coordination of **2** to the Lewis acid initially gives zwitterionic intermediate **F**, which results in the formation of cyclopropyl ring-opened zwitterionic intermediate **G** via σ -bond *a*-cleavage. Intermediate **G** also can be drawn as its delocalized representations as intermediate **H** or intermediate **H'**.^{6,8} Subsequent intramolecular Friedel–Crafts reaction of intermediate **H'** with aromatic group R^1 affords intermediate **I**. Elimination of Lewis acid and deprotonation from intermediate **I** furnish **4**.

(7) The crystal data of **4a** have been deposited in the CCDC with number 615026. Empirical formula: $\text{C}_{25}\text{H}_{22}$. Formula weight: 322.43. Crystal color, habit: colorless, prismatic. Crystal system: Triclinic. Lattice type: primitive. Lattice parameters: $a = 8.1168(19)$ Å, $b = 10.068(2)$ Å, $c = 11.493(16)$ Å, $\alpha = 93.001(4)^\circ$, $\beta = 98.563(4)^\circ$, $\gamma = 98.080(4)^\circ$, $V = 917.0(4)$ Å³. Space group: $P1$. $Z = 2$; $D_c = 1.168$ g/cm³; $F_{000} = 344$. Diffractometer: Rigaku AFC7R. Residuals: R , R_w : 0.0655, 0.1722.

(8) For the mechanism of the 1,3-proton shift, see: Carey, F. A.; Sundberg, R. J. *Advanced Organic Chemistry*, 3rd ed.; Plenum Press: New York, 1990; p 609.

Table 3. Cu(OTf)₂-Catalyzed Rearrangement of Vinylcyclopropenes **2** for the Formation of Indenes **4**




entry ^a	2 (R ¹ /R ² /R ³ /R ⁴)	yield (%) ^b
1	2a (C ₆ H ₅ /C ₆ H ₅ /Me/Me)	4a , 98
2	2b (4-FC ₆ H ₄ /C ₆ H ₅ /Me/Me)	4b , 68
3	2c (4-ClC ₆ H ₄ /C ₆ H ₅ /Me/Me)	4c , 78
4	2d (4-MeC ₆ H ₄ /C ₆ H ₅ /Me/Me)	4d , 94
5	2e (C ₆ H ₅ /4-ClC ₆ H ₄ /Me/Me)	4e , 87
6	2f (C ₆ H ₅ /4-BrC ₆ H ₄ /Me/Me)	4f , 80
7	2g (C ₆ H ₅ /4-MeC ₆ H ₄ /Me/Me)	4g , 98
8	2h (C ₆ H ₅ /C ₆ H ₅ /Me/H)	4h , 84
9	2j (C ₆ H ₅ /C ₆ H ₅ /C ₆ H ₅ /C ₆ H ₅)	4i , 78
10	2k (C ₆ H ₅ /C ₆ H ₅ /C ₆ H ₅ /C ₆ H ₅)	4j , 75

^a All reactions were carried out using **2** (0.2 mmol) in the presence of Cu(OTf)₂ (10 mol %) and DCE (5.0 mL) at 50 °C for 5 h. ^b Isolated yields.

At the present stage, we believe that the coordination of sterically bulky LA to the cyclopropene moiety of **2** exclusively produces intermediate **F** due to the steric repulsion of LA and the aromatic R² group. However, the coordination of a proton to the cyclopropene moiety of **2** gives a more stabilized intermediate **B**. The different coordination place in the cyclopropene moiety of **2** causes the different reaction products.

Scheme 4. Plausible Mechanism for the Formation of Indenes **4**



In conclusion, we have discovered efficient and mild Lewis acid catalyzed rearrangement reactions of vinylcyclopropenes **2**. In these reactions, the choice of Lewis acid catalyst is important and naphthalenes **3** or indenes **4** can be obtained in good to high yields. Stronger Lewis acid catalysts (or Brønsted acid in this case) and somewhat weak Lewis acid catalysts will result in the different σ -bond cleavage of **2**, which directly affects the subsequent Friedel–Crafts reaction and determines the product structure. Efforts are underway in the laboratory to further elucidate the mechanistic details of these reactions and to understand their scope and limitations.

Acknowledgment. Financial support from the Shanghai Municipal Committee of Science and Technology (04JC14083, 06XD14005), the National Natural Science Foundation of China (203900502, 20472096, and 20672127), and the Cheung Kong Scholar Program is greatly appreciated.

Supporting Information Available: Spectroscopic data of all the new compounds, detailed experimental procedures, and X-ray data for compounds **2b** and **4a**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL0626746