


Journal Pre-proof

Sequential catalytic carbonylation reactions for sustainable synthesis of biologically relevant entities

Liliana Damas, Fábio M.S. Rodrigues, Andreia C.S. Gonzalez, Rui M.B. Carrilho, Marta Pineiro, Mariette M. Pereira



PII: S0022-328X(20)30319-3

DOI: <https://doi.org/10.1016/j.jorganchem.2020.121417>

Reference: JOM 121417

To appear in: *Journal of Organometallic Chemistry*

Received Date: 1 June 2020

Accepted Date: 23 June 2020

Please cite this article as: L. Damas, F. M. S. Rodrigues, A. C. S. Gonzalez, R. M. B. Carrilho, M. Pineiro, M. M. Pereira, Sequential catalytic carbonylation reactions for sustainable synthesis of biologically relevant entities, *Journal of Organometallic Chemistry* (2020), doi: <https://doi.org/10.1016/j.jorganchem.2020.121417>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier B.V.

Journal Pre-proof

Sequential Catalytic Carbonylation Reactions for Sustainable Synthesis of Biologically Relevant Entities

Liliana Damas, Fábio M.S. Rodrigues, Andreia C.S. Gonzalez, Rui M.B. Carrilho*, Marta Pineiro and Mariette M. Pereira*,[‡]

University of Coimbra, Coimbra Chemistry Centre, Department of Chemistry
Rua Larga, 3004-535 Coimbra, Portugal

Abstract

The sustainable synthesis of highly functionalised formylcarboxamide compounds with biological relevance is reported through a sequential aminocarbonylation/hydroformylation approach. The optimisation of palladium-catalysed aminocarbonylation of iodoaromatic substrates, using allylamine as nucleophile was first performed, with molybdenum hexacarbonyl as alternative CO source *versus* gaseous carbon monoxide. The combination of microwave irradiation with molybdenum hexacarbonyl allowed to selectively prepare a set of *N*-heterocyclic-based allylcarboxamides. Subsequent rhodium-catalysed hydroformylation of the allylcarboxamide intermediates led to the preparation of new pyridine, pyrazoline and chalcone derivatives containing both carboxamide and formyl moieties.

Keywords: sequential reactions; aminocarbonylation; hydroformylation; microwave synthesis; *N*-heterocycles; chalcones.

* corresponding authors: mmpereira@qui.uc.pt (M. M. Pereira), ruy.carrilho@uc.pt (R. M. B. Carrilho) Address: University of Coimbra, Department of Chemistry, Rua Larga, 3004-535 Coimbra, Portugal, Telephone: +351 239 854 474.

[‡]Dedicated to Professor Laszló Kollár, on the occasion of his 65th birthday.

1. Introduction

Carbonyl compounds, such as carboxamides and aldehydes, are among the most relevant synthons for preparation of highly functionalised molecules with potential applications in medicinal chemistry [1-6]. Particularly, those containing *N*-heterocycles [7-14] and chalcone [15-17] cores are frequently present in important pharmacological entities, such as antibiotics, antibacterial, anti-hypertensive, anti-inflammatory and anticancer drugs (Figure 1).





Figure 1. Examples of *N*-heterocyclic and chalcone-based carbonyl compounds with pharmacological activity.

Therefore, there has been an increasing interest in the search for efficient and sustainable chemical processes for preparing such molecules. Among them, transition metal catalysed hydroformylation [18-25] and aminocarbonylation [26-34] reactions are two paradigmatic examples of efficient one-pot and versatile synthetic approaches to get access to carbonyl compounds, such as aldehydes and carboxylic acid derivatives. Regarding these topics, it should be highlighted the relevant contribution of L. Kollár, with more than 300 published papers [35]. Furthermore, the development of sequential carbonylation reactions have led to significant reduction in costs, purification steps and waste formation. Such sequential reactions present higher atom economy, allowing to improve the general processes efficiency and sustainability [36-44]. It should be also noted that the global demands for implementation of more sustainable synthetic processes have boosted the search for alternative CO sources [45-49], including metal-carbonyl compounds [50-53] and the development of energetically favourable processes, such as microwave-assisted carbonylation reactions [54-58].

Along the last years, we have been focused on the development of efficient transition-metal catalysed carbonylation-based sequential processes applied to biologically relevant substrates for the preparation value-added products with biological activity and/or industrial interest, including hydroformylation/arylation [59], hydroformylation/isomerisation [60],

Journal Pre-proof
hydroformylation/reductive amination [61,62], hydroformylation/Strecker [62], hydroformylation/hydrogenation [63] and aminocarbonylation/cyclisation [64].

As part of our continuing research in this field, herein we report an original sequential process for preparation of biologically relevant formylcarboxamide compounds comprising N-heterocycle and chalcone scaffolds, through microwave-assisted palladium-catalysed aminocarbonylation of iodoaromatic substrates with allylamine, followed by rhodium-catalysed hydroformylation of the allylcarboxamide intermediates, to obtain carbonylated products containing both formyl and carboxamide motifs (**Scheme 1**).




Scheme 1

2. Results and discussion

The palladium-catalysed aminocarbonylation step was first optimised, using iodobenzene (**1**) as a model substrate, allylamine as nucleophile, palladium (II) acetate as catalytic precursor, DBU as base and dioxane as solvent. The reactions were carried out in the presence of carbon monoxide or molybdenum hexacarbonyl as alternative CO carbonyl source, at 125 °C, using either conventional heating or microwave irradiation. Conversion and selectivity were determined by GC-MS analysis of the crude mixtures and the results are presented in Table 1.

Table 1. Aminocarbonylation of iodobenzene (**1**) using allylamine as nucleophile.^a



Entry	Carbonyl source	Heating	Time (min)	Conversion (%)	Selectivity (%)	TOF (h ⁻¹)
1	CO (3 bar)	Conv.	60	79	69	32
2	CO (3 bar)	MW (150W)	5	76	62	365
3	Mo(CO) ₆	Conv.	60	99	82	40
4	Mo(CO) ₆	MW (150 W)	5	99	76	475













a) Reaction conditions: iodobenzene (0.4 mmol), allylamine (0.6 mmol), Pd(OAc)₂ (0.01 mmol), Mo(CO)₆ (0.4 mmol), DBU (0.2 mL), dioxane (3 mL), 125 °C.


Using a CO pressure of 3 bar and conventional heating, a conversion of 79% was obtained in 60 min (TOF = 32 h⁻¹), with 69% selectivity for the carboxamide product (**2**) (Table 1, entry 1). Under microwave irradiation (150 W), a similar conversion was obtained in just 5 min, leading to a substantial increase in TOF (365 h⁻¹), although a slightly slower selectivity (62%) was observed (Table 1, entry 2). Alternatively, the use of Mo(CO)₆ complex as alternative carbonyl source, under conventional heating, provided full conversion in 60 min (TOF = 40 h⁻¹), with 82% selectivity for the carboxamide (**2**) (Table 1, entry 3). In addition, under microwave irradiation, the reaction proceeded with full conversion in 5 min (TOF = 475 h⁻¹) and with 76% selectivity for the carboxamide product (**2**) (Table 1, entry 4). In sum, microwave irradiation allowed to significantly reduce the required reaction time from 60 min to 5 min with enhancement of the activity (TOF = 40 and 475 h⁻¹ respectively), maintaining the reaction selectivity.

With optimised conditions, microwave-assisted Pd-catalysed aminocarbonylation reactions, using $\text{Mo}(\text{CO})_6$ as carbonyl source, were applied to different iodoaromatic substrates with biologically relevant scaffolds, including 1-bromo-4-iodo benzene (**3**), 4-iodoanisole (**4**), iodopyridine (**5**), 7-iodoindole (**6**), 5-iodoindole (**7**) and iodopyrazoline (**8**) and iodochalcone (**9**) [68]. Conversion and yields were determined by GC-MS analysis of the crude mixtures and the results are presented in Table 2.

Table 2. Microwave-assisted Pd-catalysed aminocarbonylation using allylamine as nucleophile.^a




Entry	Substrate	Product (yield, %) ^b
1		 2 (76)
2		 10 (71)
3		 11 (65)
4		 12 (76)
5		 13 (87)
6		 14 (72)
7 ^d		 15 (70)



- a) Reaction conditions: substrate (0.4 mmol), allylamine (0.6 mmol), Pd(OAc)₂ (0.01 mmol), Mo(CO)₆ (0.4 mmol), DBU (0.2 mL), dioxane (3 mL), 125 °C, 5 min, 150 W.
 b) Yield determined by GC-MS analysis of crude mixtures.
 c) 1 h.


The microwave-assisted aminocarbonylation reactions proceeded with *ca.* 99% conversion in 5 min and allylcarboxamide derivatives were obtained as major products in all cases, in yields ranging from 65% (for product **11**) to 87% (for product **13**). In addition, the aminocarbonylation of substrates **3** and **5** led to predominant formation mono-carbonylated products **10** and **12**, respectively (Table 2, entries 2 and 4), which indicate that oxidative addition to the palladium complex occurs mainly through the most reactive C–I bond, while C–Br and C–Cl bonds remain unchanged under these conditions. Moreover, in the case of iodopyrazoline (**8**), the reaction required 1 h, under microwave irradiation, to obtain the corresponding allylcarboxamide product (**15**), obtained in 70% yield. In all cases, GC-MS indicated the formation of ketocarboxamides as minor products (up to *ca.* 10 %), corroborated by the ¹³C NMR spectra which presented typical signals around 190 ppm. After work up and purification by column chromatography in silica gel, using EtOAc/*n*-hexane or EtOAc/CH₂Cl₂ mixtures as eluents, the carboxamide products **2** and **10-16** have been isolated and fully characterised. In sum, microwave-assisted palladium-catalysed aminocarbonylation of iodoaromatic substrates, using allylamine as a nucleophile, allowed optimising the first step of the sequential process and provided the access to a set of allylcarboxamide products.

In order to minimise product losses by work-up and purification procedures, a sequential catalytic aminocarbonylation/hydroformylation methodology was then carried out, without work-up or purification of the allylcarboxamide intermediates. Thus, using iodobenzene (**1**) as a model substrate, the crude mixture obtained in the microwave-assisted aminocarbonylation step was directly subjected to hydroformylation conditions (8 bar H₂/CO) with a rhodium/triphenylphosphine catalyst, at 50 °C. However, in a first attempt, no conversion was observed after 6 h. This was attributed to the formation of a sterically hindered rhodium/DBU complex of type **17** [65,66], which leads to formation of catalytically inactive species.



To overcome this issue, the crude mixture, resulting from the aminocarbonylation step, was passed through a silica-pad previous to hydroformylation. Following this procedure, full substrate conversion was obtained in 5 h, with 72% chemoselectivity for oxo-products and 44% regioselectivity for the branched aldehyde (**18**) (Table 3, entry 1). Then, aiming the preparation of biologically relevant formylcarboxamide molecules, the scope of this sequential aminocarbonylation/hydroformylation methodology was expanded to iodopyridine (**5**), iodopyrazoline (**8**) and iodochalcone (**9**) substrates.

Table 3. Synthesis of formylcarboxamides by sequential aminocarbonylation/hydroformylation methodology.^a




Entry	Substrate	Chemo (%) ^b	<i>iso</i> -Regio (%) ^c	Product (isolated yield, %)
1		72	44	 18 (21)
2		73	51	 19 (20)
3 ^d		56	46	 20 (19)
4 ^d		64	36	 21 (22)

- a) Reactions conditions: *aminocarbonylation step* substrate (0.4 mmol), allylamine (0.6 mmol), Pd(OAc)₂ (0.01 mmol), Mo(CO)₆ (0.4 mmol), DBU (0.2 mL), dioxane (3 mL), 125 °C, 150 W, 5 min.; *hydroformylation step*: Rh(acac)(CO)₂/PPh₃ = 1:5; Rh/substrate = 1:100; P = 8 bar H₂/CO; 50 °C, 5 h. Conversion was ca 99% in all cases.
- b) Chemoselectivity for oxo-products (determined by ¹H NMR).
- c) Regioselectivity for the branched aldehyde (*iso*) (determined by ¹H NMR).
- d) 12 h

In a typical sequential aminocarbonylation/hydroformylation experiment, the iodo-substrate, allylamine, Pd(OAc)₂, Mo(CO)₆ and DBU were dissolved in dioxane and placed inside a microwave vial. The mixture was then subjected to microwave irradiation (P = 150 W), the reaction proceeding at 125 °C for 5 min. After cooling down to room temperature, the obtained mixture was filtered through a silica pad and the residue was introduced *via* cannula into an autoclave containing the precursor, Rh(CO)₂acac and triphenylphosphine (PPh₃). The autoclave was then pressurised with 8 bar H₂/CO and kept at 50 °C for 5 h (or 12 h). At the end, the solvent was evaporated and the crude mixture was analysed by ¹H and ¹³C NMR spectroscopy.

The sequential aminocarbonylation/hydroformylation processes proceeded with complete conversions and the chemoselectivity for oxo-products was in the range 56-73%. The regioselectivity for the branched aldehydes varied from 36 to 51% (Table 3, entries 2-4). Upon purification by column chromatography in silica gel, the branched formylcarboxamide products **18-21** were isolated in yields ranging from 19 to 22%. These low yields can be explained by the significant formation of linear aldehydes and the respective cyclisation products (not isolated), resultant from intramolecular condensation between the carboxamide and formyl moiety of the linear aldehydes, which leads to the formation of a stable 5 membered ring (**Scheme 2**) [67], evidenced by ¹H NMR and GC-MS.



Scheme 2

It is worth mentioning that, in the case of chalcone-derived substrate (**9**), the hydroformylation of the allyl moiety occurred concomitantly with reduction of the conjugated internal C=C double bond, which resulted in the formation of formylcarboxamide **21**, obtained as major product (Table 3, entry 4).

4. Conclusions

We have developed an innovative catalytic aminocarbonylation/hydroformylation sequential approach to prepare new formylcarboxamide molecules. The optimisation of the first step of the catalytic sequence was achieved through microwave-assisted palladium-catalysed aminocarbonylation of iodoaromatic substrates, using allylamine as nucleophile and $\text{Mo}(\text{CO})_6$ as alternative carbonyl source, which provided an efficient strategy for the synthesis of allylcarboxamide derivatives. The subsequent Rh/triphenylphosphine-catalysed hydroformylation of the allylcarboxamide intermediates allowed the synthesis of a set of new *N*-heterocyclic and chalcone-based formylcarboxamides, including 6-chloro-*N*-(2-methyl-3-oxopropyl)nicotinamide, 4-(1,3-diphenyl-4,5-dihydro-1*H*-pyrazol-5-yl)-*N*-(2-methyl-3-oxopropyl)benzamide and (*E*)-*N*-(2-methyl-3-oxopropyl)-4-(3-oxo-3-phenylprop-1-en-1-yl)benzamide. In sum, this transition-metal catalysed aminocarbonylation/hydroformylation sequential process opens new perspectives regarding the preparation of biologically relevant molecules containing both carboxamide and formyl motifs.

5. Experimental section

5.1. General

Manipulation of all moisture sensitive reagents was carried out under nitrogen atmosphere by using Schlenk techniques. Glassware was dried in an oven at 200 °C and cooled under a nitrogen atmosphere. Palladium (II) acetate, triphenylphosphine, $\text{Mo}(\text{CO})_6$, iodobenzene, 1-bromo-4-iodobenzene, 4-methoxyiodobenzene, 2-chloro-5-iodopyridine, 7-iodoindole, 5-iodoindole, DBU and dioxane were purchased from Merck and used without further purification. The iodopyrazoline **8** and iodochalcone **9** substrates were synthesised through a mechanochemical synthetic methodology, recently developed by some of us [68].

Microwave-assisted experiments were performed in thick-walled glass vials under closed-vessel conditions, using a CEM Discover[®] SP Focused Microwave[™] Synthesis System. NMR spectra were recorded on a Bruker Avance 400 spectrometer, operating at 400.13 MHz for ¹H NMR and 100.62 MHz for ¹³C NMR. Chemical shifts (δ) are reported in ppm relatively to CDCl_3 (7.260 and 77.16 ppm for ¹H and ¹³C, respectively) or tetramethylsilane (TMS). High-resolution mass spectrometry analysis was carried out on a Bruker Microtof apparatus, equipped with selective ESI detector. Samples of the reactions were analysed by: a) gas chromatography carried out on Agilent-7820A GC System equipped with a non-polar capillary HP-5 column (5% diphenyl and 95% dimethylpolysiloxane), with 30 m length and 0.32 mm inside diameter and using nitrogen as carrier gas, and equipped with an FID detector; or b) gas chromatography

Journal Pre-proof

coupled with mass spectrometry (GC-MS) using an Agilent 7820A GC System, equipped with a HP-5 MS column, coupled to an Agilent 5975 MSD System Technologies spectrometer, using EI (70 eV) and helium as carrier gas. Carboxamide derivatives were analysed by gas chromatography using the following methods: a) GC-FID method: injector temp 250 °C, oven: starting temp 50 °C (hold-time 2 min), heating rate 30 °C min⁻¹, final temp 280 °C (hold-time 9 min); detector temp 280 °C, carrier gas: nitrogen (rate: 1.6 mL min⁻¹); b) GC-MS method: injector temp 250 °C, oven: starting temperature 50 °C (hold-time 1 min), heating rate 15 °C min⁻¹, final temp 280 °C (hold-time 15 min); detector temp 280 °C, carrier gas: helium (rate: 1 mL min⁻¹).

5.2. Microwave-assisted palladium-catalysed aminocarbonylation

The iodoaryl substrate (0.4 mmol), allylamine (45 µL, 0.6 mmol), Pd(OAc)₂ (2.2 mg, 0.01 mmol), Mo(CO)₆ (105.6 mg, 0.4 mmol), DBU (180 µL, 1.2 mmol) were dissolved in dioxane (3 mL) and placed into a microwave vial. The mixture was subjected to microwave irradiation (Initial Power = 150 W) and the reaction was conducted at 125 °C for 5 min. The mixture was then cooled to room temperature, filtered through a small plug of celite and evaporated to dryness with the crude being analysed by GC-MS and/or GC-FID (methods described in section 5.1). The residue obtained was dissolved in dichloromethane (20 mL) and washed with water (3×20 mL). The organic phase was dried over anhydrous Na₂SO₄, filtered and evaporated to a solid material or to a waxy residue. The reaction products were isolated and purified by column chromatography (Silicagel 60 (Merck), 0.063-0.200 mm), using EtOAc/*n*-hexane or EtOAc/CH₂Cl₂ mixtures as the eluents (specified below for each compound).

5.3. General procedure of sequential aminocarbonylation/hydroformylation

For the sequential aminocarbonylation/hydroformylation process the iodoaromatic substrate (0.4 mmol), allylamine (45 µL, 0.6 mmol, 1.5 equiv.), Pd(OAc)₂ (2.2 mg, 0.01 mmol), Mo(CO)₆ (105.6 mg, 0.4 mmol), DBU (180 µL, 1.2 mmol) and dioxane (3 mL) were placed into a microwave vial. The mixture was subjected to microwave irradiation (Initial Power = 150 W) and the reaction was conducted at 125 °C for 5 min. After cooled down to room temperature, the aminocarbonylation crude mixture was filtered through a small plug of silica gel and introduced *via* cannula into the autoclave. The rhodium precursor Rh(acac)(CO)₂ (4×10⁻³ mmol, 1.03 mg) and triphenylphosphine (0.02 mmol, 5.24 mg) were dissolved in dioxane (3 mL), under argon atmosphere, and introduced *via* cannula into the autoclave. After purging the system with three cycles of vacuum/*syngas* the reactor was pressurised with 8 bar of H₂/CO (1:1) and kept at 50 °C.

After 5 h, the reactor was cooled to room temperature and slowly depressurised. The reaction mixture was evaporated to dryness analysed by ^1H NMR spectroscopy. The reaction products were isolated and purified by column chromatography (Silicagel 60 (Merck), 0.063-0.200 mm), using EtOAc/*n*-hexane or EtOAc/ CH_2Cl_2 mixtures as the eluents (specified below for each compound).

5.4.1 Characterisation of products

The spectroscopic data of **2**, **10** and **11** are in agreement with that previously reported [69,70,71].

N-allyl-6-chloronicotinamide (**12**)

R_f ($\text{CH}_2\text{Cl}_2/\text{EtOAc}$ 1:1) = 0.74. Beige solid, Yield 23%, 18 mg (0.09 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 8.75 (d, J = 7.6 Hz, 1H), 8.10 (dd, J = 8.3, 2.5 Hz, 1H), 7.43 (d, J = 8.3 Hz, 1H), 6.18 (s, 1H), 5.97 – 5.90 (m, 1H), 5.28 (dd, J = 17.1, 1.2 Hz, 1H), 5.23 (dd, J = 10.2, 0.9 Hz, 1H), 4.12 – 4.08 (m, 2H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 164.5; 154.5; 148.0; 138.1; 133.6; 129.2; 124.6; 117.6; 42.8. HRMS (ESI) (m/z): found 197.0477 [$\text{M}+\text{H}$]⁺ (calcd. 197.0476).

N-allyl-1*H*-indole-7-carboxamide (**13**)

R_f ($\text{EtOAc}/n\text{-hex}$ 1:4) = 0.37. White solid. Yield 20%, 16 mg (0.08 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 10.32 (br s, 1H), 7.81 (d, J = 7.8 Hz, 1H), 7.38 (d, J = 7.4 Hz, 1H), 7.33 – 7.32 (m, 1H), 7.12 (t, J = 7.7 Hz, 1H), 6.58 – 6.57 (m, 1H), 6.47 (s, 1H), 6.01 – 5.94 (m, 1H), 5.30 (dd, J = 17.2, 1.4 Hz, 1H), 5.21 (dd, J = 10.2, 1.3 Hz, 1H), 4.16 – 4.13 (m, 2H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 167.8; 135.6; 134.4; 129.7; 125.8; 125.0; 118.9; 118.8; 116.8; 115.9; 102.1; 42.2. HRMS (ESI) (m/z): found 201.1023 [$\text{M}+\text{H}$]⁺ (calcd. 201.1022).

N-allyl-1*H*-indole-5-carboxamide (**14**)

R_f ($\text{CH}_2\text{Cl}_2/\text{EtOAc}$ 2:1) = 0.51. White solid. Yield 23%, 18 mg (0.09 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 8.53 (br s, 1H), 8.11 (s, 1H), 7.66 (dd, J = 8.5, 1.5 Hz, 1H), 7.42 (d, J = 8.5 Hz, 1H), 7.28 – 7.26 (m, 1H), 6.62 (s, 1H), 6.23 (br s, 1H), 6.01 – 5.94 (m, 1H), 5.29 (dd, J = 17.1, 1.5 Hz, 1H), 5.19 (dd, J = 10.2, 1.3 Hz, 1H), 4.14 – 4.12 (m, 2H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 168.6; 137.7; 134.7; 127.7; 126.6; 125.7; 121.2; 120.3; 116.6; 111.2; 103.8; 42.6. HRMS (ESI) (m/z): found 201.1024 [$\text{M}+\text{H}$]⁺ (calcd. 201.1022).

N-allyl-4-(1,3-diphenyl-4,5-dihydro-1*H*-pyrazol-5-yl)benzamide (**15**)

R_f ($\text{EtOAc}/n\text{-hex}$ 1:1) = 0.66. Yellow solid. Yield 21%, 32 mg (0.08 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 7.73 (t, J = 7.3 Hz, 4H), 7.31 – 7.25 (m, 4H), 7.34 (d, J = 7.2 Hz, 2H), 7.20 –

Journal Pre-proof
7.16 (m, 3H), 7.04 (d, $J = 7.8$ Hz, 2H), 6.80 (t, $J = 7.3$ Hz, 1H), 6.19 (br s, 1H), 5.95 – 5.88 (m, 1H), 5.30 (dd, $J = 12.4, 7.2$ Hz, 1H), 5.24 (dd, $J = 17.2, 1.4$ Hz, 1H), 5.17 (dd, $J = 10.2, 1.2$ Hz, 1H), 4.07 – 4.04 (m, 2H), 3.86 (dd, $J = 17.0, 12.4$ Hz, 1H), 3.11 (dd, $J = 17.1, 7.2$ Hz, 1H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 167.0; 146.9; 146.3; 144.7; 134.2; 134.0; 132.6; 129.1; 128.9; 128.7; 128.0; 126.3; 125.9; 119.5; 116.9; 113.5; 64.3; 43.5; 42.5. HRMS (ESI) (m/z): found 381.1835 [M^+] (calcd. 381.1841).

(E)-*N*-allyl-4-(3-oxo-3-phenylprop-1-en-1-yl)benzamide (**16**)

Rf (EtOAc/*n*-hex 1:1) = 0.56. Yellow solid. Yield 25%, 29 mg (0.1 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 8.04 – 8.02 (m, 2H), 7.85 – 7.79 (m, 3H), 7.70 (d, $J = 8.2$ Hz, 2H), 7.63 – 7.59 (m, 2H), 7.55 (d, $J = 10.8$ Hz, 1H), 7.51 (d, $J = 7.3$ Hz, 1H), 6.26 (s, 1H), 5.99 – 5.92 (m, 1H), 5.28 (dd, $J = 17.1, 1.4$ Hz, 1H), 5.21 (dd, $J = 10.2, 1.2$ Hz, 1H), 4.13 – 4.10 (m, 2H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 190.4; 166.6; 143.4; 138.0; 137.9; 136.0; 134.1; 133.2; 128.8; 128.7; 128.6; 127.7; 123.8; 117.0; 42.7. HRMS (ESI) (m/z): found 292.13302 [$\text{M}+\text{H}$] $^+$ (calcd. 292.1332).

N-(2-methyl-3-oxopropyl)benzamide (**18**)

Rf ($\text{CH}_2\text{Cl}_2/\text{EtOAc}$ 7:3) = 0.54. Yellow oil. Yield 21%, 16 mg (0.08 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 9.71 (s, 1H), 7.74 – 7.72 (m, 2H), 7.50 – 7.46 (m, 1H), 7.40 (dd, $J = 10.2, 4.6$ Hz, 2H), 6.73 (br s, 1H), 3.75 – 3.69 (m, 1H), 3.57 – 3.50 (m, 1H), 2.82 – 2.74 (m, 1H), 1.21 (d, $J = 7.5$ Hz, 3H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 204.5; 167.7; 134.3; 131.7; 128.7; 127.0; 46.9; 39.9; 11.6. HRMS (ESI) (m/z): found 214.08399 [$\text{M}+\text{Na}$] $^+$ (calcd. 214.0839).

6-chloro-*N*-(2-methyl-3-oxopropyl)nicotinamide (**19**)

Rf ($\text{CH}_2\text{Cl}_2/\text{EtOAc}$ 1:2) = 0.27. Beige solid. Yield 20%, 18 mg (0.08 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 9.70 (s, 1H), 8.73 (d, $J = 2.2$ Hz, 1H), 8.04 (dd, $J = 8.3, 2.5$ Hz, 1H), 7.39 (d, $J = 8.3$ Hz, 1H), 6.92 (s, 1H), 3.78 – 3.70 (m, 1H), 3.56 – 3.48 (m, 1H), 2.86 – 2.78 (m, 1H), 1.23 (d, $J = 7.5$ Hz, 3H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 204.6; 164.8; 154.4; 148.2; 139.2; 129.3; 124.8; 46.7; 39.0; 11.7. HRMS (ESI) (m/z): found 227.0581 [$\text{M}+\text{H}$] $^+$ (calcd. 227.0582).

4-(1,3-diphenyl-4,5-dihydro-1*H*-pyrazol-5-yl)-*N*-(2-methyl-3-oxopropyl)benzamide (**20**)

Rf (EtOAc/*n*-hex 2:1) = 0.44. Yellow solid. Yield 19%, 31 mg (0.08 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 9.69 (s, 1H), 7.73 – 7.66 (m, 4H), 7.39 – 7.34 (m, 5H), 7.19 – 7.15 (m, 2H), 7.03 (d, $J = 7.8$ Hz, 2H), 6.79 (t, $J = 7.3$ Hz, 1H), 6.60 (t, $J = 5.9$ Hz, 1H), 5.30 (dd, $J = 12.5, 7.0$ Hz, 1H), 3.86 (dd, $J = 17.1, 12.4$ Hz, 1H), 3.74 – 3.67 (m, 1H), 3.53 – 3.50 (m, 1H), 3.10 (dd, $J =$

Journal Pre-proof
17.1, 7.3 Hz, 1H), 2.80 – 2.74 (m, 1H), 1.21 (d, $J = 7.5$ Hz, 3H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 204.5; 167.3; 152.3; 146.9; 146.4; 133.8; 133.8; 129.1; 128.9; 128.7; 128.0; 126.4; 125.9; 119.5; 113.5; 64.3; 46.9; 43.5; 39.9; 11.6. HRMS (ESI) (m/z): found 412.2020 [$\text{M}+\text{H}$] $^+$ (calcd. 412.2020).

(*E*)-*N*-(2-methyl-3-oxopropyl)-4-(3-oxo-3-phenylprop-1-en-1-yl)benzamide (**21**)

Rf ($\text{CH}_2\text{Cl}_2/\text{EtOAc}$ 1:1) = 0.6. Yellow solid. Yield 22%, 28 mg (0.09 mmol). ^1H RMN (400 MHz, CDCl_3 , ppm): δ 9.71 (s, 1H), 7.95 – 7.93 (m, 2H), 7.67 (d, $J = 8.2$ Hz, 2H), 7.56 (t, $J = 7.4$ Hz, 1H), 7.45 (t, $J = 7.6$ Hz, 2H), 7.30 (d, $J = 8.1$ Hz, 2H), 6.63 (s, 1H), 3.73 (ddd, $J = 13.7, 6.7, 4.3$ Hz, 1H), 3.53 (ddd, $J = 13.8, 8.0, 5.8$ Hz, 1H), 3.30 (t, $J = 7.5$ Hz, 2H), 3.11 (t, $J = 7.5$ Hz, 2H), 2.81 – 2.76 (m, 1H), 1.22 (d, $J = 7.5$ Hz, 3H). ^{13}C RMN (100 MHz, CDCl_3 , ppm): δ 204.5; 198.9; 167.5; 145.5; 136.8; 133.3; 132.2; 128.8; 128.8; 128.1; 127.3; 46.9; 40.0; 39.9; 30.0; 11.6. HRMS (ESI) (m/z): found: 324.1591 [$\text{M}+\text{H}$] $^+$ (calcd. 324.1594).

Acknowledgements

This work was supported by national funds from FCT – Fundação para a Ciência e a Tecnologia, I.P., through projects UID/QUI/00313/2019 to the Coimbra Chemistry Centre (CQC), PTDC/QUI-OUT/27996/2017 (DUALPI) and POCI-01-0145-FEDER-016387 (SunStorage). LD thanks MATIS - Materiais e Tecnologias Industriais Sustentáveis (CENTRO-010145-FEDER- 00014) and FMSR thanks FCT PhD CATSUS program (PD/BD/114340/2016) for funding.

References

- [1] T.N. Reddy, A. Beatriz, V.J. Rao, D.P. de Lima, Carbonyl Compounds' Journey to Amide Bond Formation, *Chem. Asian J.* 14 (2019) 344–388. <https://doi.org/10.1002/asia.201801560>.
- [2] C. De Risi, G.P. Pollini, V. Zanirato, Recent Developments in General Methodologies for the Synthesis of α -Ketoamides, *Chem. Rev.* 116 (2016) 3241–3305. <https://doi.org/10.1021/acs.chemrev.5b00443>.
- [3] L. Kollár (Ed.), *Modern Carbonylation Methods*, Wiley-VCH, Weinheim, 2008.
- [4] S. Zhao, N.P. Mankad, Metal-catalysed radical carbonylation reactions, *Catal. Sci. Technol.* 9 (2019) 3603-3613. <https://doi.org/10.1039/C9CY00938H>.
- [5] S.T. Gadge, B.M. Bhanage, Recent developments in palladium catalysed carbonylation reactions, *RSC Adv.*, 4 (2014) 10367–10389. <https://doi.org/10.1039/c3ra46273k>.

- [6] R. Franke, D. Selent, A. Börner, Applied Hydroformylation, *Chem. Rev.* 112 (2012) 5675–5732. <https://doi.org/10.1021/cr3001803>.
- [7] S. Perrone, L. Troisi, A. Salomone, Heterocycle Synthesis through Pd-Catalyzed Carbonylative Coupling, *Eur. J. Org. Chem.* (2019) 4626–4643. <https://doi.org/10.1002/ejoc.201900439>.
- [8] X.-F. Wu, H. Neumann, M. Beller, Synthesis of Heterocycles via Palladium-Catalyzed Carbonylations, *Chem. Rev.* 113 (2013) 1–35. <https://doi.org/10.1021/cr300100s>.
- [9] A. Kumari, R.K. Singh, Medicinal chemistry of indole derivatives: Current to future therapeutic prospectives, *Bioorg. Chem.* 89 (2019) 103021. <https://doi.org/10.1016/j.bioorg.2019.103021>.
- [10] P. Das, M.D. Delost, M.H. Qureshi, D.T. Smith, J.T. Njardarson, A Survey of the Structures of US FDA Approved Combination Drugs, *J. Med. Chem.* 62 (2019) 4265–4311. <https://doi.org/10.1021/acs.jmedchem.8b01610>.
- [11] P.R. Kamath, D. Sunil, A.A. Ajees, K.S.R. Pai, S. Das, Some new indole–coumarin hybrids; Synthesis, anticancer and Bcl-2 docking studies, *Bioorg. Chem.* 63 (2015) 101–109. <http://dx.doi.org/10.1016/j.bioorg.2015.10.001>.
- [12] F. Nawaz, O. Alam, A. Perwez, M.A. Rizvi, M.J. Naim, N. Siddiqui, F.H. Pottoo, M. Jha, 3'-(4-(Benzyloxy)phenyl)-1'-phenyl-5-(heteroaryl/aryl)-3,4-dihydro-1'H,2H-[3,4'-bipyrazole]-2-carboxamides as EGFR kinase inhibitors: Synthesis, anticancer evaluation, and molecular docking studies, *Arch. Pharm. Chem. Life Sci.* 353 (2020) 1900262. <https://doi.org/10.1002/ardp.201900262>.
- [13] Z. Zhang, Y.-L. Gu, Z.-Y. Wang, H.-N. Wang, Y. Zhao, X.-M. Chu, C.-Y. Zhang, M.-C. Yan, Synthesis and biological evaluation of novel indoleamide derivatives as antioxidative and antitumor agents, *J. Heterocyclic Chem.* 57 (2020) 1165–1172. <https://doi.org/10.1002/jhet.3853>.
- [14] R. Tamatam, S.S. Panga, Z. Grigory V, S. Gundala, B.R. Nemallapudi, K.P. Vasikarla, P. Adivireddy, P. Venkatapuram, Synthesis, Antimicrobial and Antioxidant Activities of Pyrimidinyl Benzothiazine Carboxamides, *ChemistrySelect* 4 (2019) 6813–6820. <https://doi.org/10.1002/slct.201900525>.
- [15] C. Thirumurugan, P. Vadivel, A. Lalitha, S. Lakshmanan, Synthesis, characterization of novel quinoline-2-carboxamide based chalcone derivatives and their molecular docking, photochemical studies, *Synth. Commun.* 50 (2020) 831–839. <https://doi.org/10.1080/00397911.2020.1720737>.

- [16] S. Nisa, M. Yusuf, Synthetic and antimicrobial studies of N-substituted pyrazoline-based new bisheterocycles, *J. Heterocyclic Chem.* 57 (2020) 2024–2036. <https://doi.org/10.1002/jhet.3910>.
- [17] M.S. Abdelbaset, M. Abdel-Aziz, M. Ramadan, M.H. Abdelrahman, S.N.A. Bukhari, T. F.S. Ali, G.E.-D. A. Abuo-Rahma, Discovery of novel thienoquinoline-2-carboxamide chalcone derivatives as antiproliferative EGFR tyrosine kinase inhibitors, *Bioorg. Med. Chem.* 27 (2019) 1076–1086. <https://doi.org/10.1016/j.bmc.2019.02.012>.
- [18] A.C. Brezny, C.R. Landis, Recent Developments in the Scope, Practicality, and Mechanistic Understanding of Enantioselective Hydroformylation, *Acc. Chem. Res.* 51 (2018) 2344–2354. <https://doi.org/10.1021/acs.accounts.8b00335>.
- [19] S.S. Nurttala, P.R. Linnebank, T. Krachko, J.N. H. Reek, Supramolecular Approaches To Control Activity and Selectivity in Hydroformylation Catalysis, *ACS Catal.* 8 (2018) 3469–348. <https://doi.org/10.1021/acscatal.8b00288>.
- [20] A. Cunillera, C. Blanco, A. Gual, J.M. Marinkovic, E.J. Garcia-Suarez, A. Riisager, C. Claver, A. Ruiz, C. Godard, Highly Efficient Rh-catalysts Immobilised by π - π Stacking for the Asymmetric Hydroformylation of Norbornene under Continuous Flow Conditions, *ChemCatChem* 11 (2019) 2195–2205. <https://doi.org/10.1002/cctc.201900211>.
- [21] A. Gual, C. Godard, S. Castellón, C. Claver, Highlights of the Rh-catalysed asymmetric hydroformylation of alkenes using phosphorus donor ligands, *Tetrahedron Asymmetry* 21 (2010) 1135–1146. <https://doi.org/10.1016/j.tetasy.2010.05.037>.
- [22] F.G. Delolo, E.N. dos Santos, E.V. Gusevskaya, Anisole: a further step to sustainable hydroformylation, *Green Chem.*, 21 (2019) 1091–1098. <https://doi.org/10.1039/c8gc03750g>.
- [23] A.C. Faria, M.P. de Oliveira, A.C. Monteiro, R.L.V. Mota, K.C.B. Oliveira, E.N. dos Santos, E.V. Gusevskaya, Hydroformylation of recalcitrating biorenewable compounds containing trisubstituted double bonds, *Appl. Catal. A – Gen.* 591 (2020) 117406. <https://doi.org/10.1016/j.apcata.2019.117406>.
- [24] P. Pongrácz, B. Bartal, L. Kollár, L.T. Mika, Rhodium-catalyzed hydroformylation in γ -valerolactone as a biomass-derived solvent, *J. Organomet. Chem.* 847 (2017) 140–145. <https://doi.org/10.1016/j.jorganchem.2017.04.029>.
- [25] P. Pongrácz, H. Szentjóbi, T. Tóth, P. Huszthy, L. Kollár, Enantioselective hydroformylation of styrene in the presence of platinum(II)–monophospho-crown ether complexes, *Mol. Catal.* 439 (2017) 128–133. <https://doi.org/10.1016/j.mcat.2017.06.037>.

- [26] V.M. Fernández-Alvarez, V. de la Fuente, C. Godard, S. Castellón, C. Claver, F. Maseras, J.J. Carbó, Pd-Catalysed Mono- and Dicarbonylation of Aryl Iodides: Insights into the Mechanism and the Selectivity, *Chem. Eur. J.* 20 (2014) 10982–10989. <https://doi.org/10.1002/chem.201403022>.
- [27] M. Gergely, A. Bényei, L. Kollár, 2-Aminobenzimidazole and -benzoxazole as N-nucleophile in palladium-catalysed aminocarbonylation, *Tetrahedron* 76 (2020) 131079. <https://doi.org/10.1016/j.tet.2020.1310>.
- [28] P. Szuroczki, L. Molnár, A. Dörnyei, L. Kollár, Facile, High-Yielding Synthesis of 4-Functionalised 1,2,3-Triazoles via Amino- and Aryloxycarbonylation, *ChemistrySelect* 5 (2020) 448–451. <https://doi.org/10.1002/slct.201903801>.
- [29] L. Kollár, M.G. Varga, A. Dörnyei, A. Takács, Functionalisation of the uracil ring via palladium-catalysed aminocarbonylation, *Tetrahedron* 75 (2019) 4632–4639. <https://doi.org/10.1016/j.tet.2019.07.008>.
- [30] A. Brennfürer, H. Neumann, M. Beller, Palladium-Catalyzed Carbonylation Reactions of Aryl Halides and Related Compounds, *Angew. Chem. Int. Ed.* 48 (2009) 4114–4133. <https://doi.org/10.1002/anie.200900013>.
- [31] A. Takács, D. Marosvölgyi-Haskó, Z. Kabak-Solt, L. Damas, F.M.S. Rodrigues, R.M.B. Carrilho, M. Pineiro, M.M. Pereira, L. Kollár, Functionalization of indole at C-5 or C-7 via palladium-catalysed double carbonylation. A facile synthesis of indole ketocarboxamides and carboxamide dimers, *Tetrahedron* 72 (2016) 247–256. <https://doi.org/10.1016/j.tet.2015.11.007>.
- [32] R.M.B. Carrilho, M.M. Pereira, M.J.S.M. Moreno, A. Takács, L. Kollár, A new facile synthesis of steroid dimers containing 17,17'-dicarboxamide spacers, *Tetrahedron Lett.* 54 (2013) 2763–2765. <https://doi.org/10.1016/j.tetlet.2013.02.108>.
- [33] R.M.B. Carrilho, A.R. Almeida, M. Kiss, L. Kollár, R. Skoda-Földes, J.M. Dabrowski, M.J.S.M. Moreno, M.M. Pereira, One-Step Synthesis of Dicarboxamides through Pd-Catalysed Aminocarbonylation with Diamines as N-Nucleophiles, *Eur. J. Org. Chem.* (2015) 1840–1847. <https://doi.org/10.1002/ejoc.201403444>.
- [34] R.M.B. Carrilho, A.R. Almeida, M. Kiss, L. Kollár, R. Skoda-Földes, J.M. Dabrowski, M.J.S.M. Moreno, M.M. Pereira, One-Step Synthesis of Dicarboxamides through Pd-Catalysed Aminocarbonylation with Diamines as N-Nucleophiles, *Eur. J. Org. Chem.* (2015) 1840–1847. <https://doi.org/10.1002/ejoc.201403444>.
- [35] Web of ScienceTM (accessed on May 27th 2020). <https://apps.webofknowledge.com/>.

- [36] J. Biemolt, E. Ruijter, Advances in Palladium-Catalyzed Cascade Cyclizations, *Adv. Synth. Catal.* 360 (2018) 3821–3871. <https://doi.org/10.1002/adsc.201800526>.
- [37] L. Åkerbladh, S.Y. Chow, L.R. Odell, M. Larhed, Synthesis of 4*H*-Benzo[*e*][1,3]oxazin-4-ones by a Carbonylation–Cyclization Domino Reaction of *ortho*-Halophenols and Cyanamide, *ChemistryOpen* 6 (2017) 620–628. <https://doi.org/10.1002/open.201700130>.
- [38] P. Kalck, M. Urrutigoity, Tandem Hydroaminomethylation Reaction to Synthesize Amines from Alkenes, *Chem. Rev.* 2018, 118, 3833–3861. <https://doi.org/10.1021/acs.chemrev.7b00667>.
- [39] B.P. Bondžić, Rh catalyzed multicomponent tandem and one-pot reactions under hydroformylation conditions, *J. Mol. Catal. A: Chem.* 408 (2015) 310–334. <https://doi.org/10.1016/j.molcata.2015.07.026>.
- [40] M. Vilches-Herrera, L. Domke, A. Börner, Isomerization–Hydroformylation Tandem Reactions, *ACS Catal.* 4 (2014) 1706–1724. <https://doi.org/10.1021/cs500273d>.
- [41] A. Cunillera, M.D. de los Bernardos, M. Urrutigoity, C. Claver, A. Ruiz, C. Godard, Efficient Synthesis of Chiral γ -Aminobutyric Esters via direct Rhodium-Catalysed Enantioselective Hydroaminomethylation of acrylates, *Catal. Sci. Technol.* 10 (2020) 630–634. <https://doi.org/10.1039/C9CY01797F>.
- [42] D.E. Fogg, E.N. dos Santos, Tandem catalysis: a taxonomy and illustrative review, *Coord. Chem. Rev.* 248 (2004) 2365–2379. <https://doi.org/10.1016/j.ccr.2004.05.012>.
- [43] A.O. Dias, M.G.P. Gutiérrez, J.A.A. Villarreal, R.L.L. Carmo, K.C.B. Oliveira, A.G. Santos, E.N. dos Santos, E.V. Gusevskaya, Sustainable route to biomass-based amines: rhodium catalyzed hydroaminomethylation in green solvents, *Appl. Catal. A – Gen.* 574 (2019) 97–104. <https://doi.org/10.1016/j.apcata.2019.02.003>.
- [44] L. Kollár, P. Pongrácz, Tandem hydroformylation/aldol condensation reactions: Synthesis of unsaturated ketones from olefins, *J. Organomet. Chem.* 866 (2018) 184–188. <https://doi.org/10.1016/j.jorganchem.2018.04.031>.
- [45] B. Bartal, G. Mikle, L. Kollár, P. Pongrácz, Palladium catalyzed carbonylations of alkenyl halides with formic acid to get corresponding α,β -unsaturated carboxylic acids and esters, *Mol. Catal.* 467 (2019) 143–149. <https://doi.org/10.1016/j.mcat.2018.12.026>.
- [46] J. Cao, Z.-J. Zheng, Z. Xu, L.-W. Xu, Transition-metal-catalyzed transfer carbonylation with HCOOH or HCHO as non-gaseous C1 source, *Coord. Chem. Rev.* 336 (2017) 43–53. <https://doi.org/10.1016/j.ccr.2017.01.005>.

- [47] S.D. Friis, A.T. Lindhardt, T. Skrydstrup, The Development and Application of Two-Chamber Reactors and Carbon Monoxide Precursors for Safe Carbonylation Reactions, *Acc. Chem. Res.* 49 (2016) 594–605. <https://doi.org/10.1021/acs.accounts.5b00471>.
- [48] F. Xu, D. Lib, W. Han, Transition-metal-free carbonylation of aryl halides with arylboronic acids by utilizing stoichiometric CHCl_3 as the carbon monoxide-precursor, *Green Chem.* 21 (2019) 2911–2915. <https://doi.org/10.1039/c9gc00598f>.
- [49] J.-B. Penga, X. Qia, X.-F. Wu, Recent Achievements in Carbonylation Reactions: A Personal Account, *Synlett* 27 (2016) A–T. <https://doi.org/10.1055/s-0036-1588351>.
- [50] R.I. Khusnutdinov, T.M. Oshnyakova, U.M. Dzhemilev, Molybdenum compounds in organic synthesis, *Russ. Chem. Rev.* 86 (2017) 128–163. <https://doi.org/10.1070/RCR4617>.
- [51] M. Mamone, J. Aziz, J.L. Bescont, S. Piguel, Aminocarbonylation of N-Containing Heterocycles with Aromatic Amines Using $\text{Mo}(\text{CO})_6$, *Synthesis* 50 (2018) 1521–1526. <https://doi.org/10.1055/s-0037-1609152>.
- [52] L. Wu, Q. Liu, R. Jackstell, M. Beller, Carbonylations of Alkenes with CO Surrogates, *Angew. Chem. Int. Ed.* 53 (2014) 6310–6320. <https://doi.org/10.1002/anie.201400793>.
- [53] L. Åkerbladh, L.R. Odell, M. Larhed, Palladium-Catalyzed Molybdenum Hexacarbonyl-Mediated Gas-Free Carbonylative Reactions, *Synlett* 30 (2019) 141–155. <https://doi.org/10.1055/s-0037-1610294>.
- [54] M. Pineiro, L.D. Dias, L. Damas, G.L.B. Aquino, M.J.F. Calvete, M.M. Pereira, Microwave irradiation as a sustainable tool for catalytic carbonylation reactions, *Inorg. Chim. Acta* 455 (2017) 364–377. <https://doi.org/10.1016/j.ica.2016.06.043>.
- [55] C. Oliver Kappe, My Twenty Years in Microwave Chemistry: From Kitchen Ovens to Microwaves that aren't Microwaves, *Chem. Rec.* 19 (2019) 15–39. <https://doi.org/10.1002/tcr.201800045>.
- [56] S. Horikoshi, N. Serpone (Eds.), *Microwaves in Catalysis Methodology and Applications*, Wiley-VCH Verlag GmbH & Co. KGaA., Weinheim, 2016.
- [57] Elena Cini, E. Petricci, M. Taddei, Pd/C Catalysis under Microwave Dielectric Heating, *Catalysts* 7 (2017) 89. <https://doi.org/10.3390/catal7030089>.
- [58] I.M.Z. Fares, A.E.M. Mekky, A.H.M. Elwahy, I.A. Abdelhamid, Microwave-assisted three component synthesis of novel bis-fused quinazolin-8(4*H*)-ones linked to aliphatic or aromatic

spacer *via* amide linkages, Synth. Commun. 50 (2020) 893–903. <https://doi.org/10.1080/00397911.2020.1725575>.

[59] A.R. Almeida, R.D. Dias, C.J.P. Monteiro, A.R. Abreu, P.M.P. Gois, J.C. Bayon, M.M. Pereira, Rhodium-Catalysed Tandem Hydroformylation/Arylation Reaction with Boronic Acids, Adv. Synth. Catal. 356 (2014) 1223-1227. <https://doi.org/10.1002/adsc.201300968>.

[60] L. Damas, G.N.Costa, J.C. Ruas, R.M.B. Carrilho, A.R. Abreu, G. Aquino, M.J.F. Calvete, M. Pineiro, M.M. Pereira, Microwave assisted reactions of natural oils: transesterification and hydroformylation/isomerization as tools for high-value compounds, Curr. Microw. Chem. 2 (2015) 53-60, <https://doi.org/10.2174/221333560201150212110805>.

[61] F.M.S. Rodrigues, M.J.F. Calvete, C.J.P. Monteiro, S. A. C. Carabineiro, T.M.R. Maria, J.L. Figueiredo, M.M. Pereira, Hydroaminomethylation reaction as powerful tool for preparation of rhodium/phosphine functionalized nanomaterials. Catalytic evaluation in styrene hydroformylation, Catal. Today (2019), *in press*. <https://doi.org/10.1016/j.cattod.2019.05.045>.

[62] A.R. Almeida, R.M.B. Carrilho, A.F. Peixoto, A.R. Abreu, A. Silva, M.M. Pereira, Sequential reactions from catalytic hydroformylation toward the synthesis of amino compounds, Tetrahedron 73 (2017) 2389-2395. <https://doi.org/10.1016/j.tet.2017.03.023>.

[63] F.M.S. Rodrigues, P.K. Kucmierczyk, M. Pineiro, R. Jackstell, R. Franke, M. Beller, M.M. Pereira, Dual Rh-Ru catalysts for reductive hydroformylations of olefins to alcohols, ChemSusChem. 11 (2018) 2310-2314. <https://doi.org/10.1002/cssc.201800488>.

[64] L. Damas, R.M.B. Carrilho, S.C.C. Nunes, A.A.C.C. Pais, L. Kollar, M. Pineiro, M.M. Pereira, A novel Pd-catalysed sequential carbonylation/cyclization approach toward bis-N-heterocycles: rationalization by electronic structure calculations, R. Soc. open sci. 5, (2018) 181140. <https://doi.org/10.1098/rsos.181140>.

[65] J. Schniedermeier, H.J. Haupt, New rhodium(I)- π -chelate complexes with coordinated amidine bases (dbu, dbn) and their catalytic properties to polymerize phenylacetylene, J. Organomet Chem. 506 (1996) 41-47. [https://doi.org/10.1016/0022-328X\(95\)05689-M](https://doi.org/10.1016/0022-328X(95)05689-M).

[66] J.V. Barkley, C.J. Davies, B.T. Heaton, C. Jacob, Rhodium complexes with 1,8-diazabicyclo[5.4.0]undec-7-ene (dbu): crystal structure of *trans*-[Rh(CO)(dbu)(PPh₃)₂]ClO₄, J. Chem. Soc. Dalton Trans. (1995) 2861-2863. <https://doi.org/10.1039/DT9950002861>.

[67] A.J. von Wangelin, H. Neumann, D. Gördes, S. Klaus, D. Strübing, M. Beller, Multicomponent Coupling Reactions for Organic Synthesis: Chemoselective Reactions with

Amide–Aldehyde Mixtures, *Chem. Eur. J.* 9 (2003) 4286-4294.
<https://doi.org/10.1002/chem.200305048>.

[68] C. Gomes, C.S. Vinagreiro, L. Damas, G. Aquino, J. Quaresma, C. Chaves, J. Pimenta, J. Campos, M. Pereira, M. Pineiro, Advanced Mechanochemistry Device for Sustainable Synthetic Processes, *ACS Omega* 5 (2020) 10868-10877. <https://doi.org/10.1021/acsomega.0c00521>.

[69] J-F. Soulé, H. Miyamura, S. Kobayashi, Powerful Amide Synthesis from Alcohols and Amines under Aerobic Conditions Catalyzed by Gold or Gold/Iron, -Nickel or -Cobalt Nanoparticles, *J. Am. Chem. Soc.* 133 (2011) 18550–18553. <https://doi.org/10.1021/ja2080086>.

[70] D.F. Harvey, D.M. Sigano, Synthesis of Cyclopropylpyrrolidines via Reaction of *N*-Allyl-*N*-propargylamides with a Molybdenum Carbene Complex. Effect of Substituents and Reaction Conditions, *J. Org. Chem.* 61 (1996) 2268-2272. <https://doi.org/10.1021/jo9519930>.

[71] N.G. Moon, A.M. Harned, Iodine(III)-promoted synthesis of oxazolines from *N*-allylamides, *Tetrahedron Lett.* 54 (2013) 2960–2963. <http://doi.org/10.1016/j.tetlet.2013.03.140>.

Highlights

- Pd-catalysed aminocarbonylation of iodoaromatics using MW irradiation and Mo(CO)_6 as CO source
- Synthesis of *N*-heterocyclic and chalcone-derived allylcarboxamides
- Catalytic aminocarbonylation/hydroformylation sequential process
- Synthesis of biologically relevant *N*-heterocyclic and chalcone derived formylcarboxamides

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof