

Published on Web 03/03/2007

Palladium-Catalyzed Methylation and Arylation of sp² and sp³ C–H Bonds in Simple Carboxylic Acids

Ramesh Giri, Nathan Maugel, Jiao-Jie Li, Dong-Hui Wang, Steven P. Breazzano,

Lindsey B. Saunders, and Jin-Quan Yu*

Department of Chemistry MS015, Brandeis University, Waltham, Massachusetts 02454-9110 Received January 13, 2007; E-mail: yu200@brandeis.edu

Palladium-catalyzed cross-coupling reactions are among the most widely used carbon–carbon bond-forming reactions in organic synthesis.¹ Extensive development of a wide range of ligands has allowed the use of ArCl,^{2a-c} alkyl halides,^{2d} and sterically hindered 2,6-substituted ArX^{2e} as coupling partners. The decarboxylative Heck or Suzuki coupling using ArCOOH is another practically appealing approach since carboxylic acids are readily available.³ The potential of Pd-catalyzed C–H activation reactions for developing synthetically useful carbon–carbon bond forming reactions prompted our initial efforts to couple sp² and sp³ C–H bonds in oxazoline and pyridine substrates with organotin and organoboron reagents.^{4,5} However, these types of substrates seriously restrict the substrate scope since the oxazolines require installation and removal and pyridines lack versatility for further synthetic manipulations.

Considering the broad utility of hydroxyl and carboxyl functionalities as directing groups in asymmetric catalysis,⁶ the use of simple functional groups to direct C-H activation/C-C coupling processes could prove fruitful in developing C-C bond forming reactions.7 Carboxyl-directed lactonization of sp3 C-H bonds catalyzed by K2PtCl4 has been previously observed by Sen, Sames, and others.8 Ortho alkenylation of benzoic acid catalyzed by Pd-(OAc)₂ has been reported by Miura.⁹ Despite these significant advances, Pd-catalyzed C-H activation/C-C coupling reactions using a carboxyl group (eqs 1,2) have not been developed. In particular, Pd-catalyzed coupling of sp³ β -C-H bonds in aliphatic acids with organometallic reagents is an unanswered challenge and yet a significant goal from the viewpoint of synthetic applications. Herein, we report the first catalytic protocol for the coupling of both o-C-H bonds in benzoic acids and β -C-H bonds in aliphatic acids with organoboron reagents via PdII/Pd⁰ catalysis. The generality of this newly observed Pd-insertion into β -C-H bonds was further demonstrated by β -arylation of aliphatic acids using ArI in which Pd^{II}/Pd^{IV} catalysis is likely responsible.¹⁰⁻¹¹

$$\square H \xrightarrow{\text{COOH}}_{\text{Pd}^{II}} \xrightarrow{\text{R-M}}_{\text{Pd}^{II}} \square \xrightarrow{\text{COOH}}_{\text{R}} (1) \xrightarrow{\text{R-M}}_{\text{R_1}} \xrightarrow{\text{R-M}}_{\text{R_2}} \xrightarrow{\text{R-M}}_{\text{Pd}^{II}} \xrightarrow{\text{R-M}}_{\text{R_1}} \xrightarrow{\text{R-M}}_{\text{R_2}} (2)$$

The formation of phenoxide in the presence of Cs_2CO_3 accelerates both intra-^{12a} and intermolecular^{12b} ortho arylation of phenols using a combination of Pd⁰ and ArI. This effect was attributed to either an increase of the electron density of the phenyl ring or an enhanced binding of the ArPdI species to the phenoxide. Inspired by these observations, we focused on the development of Pd(OAc)₂catalyzed C–H functionalization processes directed by carboxyl groups. If successful, these types of processes could be applicable to a wide range of simple substrates containing acidic protons. On the basis of the Pd-catalyzed coupling protocols recently developed in our laboratory,^{4b} we began to search for coupling conditions using simple carboxylic acids as substrates. Extensive screening of conditions (see Supporting Information) led us to discover that the use of sodium carboxylates as substrates was effective for the desired transformation. Thus, stirring sodium toluate **1a** with 0.5 equiv of benzoquinone, 1 equiv of Ag₂CO₃,¹³ 3 equiv of MeB(OH)₂ and 10 mol % Pd(OAc)₂ in *tert*-amyl alcohol or *tert*-BuOH at 100 °C for 3 h affords ortho methylated product **1b** in 50% isolated yield (eq 3). Arylation of **1a** using a phenylboronate **2** as the coupling partner proceeds under similar conditions at 120 °C to give ortho arylated product **1c** in 45% isolated yield (eq 3). Based on our previous studies,^{4b} this catalytic reaction most likely proceeds via a Pd^{II}/Pd⁰ catalysis. *Although the role of sodium counterions remains to be elucidated, we have shown that the preformed palladium toluate*¹⁴ *in the absence of sodium counterions is not reactive.* This led us to hypothesize that the electronically enriched carbonyl instead of the O-anion of the sodium carboxylate binds the Pd^{II} in the C–H cleavage step.



Guided by the mechanism of the Suzuki coupling reaction, we reasoned that the yield could be further improved by employing a suitable base to enhance the transmetalation step in the coupling reaction. We screened a wide range of bases and found that K_2HPO_4 increases the yields of **1b** and **1c** to 75% and 63%, respectively (Table 1, entries 1, 3). Since the presence of K_2HPO_4 leads to the in situ formation of carboxylates, benzoic acids instead of sodium carboxylates are used as substrates under these new conditions.

Although the arylation of benzoic acid gives a mixture of monoand diarylated compounds in 50% yield (3:2 ratio), we found that the meta substituted benzoic acids lead to regioselective arylation, albeit in lower yields (Table 1, entries 4-6). The tolerance of OMe and CO₂Me groups allows for the preparation of a wide range of 1,2,4-substituted arenes.

Encouraged by these results, we attempted the coupling of β -C-H bonds in aliphatic acids with **2**. The sodium carboxylate of **6** was subjected to the arylation conditions in eq 3 to give

Table 1. Ortho Methylation and Arylation of Benzoic Acids^a



^{*a*} Conditions: 10 mol % Pd(OAc)₂, 0.5 equiv of benzoquinone, 1 equiv of Ag₂CO₃, 1.5 equiv of K₂HPO₄, 2 equiv of MeB(OH)₂ or 1 equiv of **2**, *tert*-BuOH, 120 °C, 3 h. ^{*b*} Yield at 100 °C.



^a Conditions: 10 mol % Pd(OAc)₂, 1 equiv of 2, 0.5 equiv of benzoquinone, 1 equiv of Ag₂CO₃ and 1.5 equiv of K₂HPO₄. ^b Yields of their methyl esters. Less than 2% diarylated products were observed in 6-10.

Table 3. β -Arylation of Aliphatic Acids Using Arl^a



entry	product	yield (%) ^b	entry	product	yield (%) ^b
1	R = Me, Ar = Ph 6a/6b , 5:2	70	5	$R = {}^{r}Bu_{,,} Ar = Ph$ 8a/8b , 4:1	62
2	R = Et, Ar = Ph 7a/7b, 4:1	72	6	$R = (CH_2)_3OBn, Ar = Ph$ 9a/9b, 5:1	45
3	R = Pr, Ar = pMePh 12a/12b , 3:1	60	7	$R = (CH_2)_2 CO_2 Me$, $Ar = Ph$ 10a/10b , 5:1	42 ^c
4	$R = {}^{n}Pr, Ar = p-BrPh$ 12c/12d, 3:1	63	8	$R = (CH_2)_2CO_2Me$, $Ar = p$ -MePh 10c/10d , 5:1	43

^a Conditions: 10 mol % Pd(OAc)₂, 2 equiv of aryl iodide, 2 equiv of Ag₂CO₃, 1 equiv of K₂HPO₄, and 2 equiv of NaOAc. ^b Yields of their methyl esters. ^c 10b was not isolated. The yield is based on 10a.

predominantly the monoarylated product 6a in 30% isolated yield. The use of Ag₂O or AgOAc in place of Ag₂CO₃ as an oxidant afforded less than 5% of 6a. The presence of Na₂CO₃ results in a complete loss of reactivity with aliphatic acids. However, the potassium carboxylate of 6 generated in situ using K₂HPO₄ affords 6a in 38% isolated yield (Table 2, entry 1).

While the catalytic turnover remains to be improved, the observed mono-selectivity is a highly desirable advantage. Benzylethers and esters were also tolerated (Table 2, entries 4-5), thus making this protocol potentially applicable to organic synthesis. The preferential arylation of the cyclopropyl C-H bond is worth noting since examples of Pd insertion into methylene C-H bonds are still rare (entry 6).15

Importantly, this coupling reaction provides the first example for carboxyl-directed Pd-insertion into sp³ β -C-H bonds in simple aliphatic acids. To demonstrate the generality of this C-H cleavage reactivity in C-C bond forming reactions, we carried out arylation reactions using ArI as the arylating reagents.^{15–17} We found that alteration of our coupling protocol by omitting the benzoquinone and using PhI as the coupling partner led to mono- and diarylation of aliphatic acid **6** in 40% combined yield (6a/6b = 5:2). We further discovered that the use of 2 equiv of NaOAc as an additive substantially increases the combined yield of 6a and 6b to 70% (Table 3).¹⁸

This arylation reaction most likely involves a COOH directed Pd insertion into C-H bonds and subsequent oxidation of the RPdIIcomplex to (R)(Ar)Pd^{IV}I intermediate by ArI. The formation of diarylated products (Table 3) is consistent with the PdII/PdIV pathway in which the Pd^{II}, unlike the Pd⁰ in the cross-coupling protocol, remains bound to the carboxylate and results in further arylation. Ag₂CO₃ is mainly responsible for the catalytic turnover by converting PdI2 into the reactive PdII species.^{16a} The excess AcO-

could also displace the iodide from the (R)(Ar)Pd^{IV}I intermediate and increase the turnover number as observed.

In summary, we have observed the first example of Pd-insertion into sp³ β -C-H bonds in simple aliphatic acids. A promising protocol for the coupling of sp² and sp³ C-H bonds in simple carboxylic acids with organoboron reagents has been established. The potential of carboxyl-directed C-H activation in developing C-C bond-forming reactions is also demonstrated by the arylation of β -C-H bonds in aliphatic acids using ArI. We are currently optimizing the conditions to improve the yields of these reactions.

Acknowledgment. We thank Brandeis University for financial support and the Camille and Henry Dreyfus Foundation for a New Faculty Award. We are also grateful to the Myron Rosenblum Endowed Fellowship to R.G and the NSF for a REU Fellowship to S.P.B.

Supporting Information Available: Experimental procedure and characterization of all new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

References

- For reviews see: (a) Diederich, F., Stang, P. J., Eds. *Metal-Catalyzed Cross-Coupling Reactions*; Wiley-VCH: New York, 1998. (b) Nicolaou, K. C.; Bulger, P. G.; Sarlah, D. Angew. Chem., Int. Ed. 2005, 44, 4442.
- (2) (a) Littke, A. F.; Fu, G. C. Angew. Chem., Int. Ed. 1999, 38, 2411. (b) Wolfe, J. P.; Buchwald, S. L. Angew. Chem., Int. Ed. 1999, 38, 2413. (c) Kawatsura, M.; Hartwig, J. F. J. Am. Chem. Soc. **1999**, *121*, 1473. (d) Netherton, M. R.; Dai, C.; Neuschütz, K.; Fu, G. C. J. Am. Chem. Soc. 2001, 123, 10099. (e) Barder, T. E.; Walker, S. D.; Martinelli, J. R.; Buchwald, S. L. J. Am. Chem. Soc. 2005, 127, 4685.
 (3) (a) Myers, A. G.; Tanaka, D.; Mannion, M. R. J. Am. Chem. Soc. 2002,
- 124, 11250. (b) Gooßen, L. J.; Deng, G.; Levy, L. M. Science 2006, 313, 662.
- (4) (a) Chen, X.; Li, J.-J.; Hao, X.-S.; Goodhue, C. E.; Yu, J.-Q. J. Am. Chem. Soc. 2006, 128, 78. (b) Chen, X.; Goodhue, C. E.; Yu, J.-Q. J. Am. Chem. Soc., 2006, 128, 12634.
- (5) For a Rh-catalyzed ortho arylation of 2-phenylpyridine using tetraaryltin see: Oi, S.; Fukita, S.; Inoue, Y. *Chem. Commun.* **1998**, 2439. (a) Katsuki, T.; Sharpless, K. B. *J. Am. Chem. Soc.* **1980**, *102*, 5974. (b)
- Brown, J. M. Chem. Soc. Rev. 1993, 22, 25. (c) Halpern, J. Science 1982, 217.401.
- (7) (a) For a remarkable Ru-catalyzed ortho arylation of acetophenone using a phenylboronate see: Kakiuchi, F.; Kan, S.; Igi, K.; Chatani, N.; Murai, S. J. Am. Chem. Soc. 2003, 125, 1698. (b) For an example of Rh-catalyzed alkenylation of aryl C–H bonds directed by imine see: Thali, R. K.; Ellman, J. A.; Bergman, R. G. J. Am. Chem. Soc. **2004**, *126*, 7192. (c) For a Pd-catalyzed arylation of sp³ C–H bonds in 2,4,6-tri-tert-butylbromobenzene with PhB(OH)₂ see: Barder, T. E.; Walker, S. D.; Martinelli, J. R.; Buchwald, S. L. J. Am. Chem. Soc. **2005**, *127*, 4685. (d) For a Pd-catalyzed arylation of vinylic C-H bonds with PhB(OH)₂ see: Delcamp, J. H.; White, M. C. J. Am. Chem. Soc. 2006, 128, 15076.
- (a) Kao, L.-C.; Sen, A. J. Chem. Soc., Chem. Commun. 1991, 1242. (b) (d) Rub, J. S., Johnson, J. A.; Sames, D. J. Am. Chem. Soc. 2001, 123, 8149. (c) Lee, J. M.; Chang, S. Tetrahedron Lett. 2006, 47, 1375.
 (9) Miura, M.; Tsuda, T.; Satoh, T.; Pivsa-Art, S.; Nomura, M. J. Org. Chem.
- 1998, 63, 5211.
- (10) For avlation of sp³ C-H bonds using an amide-pyridine directing group and ArI as the arylating reagent see: Zaitsev, V. G.; Shabashov, D.; Daugulis, O. J. Am. Chem. Soc. 2005, 127, 13154.
- For arylation of aryl and benzylic C-H bonds using [Ph₂I]BF₄ or [Ph₂I] For arylation of aryl and benzylic C-H bonds using [Ph₂I]BF₄ or [Ph₂I] FF₆ see: (a) Kalyani, D.; Deprez, N. R.; Desai, L. V.; Sanford, M. S. J. Am. Chem. Soc. **2005**, *127*, 7330. (b) Daugulis, O.; Zaitsev, V. G. Angew. Chem., Int. Ed. **2005**, *44*, 4046. (11)
- (12) (a) Hennings, D. D.; Iwasa, S.; Rawal, V. H. J. Org. Chem. 1997, 62, 2. (b) Satoh, T.; Kawamura, Y.; Miura, M.; Nomura, M. Angew. Chem., Int. Ed. 1997, 36, 1740.
- (13) Ag₂CO₃ releases Ag⁺ and CO₃²⁻ slowly during the reaction which appears to be important for the catalysis
- (14) Hermans, S.; Wenkin, M.; Devillers, M. J. Mol. Catal. A: Chem. 1998, 136. 59.
- (15) Reddy, B. V. S.; Reddy, L. R.; Corey, E. J. Org. Lett. 2006, 8, 3391.
- (16) For examples of Pd-catalyzed methylation or arylation of sp² C-H bonds using MeI or ArX see: (a) Tremont, S. J.; Rahman, H. U. J. Am. Chem. Soc. 1984, 106, 5759. (b) Catellani, M.; Chiusoli, G. P. J. Organomet. Chem. 1985, 286, C13. (c) Catellani, M.; Frignani, F.; Rangoni, A. Angew. *Chem., Int. Ed.* **1997**, *36*, 119. (d) Campo, M. A.; Huang, Q.; Yao, T.; Tian, Q.; Larock, R. C. *J. Am. Chem. Soc.* **2003**, *125*, 11506. (e) Bressy, C.; Alberico, D.; Lautens, M. *J. Am. Chem. Soc.* **2005**, *127*, 13148.
- (17) For arylation of sp3 C-H bonds in 1-tert-butyl-2-iodobenzene see: Dyker, G. Angew. Chem., Int. Ed. 1992, 31, 1023.
- (18) The arylation of toluic acid 1 under the same conditions gives the ortho arylated product in 95% yield.

JA0701614