



Subscriber access provided by Nottingham Trent University

## Communication

### 1,2-Boron Shifts of #-Boryl Radicals Generated from Bis-Boronic Esters using Photoredox Catalysis

Daniel Kaiser, Adam Noble, Valerio Fasano, and Varinder K. Aggarwal

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.9b07564 • Publication Date (Web): 28 Aug 2019

Downloaded from pubs.acs.org on August 28, 2019

#### Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

## **1,2-Boron Shifts of β-Boryl Radicals Generated from Bis-Boronic Esters** using Photoredox Catalysis

Daniel Kaiser, Adam Noble, Valerio Fasano, and Varinder K. Aggarwal\*

School of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, U.K.

Supporting Information Placeholder

**ABSTRACT:** 1,2-Bis-boronic esters are versatile intermediates that enable the rapid elaboration of simple alkene precursors. Previous reports on their selective mono-functionalization have targeted the most accessible position, retaining the more hindered secondary boronic ester. In contrast, we have found that photoredox-catalyzed mono-deboronation generates primary  $\beta$ -boryl radicals that undergo rapid 1,2-boron shift to form thermodynamically favored secondary radicals, allowing for selective transformation of the more hindered boronic ester. The pivotal 1,2-boron shift, which has been demonstrated to be stereoretentive, enables access to a wide range of functionalized boronic esters and has been applied to highly diastereoselective fragmentation and transannular cyclization reactions. Furthermore, its generality has been shown in a radical cascade reaction with an allylboronic ester.

Photoredox chemistry enables radical species to be generated from a broad range of functional groups under exceptionally mild conditions.<sup>1</sup> We and others have shown that readily available alkylboron reagents, including boronic esters<sup>2</sup> and trifluoroborate salts,<sup>3</sup> undergo deboronative single-electron oxidation to give alkyl radical intermediates that engage in Giese-type additions,<sup>2a-c,3a-d</sup> hydrogen atom transfer,<sup>2f</sup> nickelcatalyzed cross couplings,<sup>2e,3e-h</sup> and radical-polar cross-over reactions (Scheme 1A).<sup>2d,3l</sup> Despite these diverse photoredoxcatalyzed transformations of alkylborons, related reactions that use 1,2-bis-boron species as radical precursors have yet to be explored. However, such processes would be highly valuable since (i) 1,2-bis-boronic esters are easily prepared from alkenes, often with high enantioselectivity;<sup>4</sup> and (ii) following the deboronative radical reaction, a boronic ester is retained for use in further transformations.5

Previous reports of mono-functionalizations of 1,2-bis-boronic esters (1) have involved selective reaction at the primary boronic ester, which is due to favorable activation (boronate complex formation) of the sterically less hindered boron atom (Scheme 1B).<sup>6</sup> Based on this precedent, selective formation of boronate complex 2 followed by photoredox-catalyzed singleelectron oxidation and reaction of the resulting primary radical would give a product bearing a secondary boronic ester (Scheme 1C, path A). However, given the instability of primary alkyl radicals, we considered the possibility of a radical 1,2boron shift to provide the thermodynamically more stable secondary radical (Scheme 1C, path B). Thermodynamically driven 1,2-group transfers of alkyl radicals

Scheme 1. Alkylborons as radical precursors and functionalizations of 1,2-bis-boronic esters.

A Radical precursors based on alkylboron species



B Selective reaction at the less hindered position of 1,2-bis-boronic esters



C Possible pathways for radicals generated from 1,2-bis-boronic esters



are mainly limited to migrations of  $\pi$ -systems, such as the well-established neophyl and Dowd-Beckwith rearrangements.7 Related heteroatom-transfers are rare but have been proposed for halogen<sup>8</sup> and silicon groups.<sup>9</sup> There have been isolated reports of 1,2-boron migrations, including via cationic<sup>10</sup> or anionic intermediates,<sup>11</sup> and a single report by Batey in 1999 which showed that  $\beta$ -boryl radicals are capable of undergoing intramolecular homolytic substitution.<sup>12</sup> In our proposed strategy, a radical migration of the boronic ester group would enable a cascade sequence of 1,2-shift and subsequent reaction of the resulting thermodynamically favored secondary alkyl radical (Scheme 1C, path B). This would lead to a product in which it appeared as if the more hindered secondary boronic ester had been activated, providing the opposite selectivity to that typically observed for functionalizations of 1,2-bis-boronic esters.<sup>6,13</sup> We now report that, in photoredox-catalyzed deboronative Giese reactions of 1,2-bis-boronic esters, following activation of the less hindered primary boronic ester, the

59

60

exclusive product obtained is that derived from substitution of the secondary boronic ester.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

59

60

Our initial investigations of the reaction of 1,2-bis-boronic ester **1a** were based on our recent report of a deboronative cvclobutane synthesis, where phenyllithium was successfully employed to form a highly reducing boronate complex (Table 1).2d Thus, a boronate complex was formed by reaction of **1a** with phenyllithium (A) in THF, and a solution of photoredox catalyst (4CzIPN), tert-butyl acrylate, and tert-butanol in DMF was added. The subsequent reaction under blue light irradiation afforded the product of functionalization at the secondary position (3a) in a promising 48% yield (entry 1). Two characteristics of this initial reaction are noteworthy: (i) no trace of regioisomeric product 4 was detected in the crude reaction mixture, confirming that a 1,2-boron shift occurred to form the more stable secondary radical: and (ii) we observed 12% of doubly functionalized product 5 and recovered 27% of 1a. This suggested low selectivity between single and double addition of A to 1a. Hoping to increase the selectivity, we turned to more sterically hindered aryllithium reagents (B-G, entries 2-7). Pleasingly, this approach proved successful, with ortho-substituted aryllithiums suppressing the formation of 5. From this series of reactions, B emerged as the ideal reagent in terms of vield (68%), cost and reproducibility. A subsequent screen of solvents identified acetonitrile as optimal (entry 8).14 Finally, a slight increase in concentration (compare entries 8-10) gave **3a** in 90% isolated yield, while the absence of <sup>t</sup>BuOH (required for protonation of the enolate intermediate; entry 11), light (entry 12) or photocatalyst (entry 13) led to dramatically diminished yields.

28 With the optimized conditions in hand, we investigated the 29 scope of bis-boronic esters (Scheme 2A). Initially, we probed 30 the effects of steric hindrance on the reaction efficiency and were pleased to see that products with  $\alpha$ -secondary (3b) and 31  $\alpha$ -tertiary centres (3c) were formed readily. Moreover, sub-32 strates containing a tertiary boronic ester (3d-3f) provided ex-33 cellent yields of the products of tertiary functionalization. It 34 should be noted that for substrates showing considerable steric 35 bias between the two boronic ester groups (e.g., 3e), using phe-36 nyllithium instead of B provided almost identical results (81% 37 compared to 85% for B). Surprisingly, phenyl-containing 3g 38 was only isolated in moderate yield (35%) and an equimolar amount of starting material was recovered. We speculated that 39 competitive addition of the transient radical intermediate to 40 the aryl ring could (i) lead to unspecific degradation, as indi-41 cated by the low mass balance; and (ii) slow down turnover of 42 the photocatalyst, thereby allowing the boronate complex in-43 termediate to decompose over time.14 When a homologated 44 substrate was used, product **3h** was afforded in considerably increased yield (66%), suggesting the vicinity of the phenyl 45 moiety was in fact inhibiting formation of 3g. Unfortunately, 46 styrene-derived bis-boronic esters failed to give the desired 47 products.14 In addition to substrates of varying steric demand, 48 various functional groups were also tolerated, including esters 49 (3i), nitriles (3j) and silvl ethers (3k)—the former two requir-50 ing the use of less hindered aryllithium **D** in order to suppress 51 competing  $\alpha$ -deprotonation. Additionally, using two equiva-52 lents of arvllithium allowed isolation of secondary carbamate 53 31 in moderate vield. Bis-boronic esters derived from 3-carene and camphene were also competent substrates, affording the 54 corresponding products 3m and 3n in good yields, with excel-55 lent diastereoselectivity observed for the latter product. The 56 1,2-bis-boronic ester derivative of  $\beta$ -pinene (10) provided 57 monocyclic product 30 in 53% yield. This compound is the 58

product of ring-opening of the strained cyclobutane moiety after 1,2-boron shift, highlighting the radical nature of this transformation.

#### **Table 1. Optimization Studies**

S Bpin		ArLi (1.1 equiv), THF then			Bpin	
Bpin 1a	<sup>t</sup> Bu-ac sol	4CzIPN (5 n <mark>rylate</mark> (2 equiv), vent, blue LEDs	nol %) <sup>t</sup> BuOH (3 eq , 35 °C, 16 h	uiv)	CO <sub>2</sub> <sup>t</sup> Bu 3a	
Entry <sup>a</sup>	ArLi	Solvent	c (M)	<b>3a</b> (%) <sup>b</sup>	<b>1a</b> (%) <sup>b</sup>	
1	Α	DMF	0.05	48	27	
2	В	DMF	0.05	68	12	
3	С	DMF	0.05	38	27	
4	D	DMF	0.05	75	15	
5	Ε	DMF	0.05	70	9	
6	F	DMF	0.05	70	2	
7	G	DMF	0.05	29	11	
8	В	MeCN	0.05	88	2	
9	В	MeCN	0.1	100 (90)	0	
10	В	MeCN	0.2	89	5	
11 <sup>c</sup>	В	MeCN	0.1	11	37	
$12^d$	В	MeCN	0.1	0	17	
13 <sup>e</sup>	В	MeCN	0.1	<1	33	
4, not dete	CO <sub>2</sub> <sup>t</sup> Bu	Li <sup>i</sup> P A			Li D	
5	℃O₂ <sup>t</sup> Bu CO₂ <sup>t</sup> Bu	E	I (	F	G	

<sup>*a*</sup> Reactions were run on 0.2 mmol scale. <sup>*b*</sup> Determined by GC/MS analysis; values in parentheses correspond to isolated yields. <sup>*c*</sup> Without <sup>*t*</sup>BuOH. <sup>*d*</sup> In the dark. <sup>*e*</sup> Without 4CzIPN.

Our interest subsequently shifted to investigation of the radical acceptor (Scheme 2B). Radical conjugate addition to a range of electron-deficient alkenes afforded the desired products (**3p**-**3s**) in good to excellent yields. Styrene derivatives were also found to be competent radical acceptors, with pentafluorostyrene efficiently affording **3t**. On the other hand, 4-vinylpyridine, a comparatively weaker acceptor, only reacted with more nucleophilic tertiary radicals, forming **3u** in 75% yield. Employing an enantiopure dehydroalanine derivative

provided product **3v** in only moderate yield, but with complete diastereocontrol. While the aforementioned reactions all require protonation to form the corresponding products, we were interested in exploring alternative modes of termination. In this sense, enoate **3w** was synthesized via two mechanistically distinct addition-elimination sequences from either the allylic sulfone (radical elimination, 45%) or the allylic acetate (polar elimination, 55%). Following our recent report on **Scheme 2. Reaction Scope.**<sup>*a*</sup>

radical addition-polar cyclization cascades,<sup>2d</sup> formation of cyclopropane 3x was also shown to be a viable pathway for this transformation. Additionally, acridine reacted readily with the secondary radical generated from **1a**, providing **3y** in excellent yield.



<sup>*a*</sup> Reactions were run on 0.2 mmol scale, unless otherwise noted. Yields are of isolated products. <sup>*b*</sup> Isolated as the corresponding alcohols after oxidation. <sup>*c*</sup> Using aryllithium **D**. <sup>*d*</sup> Using 2.0 equiv **B**. <sup>*e*</sup> Run on 0.15 mmol scale. <sup>*f*</sup> Without <sup>*t*</sup>BuOH.

As a demonstration of the synthetic utility of this methodology, we sought to highlight its application in the synthesis of valuable structures (Scheme 2C). By using a diboration/photoredox-catalyzed deboronative Giese reaction/oxidation/lactonization sequence, we were able to transform unfunctionalized alkenes into lactones **6a** and **6b** in good yields over four steps.

Based on previous reports on the oxidation of boronate complexes,<sup>2,3</sup> and the observed regioselectivity of our reaction, we propose the mechanism depicted in Scheme 3. Initial oxidation of primary boronate complex **2** by the excited state photoredox catalyst generates a primary radical with concomitant loss of an equivalent of arylboronic ester **7**.<sup>15</sup> Subsequent equilibration of the radical species through 1,2-boron shift of **8** affords thermodynamically favored secondary radical **9**. Addition of **9** to the radical acceptor forms electron deficient radical **10**, which accepts an electron from the reduced form of the

photoredox catalyst. Finally, anion  $\mathbf{11}$  is protonated to afford product  $\mathbf{3}$ .

Scheme 3. Mechanistic Proposal.



At this point, we were intrigued as to whether our approach could be extended to the selective functionalization of 1,2,3tris-boronic esters (Scheme 4A). We therefore subjected **12** to our reaction conditions and were pleased to isolate **13**, the product of sequential double 1,2-shift of two boronic esters in 79% yield. Notably, this product is formed with impeccable selectivity, highlighting the thermodynamic control that favors the most stabilized radical center as the site of reaction.

Given the ease in which 1,2-bis-boronic esters can be prepared from readily available alkenes, we were keen to explore substrates derived from dienes, which could potentially undergo radical cascade reactions. We were attracted by cyclooctadiene derivatives, which could undergo transannular radical reactions (Scheme 4B). Pleasingly, under the standard conditions, diborated cyclooctadiene (**14**) reacted to give bicyclic product **15** in 51% yield and with excellent diastereoselectivity, alongside uncyclized product **16** (33%). This unusual radical cyclization can be viewed as both a *5-exo-* and *5-endo-trig* cyclization, accounting for its relatively low rate of ring closure ( $k = 3.3 \times 10^4 \text{ s}^{-1}$ ),<sup>16</sup> and consequently significant formation of the direct trapping product **16**.<sup>17</sup>

# Scheme 4. Additional Studies.



<sup>*a*</sup> Isolated as the corresponding alcohols after oxidation.

We subsequently turned our attention to the 1,2-boron shift. If this occurs via an intramolecular homolytic substitution, we reasoned that it could proceed with high stereospecificity. Thus, we investigated the stereochemical outcome of the reaction using diastereomerically pure **1**z, derived from  $\alpha$ -pinene (Scheme 5A). The expected product of 1,2-shift and subsequent ring opening (**3**z) was obtained in 52% yield and with >20:1 d.r., in which the boronic ester migrated with high stereochemical fidelity. Further support for this was provided by the

formation of **3m** from diborated 3-carene (Scheme 2A), in which boron migration occurred with complete stereospecificity but subsequent reaction of the resulting tertiary radical proceeded with poor diastereoselectivity.

Unequivocal proof for the 1,2-boron shift was obtained by treatment of allylboronic ester 17 with Langlois' reagent under Nicewicz's hydrotrifluoromethylation conditions, which gave 20 in 96% yield (Scheme 5B).<sup>18</sup>. This reaction proceeds via addition of a trifluoromethyl radical to 17 to provide secondary β-boryl radical 18. 1,2-Transposition of the boronic ester affords the thermodynamically favored tertiary radical 19, which undergoes hydrogen atom transfer (HAT) with thiophenol to form **20**. The high regioselectivity observed implies that 1,2boron shift from a tertiary (18) to a secondary position (19) is much more rapid than HAT with thiophenol, which has a rate of  $k = 1 \times 10^8 \,\text{M}^{-1} \,\text{s}^{-1}$  for secondary alkyl radicals.<sup>19</sup> Further insight into the facile nature of the 1,2-shift was provided by modelling the process using DFT, which revealed a barrier of only 8.1 kcal mol<sup>-1</sup> for migration of a secondary boronic ester to a primary radical (Scheme 5C). Finally, boron-isotope labelling studies using mono-labelled 1,2-bis-boronic ester 10B-1b yielded unlabelled product **3b** and labelled arylboronic ester <sup>10</sup>B-7, confirming that boronate complex formation with aryllithium B occurs exclusively at the primary boronic ester (Scheme 5D).

#### Scheme 5. Investigations into the 1,2-Boron Shift.

2

3

4

5

6 7

8

9

10

11 12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58 59

60



 $^a$  Isolated as the corresponding alcohols after oxidation.  $^b$  Determined by  $^1\mathrm{H}$  NMR analysis.

In conclusion, we have described the use of boronate complexes derived from 1,2-bis-boronic esters in photoredox-catalyzed radical transformations. This reaction demonstrates, for the first time, a radical 1,2-boron shift under thermodynamic control, allowing for counter-intuitive selective functionalization of the more hindered position of 1,2-bis-boronic esters. We have demonstrated a broad substrate scope and highlighted the diversity of the method in ring fragmentation and transannular reactions. Furthermore, the stereoretentive nature of the 1,2-boron shift was shown and we have showcased the application of this approach in different settings by extending it to radical cascades with allylboronic esters.

#### ASSOCIATED CONTENT

#### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures and characterization data for new compounds (PDF)

#### AUTHOR INFORMATION

#### Corresponding Author

\*v.aggarwal@bristol.ac.uk.

#### Notes

The authors declare no competing financial interests.

#### ACKNOWLEDGMENT

We thank H2020 ERC (670668) for financial support. D.K. thanks the Austrian Science Fund (FWF) for an Erwin Schrödinger Fellowship (J 4202-N28). V.F. thanks the EPSRC for a Doctoral Prize Fellowship. We thank Dr. Amadeu Bonet for useful discussions.

#### REFERENCES

(1) (a) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. Photoredox Catalysis in Organic Chemistry. *J. Org. Chem.* **2016**, *81*, 6898. (b) Narayanama, J. M. R.; Stephenson, C. R. J. Visible Light Photoredox Catalysis: Applications in Organic Synthesis. *Chem. Soc. Rev.* **2011**, *40*, 102. (c) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis. *Chem. Rev.* **2013**, *113*, 5322. (d) Goddard, J.-P.; Ollivier, C.; Fensterbank, L. Photoredox Catalysis for the Generation of Carbon Centered Radicals. *Acc. Chem. Res.* **2016**, *49*, 1924. (e) Staveness, D.; Bosque, I.; Stephenson, C. R. J. Free Radical Chemistry Enabled by Visible Light-Induced Electron Transfer. *Acc. Chem. Res.* **2016**, *49*, 2295. (f) Marzo, L.; Pagire, S. K.; Reiser, O.; König, B. Visible-Light Photocatalysis: Does It Make a Difference in Organic Synthesis? *Angew. Chem. Int. Ed.* **2018**, *57*, 10034.

(2) (a) Yasu, Y.; Koike, T.; Akita, M. Visible Light-Induced Selective Generation of Radicals from Organoborates by Photoredox Catalysis. Adv. Synth. Catal. 2012, 354, 3414. (b) Lima, F.; Sharma, U. K.; Grunenberg, L.; Saha, D.; Johannsen, S.; Sedelmeier, J.; Van der Eycken, E. V.; Ley, S. V. A Lewis Base Catalysis Approach for the Photoredox Activation of Boronic Acids and Esters. Angew. Chem. Int. Ed. 2017, 56, 15136. (c) Lima, F.; Grunenberg, L.; Rahman, H. B. A.; Labes, R.; Sedelmeier, J.; Ley, S. V. Organic Photocatalysis for the Radical Couplings of Boronic Acid Derivatives in Batch and Flow. Chem. Commun. 2018, 54, 5606. (d) Shu, C.; Noble, A.; Aggarwal, V. K. Photoredox-Catalyzed Cyclobutane Synthesis by a Deboronative Radical Addition-Polar Cyclization Cascade. Angew. Chem. Int. Ed. 2019, 58, 3870. (e) Lima, F.; Kabeshov, M. A.; Tran, D. N.; Battilocchio, C.; Sedelmeier, J.; Sedelmeier, G.; Schenkel, B.; Lev, S. V. Visible Light Activation of Boronic Esters Enables Efficient Photoredox C(sp<sup>2</sup>)-C(sp<sup>3</sup>) Cross-Couplings in Flow. Angew. Chem. Int. Ed. 2016, 55, 14085. (f) Clausen, F.; Kischkewitz, M.; Bergander, K.; Studer, A. Catalytic Protodeboronation of Pinacol Boronic Esters: Formal Anti-Markovnikov Hydromethylation of Alkenes. Chem. Sci. 2019, 10, 6210. (g) Duret, G.; Quinlan, R.; Bisseret, P.; Blanchard, N. Boron Chemistry in a New Light. Chem. Sci. 2015, 6, 5366.

(3) (a) Miyazawa, K.; Yasu, Y.; Koike, T.; Akita, M. Visible-Light-Induced Hydroalkoxymethylation of Electron-Deficient Alkenes by Photoredox Catalysis. Chem. Commun. 2013, 49, 7249. (b) Miyazawa, K.; Koike, T.; Akita, M. Hydroaminomethylation of Olefins with Aminomethyltrifluoroborate by Photoredox Catalysis. Adv. Synth. Catal. 2014, 356, 2749. (c) Chinzei, T.; Miyazawa, K.; Yasu, Y.; Koike, T.; Akita, M. Redox-Economical Radical Generation from Organoborates and Carboxylic Acids by Organic Photoredox Catalysis. RSC Adv. 2015, 5, 21297. (d) Huo, H.; Harms, K.; Meggers, E. Catalytic, Enantioselective Addition of Alkyl Radicals to Alkenes via Visible-Light-Activated Photoredox Catalysis with a Chiral Rhodium Complex. J. Am. Chem. Soc. 2016, 138, 6936. (e) Tellis, J. C.; Primer, D. N.; Molander, G. A. Single-Electron Transmetalation in Organoboron Cross-Coupling by Photoredox/Nickel Dual Catalysis. Science 2014, 345, 433. (f) Primer, D. N.; Karakaya, I.; Tellis, J. C.; Molander, G. A. Single-Electron Transmetalation: An Enabling Technology for Secondary Alkylboron Cross-Coupling. J. Am. Chem. Soc. 2015, 137, 2195. (g) Primer, D. N.; Molander, G. A. Enabling the Cross-Coupling of Tertiary Organoboron Nucleophiles through Radical-Mediated Alkyl Transfer. J. Am. Chem. Soc. 2017, 139, 9847. (h) Amani, J.; Molander, G. A. Synergistic Photoredox/Nickel Coupling of Acyl Chlorides with Secondary Alkyltrifluoroborates: Dialkyl Ketone Synthesis. J. Org. Chem. 2017, 82, 1856. (i) Huang, H.; Zhang, G.;
Gong, L.; Zhang, S.; Chen, Y. Visible-Light-Induced Chemoselective Deboronative Alkynylation under Biomolecule-Compatible Conditions. J. Am. Chem. Soc. 2014, 136, 2280. (j) Huang, H.; Jia, K.; Chen, Y. Hypervalent Iodine Reagents Enable Chemoselective Deboronative/Decarboxylative Alkenylation by Photoredox Catalysis. Angew. Chem. Int. Ed. 2015, 54, 1881. (k) Heitz, D. R.; Rizwan, K.; Molander, G. A. Visible-Light-Mediated Alkenylation, Allylation, and Cyanation of Potassium Alkyltrifluoroborates with Organic Photoredox Catalysts. J. Org. Chem. 2016, 81, 7308. (l) Lang, S. B.; Wiles, R. J.; Kelly, C. B.; Molander, G. A. Photoredox Generation of Carbon-Centered Radicals Enables the Construction of 1,1-Difluoroalkene Carbonyl Mimics. Angew. Chem. Int. Ed. 2017, 56, 15073.

1

2

3

4

5

6

7

8

9

10

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57 58 59

60

11 (4) (a) Bonet, A. Pubill-Ulldemolins, C.; Bo, C.; Gulyás, H.; Fernández, 12 E. Transition-Metal-Free Diboration Reaction by Activation of Diboron 13 Compounds with Simple Lewis Bases. Angew. Chem. Int. Ed. 2011, 50, 7158. (b) Miralles, N.; Cid, J.; Cuenca, A. B.; Carbó, J. J.; Fernández, E. 14 Mixed Diboration of Alkenes in a Metal-Free Context. Chem. Commun. 15 2015, 51, 1693. (c) Farre, A.; Soares, K.; Briggs, R. A.; Balanta, A.; Benoit, 16 D. M.; Bonet, A. Amine Catalysis for the Organocatalytic Diboration of 17 Challenging Alkenes. Chem. Eur. J. 2016, 22, 17552. (d) Farre, A.; Briggs, R.; Pubill-Ulldemolins, C.; Bonet, A. Developing a Bench-Scale Green Di-18 boration Reaction toward Industrial Application. Synthesis 2017, 49, 19 4775. (e) Bonet, A.; Sole, C.; Gulyás, H.; Fernández, E. Asymmetric Or-20 ganocatalytic Diboration of Alkenes. Org. Biomol. Chem. 2012, 10, 6621. (f) Blaisdell, T. P.; Caya, T. C.; Zhang, L.; Sanz-Marco, A.; Morken, J. P. 21 Hydroxyl-Directed Stereoselective Diboration of Alkenes. J. Am. Chem. 22 Soc. 2014, 136, 9264. (g) Fang, L.; Yan, L.; Haeffner, F.; Morken, J. P. Car-23 bohydrate-Catalyzed Enantioselective Alkene Diboration: Enhanced 24 Reactivity of 1,2-Bonded Diboron Complexes. J. Am. Chem. Soc. 2016, 138, 2508. (h) Yan, L. Meng, Y.; Haeffner, F.; Leon, R. M.; Crockett, M. P.; 25 Morken, J. P. Carbohydrate/DBU Cocatalyzed Alkene Diboration: Mech-26 anistic Insight Provides Enhanced Catalytic Efficiency and Substrate 27 Scope. J. Am. Chem. Soc. 2018, 140, 3663. (i) Morgan, J. B.; Miller, S. P.; Morken, J. P. Rhodium-Catalyzed Enantioselective Diboration of Simple 28 Alkenes. J. Am. Chem. Soc. 2003, 125, 8702. (j) Trudeau, S.; Morgan, J. B.; 29 Shrestha, M.; Morken, J. P. Rh-Catalyzed Enantioselective Diboration of 30 Simple Alkenes: Reaction Development and Substrate Scope. J. Org. 31 Chem. 2005, 70, 9538. (k) Kliman, L. T.; Mlynarski, S. N.; Morken, J. P. Pt-Catalyzed Enantioselective Diboration of Terminal Alkenes with 32 B2(pin)2. J. Am. Chem. Soc. 2009, 131, 13210. (l) Coombs, J. R.; Haeffner, 33 F.; Kliman, L. T.; Morken, J. P. Scope and Mechanism of the Pt-Catalyzed 34 Enantioselective Diboration of Monosubstituted Alkenes. J. Am. Chem. 35 Soc. 2013, 135, 11222. (m) Toribatake, K.; Nishiyama, H. Asymmetric Diboration of Terminal Alkenes with a Rhodium Catalyst and Subse-36 quent Oxidation: Enantioselective Synthesis of Optically Active 1,2-Di-37 ols. Angew. Chem. Int. Ed. 2013, 52, 11011. (n) Toribatake, K.; 38 Nishiyama, H. Miyata, S.; Naganawa, Y.; Nishiyama, H. Asymmetric Synthesis of Optically Active 3-Amino-1,2-Diols from N-Acyl-Protected Al-39 lylamines via Catalytic Diboration with Rh[bis(oxazolinyl)phenyl] Cat-40 alysts. Tetrahedron 2015, 71, 3203. 41

(5) Sandford, C.; Aggarwal, V. K. Stereospecific Functionalizations and Transformations of Secondary and Tertiary Boronic Esters. *Chem. Commun.* **2017**, *53*, 5481.

(6) (a) Miller, S. P.; Morgan, J. B.; Nepveux, F. J., V; Morken, J. P. Catalytic Asymmetric Carbohydroxylation of Alkenes by a Tandem Diboration/Suzuki Cross-Coupling/Oxidation Reaction. Org. Lett. 2004, 6, 131. (b) Lee, Y.; Jang, H.; Hoveyda, A. H. Vicinal Diboronates in High Enantiomeric Purity through Tandem Site-Selective NHC-Cu-Catalyzed Boron-Copper Additions to Terminal Alkynes. J. Am. Chem. Soc. 2009, 131, 18234. (c) Mlynarski, S. N.; Schuster, C. H.; Morken, J. P. Asymmetric Synthesis from Terminal Alkenes by Cascades of Diboration and Cross-Coupling. Nature 2014, 505, 386. (d) Fawcett, A.; Nitsch, D.; Ali, M.; Bateman, J. M.; Myers, E. L.; Aggarwal, V. K. Regio- and Stereoselective Homologation of 1,2-Bis(Boronic Esters): Stereocontrolled Synthesis of 1,3-Diols and Sch 725674. Angew. Chem. Int. Ed. 2016, 55, 14663. (e) Liu, X.; Sun, C.; Mlynarski, S.; Morken, J. P. Synthesis and Stereochemical Assignment of Arenolide. Org. Lett. 2018, 20, 1898.

(7) (a) Urry, W. H.; Kharasch, M. S. Factors Influencing the Course and Mechanism of Grignard Reactions. XV. The Reaction of β,β-Dimethylphenethyl Chloride with Phenylmagnesium Bromide in the Presence of Cobaltous Chloride. *J. Am. Chem. Soc.* **1944**, *66*, 1438. (b) Dowd, P.; Choi, S.C. A New Tri-*n*-butyltin Hydride Based Rearrangement of Bromomethyl β-Keto Esters. A Synthetically Useful Ring Expansion to  $\gamma$ -Keto Esters. *J. Am. Chem. Soc.* **1987**, *109*, 3493. (c) Athelstan L. J. Beckwith, A. L. J.; O'Shea, D. M.; Gerba, S.; Westwood, S. W. Cyano or Acyl Group Migration by Consecutive Homolytic Addition and β-Fission. *J. Chem. Soc., Chem. Commun.* **1987**, 666. For reviews, see: (d) Dowd, P.; Zhang W. Free Radical Mediated Ring Expansion and Related Annulations. *Chem. Rev.* **1993**, *93*, 2091. (e) Chen, Z.-M.; Zhang, X.-M.; Tu Y.-Q. Radical Aryl Migration Reactions and Synthetic Applications. *Chem. Soc. Rev.* **2015**, *44*, 5220. (f) Weng, W.-Z.; Zhang, B. Recent Advances in the Synthesis of β-Functionalized Ketones by Radical-Mediated 1,2-Rearrangement of Allylic Alcohols. *Chem. Eur. J.* **2018**, *24*, 10934.

(8) (a) Skell, P. S.; Allen, R. G.; Gilmour, N. D. Radical Rearrangements in Bromoalkyl Radicals. J. Am. Chem. Soc. **1961**, *81*, 504. (b) Chen, K. S.; Elson, I. H.; Kochi, J. K. Bridging in β-Chloroalkyl Radicals by Electron Spin Resonance. J. Am. Chem. Soc. **1973**, *95*, 5341. (c) Tan, E. W.; Chan, B.; Blackman, A. G. A Polar Effects Controlled Enantioselective 1,2-Chlorine Atom Migration via a Chlorine-Bridged Radical Intermediate. J. Am. Chem. Soc. **2002**, *124*, 2078. (d) Neumann, B.; Zipse, H. 1,2-Chlorine Atom Migration in 3-Chloro-2-butyl Radicals: A Computational Study. Org. Biomol. Chem. **2003**, *1*, 168. (e) Chen, H.-L.; Wei, D.; Zhang, J.-W.; Li, C.-L.; Yu, W.; Han, B. Synthesis of Halomethyl Isoxazoles/Cyclic Nitrones via Cascade Sequence: 1,2-Halogen Radical Shift as a Key Link. Org. Lett. **2018**, *20*, 2906.

(9) (a) Schiesser, C. H.; Styles, M. L. On the Radical Brook and Related Reactions: An Ab Initio Study of some (1,2)-Silyl, Germyl and Stannyl Translocations. *J. Chem. Soc., Perkin Trans. 2* **1997**, 2335. (b) Shuto, S.; Kanazaki, M.; Ichikawa, S.; Minakawa, N.; Matsuda, A. Stereo- and Regioselective Introduction of 1- or 2-Hydroxyethyl Group via Intramolecular Radical Cyclization Reaction with a Novel Silicon-Containing Tether. An Efficient Synthesis of 4' $\alpha$ -Branched 2'-Deoxyadenosines. *J. Org. Chem.* **1998**, 63, 746. (c) Sugimoto, I.; Shuto, S.; Matsuda, A. Kinetics of a Novel 1,2-Rearrangement Reaction of  $\beta$ -Silyl Radicals. The Ringexpansion of (3-Oxa-2-silacyclopentyl)methyl Radical into 4-Oxa-3-silacyclohexyl Radical Is Irreversible. *Synlett* **1999**, *10*, 1766. (d) Walton, J. C.; Kanada, R.; Takeaki, I.; Shuto, S.; Abe, H. Tethered 1,2-Si-Group Migrations in Radical-Mediated Ring Enlargements of Cyclic Alkoxysilanes: An EPR Spectroscopic and Computational Investigation. *J. Org. Chem.* **2017**, *82*, 6886.

(10) Lee, C. F.; Diaz, D. B.; Holownia, A.; Kaldas, S. J.; Liew, S. K.; Garrett, G. E.; Dudding, T.; Yudin, A. K. Amine Hemilability Enables Boron to Mechanistically Resemble either Hydride or Proton. *Nat. Chem.* **2018**, *10*, 1062.

(11) Tao, Z.; Robb, K. A.; Panger, J. L.; Denmark, S. E. Enantioselective, Lewis Base-Catalyzed Carbosulfenylation of Alkenylboronates by 1,2-Boronate Migration. *J. Am. Chem. Soc.* **2018**, *140*, 15621. Although an anionic pathway for boryl migration was proposed, it is possible that the reaction occurs through a radical process similar to that described in this paper.

(12) Batey, R. A.; Smil, D. V. The First Boron-Tethered Radical Cyclizations and Intramolecular Homolytic Substitutions at Boron. *Angew. Chem. Int. Ed.* **1999**, *38*, 1798.

(13) For the only examples of selective functionalization of the secondary position of 1,2-bis-boronic esters, see: (a) Blaisdell, T. P.; Morken, J. P. Hydroxyl-Directed Cross-Coupling: A Scalable Synthesis of Debromohamigeran E and Other Targets of Interest. *J. Am. Chem. Soc.* **2015**, *137*, 8712. (b) Yan, L.; Morken, J. P. Site-Selective Mono-Oxidation of 1,2-Bis(boronates). *Org. Lett.* **2019**, *21*, 3760. For an example of selective deboronative functionalisation of internal position of a 1,2,3triboronic ester, see: (c) Davenport, E.; Fernandez, E. Transition-Metal-Free Synthesis of Vicinal Triborated Compounds and Selective Functionalisation of the Internal C-B Bond. *Chem. Commun.* **2018**, *54*, 10104.

(14) See Supporting Information for details.

(15) Aryl boronic ester **7** (Ar = 2,4,6-triisopropylphenyl) was isolated in 82% yield from the reaction of **1d** with *tert*-butyl acrylate.

(16) Villa, G.; Povie, G.; Renaud, P. Radical Chain Reduction of Alkylboron Compounds with Catechols. J. Am. Chem. Soc. **2011**, *133*, 5913.

(17) Performing the reaction under more dilute conditions (0.02 M) led to an enhanced ratio (15:16 = 4.1:1). However, the overall efficiency suffered and 15 was isolated in a similar yield of 53%.

(18) Wilger, D. J.; Gesmundo, N. J.; Nicewicz, D. A. Catalytic Hydrotri-

fluoromethylation of Styrenes and Unactivated Aliphatic Alkenes via an

Organic Photoredox System. Chem. Sci. 2013, 4, 3160.

ACS Paragon Plus Environment

(19) Dénès, F.; Pichowicz, M.; Povie, G.; Renaud, P. Thiyl Radicals in Organic Synthesis. Chem. Rev. 2014, 114, 2587.

