

α -Amino acid Tröger base derivatives, possible conformationally restricted scaffolds?^{†‡}

Sean P. Bew,* Laurent Legentil, Vincent Scholier and Sunil V. Sharma

Received (in Cambridge, UK) 4th October 2006, Accepted 7th December 2006

First published as an Advance Article on the web 15th December 2006

DOI: 10.1039/b614371g

The first synthesis of innovative α -amino acid conjugates of Tröger base is reported; their potential application as conformationally restricted scaffolds is proposed and has been investigated using high level *ab initio* calculations.

We report studies on 2,8-*bis*- α -amino acid Tröger base adducts (**1**, Fig. 1) and propose them as scaffolds for use as peptide chain directors *i.e.* **2**. Tröger base¹ is a C_2 -symmetric heterocycle with a relatively rigid backbone, hydrophobic cavity and concave conformation. Due to its sharply folded geometry the aryl rings reside in a near perpendicular arrangement (generally 90–100°).²

Gaining insights into the biological mode of action of natural/non-natural proteins, polypeptides and enzymes is critical if a comprehensive understanding of protein action is to be acquired. In this respect many studies have been undertaken on the application/development of innovative mimics of β -turns and hairpins.³ The synthesis of an α -amino acid derived Tröger base scaffold⁴ and its appendage with additional α -amino acid derivatives affords a new opportunity to investigate the loop and hinge regions of proteins. α -Amino acid Tröger base derivatives similar to **3** (Fig. 2) appear not only to be able to act as scaffolds, but also to have the capacity to direct appended peptides within a 90–100° range. Small peptide chains have been identified that contain 90–100° bends. For example, the polypeptide TNYLFSPNGPIARAW that binds to EphB4 (IC₅₀ 15 nM) embodies a 90° turn induced by the GP dipeptide. This turn within the pentadecapeptide is *critical* for high affinity binding of the polypeptide into the hydrophobic upper convex portion of the active site within EphB4.⁵

The transition-metal mediated synthesis of adducts based on **3** requires a practical, efficient and convenient synthesis of building block **4** (Scheme 1).⁶ The *bis*-2,8-dibromo analog of **4** was not considered due to its recalcitrant nature towards Sonogashira




Fig. 2 Chem 3D representation of *bis*-2,8-(*N,C*-protected α -amino acid) Tröger base adduct **3**.

couplings.⁸ However, the synthesis of milligram quantities of racemic *bis*-2,8-diiodo **4** has been reported by Wärmark *et al.*⁷ Translating this protocol directly to the production of multigram quantities of **4** was problematic with low yields (~10%) resulting. After careful optimisation of the experimental and purification procedures we were able to accrue multigram quantities of racemic **4** in ~46% yield. Employing a single enantiomer of **4** would significantly simplify the analysis of the resulting diastereomeric α -amino acid Tröger base conjugates. That said, all our attempts at resolving **4** using literature protocols failed.⁸ Undeterred, we transformed racemic **4** into *bis*-2,8-dialkyne **5** via a Sonogashira coupling with TMSA. Desilylation of **5** using TBAF afforded **6** in a poor 23% yield. Switching to sodium hydroxide in a methanol–THF mix negated this problem; an excellent 91% yield of *bis*-2,8-ethynyl **6** resulted. Coupling **6** with 4-iodo-*N*-Boc-(*S*)-phenylalanine methyl ester **7** was attempted. Utilising standard Sonogashira coupling conditions either no reaction took place or poor yields of **9** resulted (~15%). Gratifyingly, when DMF was employed, and freshly synthesised *tetrakis*(triphenylphosphine)palladium(0) and 2.2 equiv. of **7** were utilised the desired Tröger base adduct **9** was returned in an unoptimised 60% yield (Scheme 1).





Fig. 1 α -Amino acid Tröger base scaffolds as peptide chain directors.

School of Chemical Sciences & Pharmacy, University of East Anglia, Norwich, Norfolk, UK NR4 7TJ. E-mail: s.bew@uea.ac.uk;

Fax: 44 (0)1603 592003; Tel: 44(0)1603 593142

[†] Electronic supplementary information (ESI) available: Experimental details. See DOI: 10.1039/b614371g

[‡] The HTML version of this article has been enhanced with colour images.



Scheme 1 Synthesis of 2,8-difunctionalised Tröger base adducts.

Curiously, attempting to couple **4** and *N*-Boc-4-ethynyl-*(S)*-phenylalanine methyl ester (**8**) using the previously successful Sonogashira reaction conditions for **6** and **7** *i.e.* DMF, Et₃N, CuI, Pd(PPh₃)₄, failed to return any **9**.

We considered the possibility that the transition-metal mediated coupling reaction between *(S)*-**7** and racemic **4** may result in a chiral resolution of racemic **4** affording diastereomerically enriched **9**. Subjecting **9** to chiral HPLC analysis (Chiralpack AD, 24 × 0.46 cm) clearly showed the *(S,S,S)*- and *(S,R,S)*-**9** adducts in an equal ratio.⁹ Our attempts at separating, using flash chromatography, *(S,S,S)*- and *(S,R,S)*-**9** failed. Similar separation problems with different Tröger base adducts have been reported by Maitra *et al.*¹⁰

The synthesis of an alicyclic, *bis*-2,8-*(N,C)*-protected- α -amino acid Tröger base was attempted. Incorporating **4**, *N*-Boc-*(S)*-propargylglycine ethyl ester (2.5 equivalents) and the reaction conditions/catalysts employed to couple **6** and **7** (Scheme 1), we were surprised to find that from the myriad of by- and decomposition products the desired adduct **10** (Fig. 3) was afforded in a disappointing 10% yield; furthermore we also isolated, albeit in mediocre 28% yield, the corresponding 2-*(N*-Boc-*(S)*-propargylglycine ethyl ester)-8-iodo Tröger base.

Hydrogenation (5% Pd on C, H₂, MeOH, 1 atmosphere) of the rigid *bis*-2,8-ethynyl linkers within **9** and **10** afforded, in quantitative yields, the corresponding alkane linked adducts **12** and **11** respectively. The reported sensitivity of Tröger base to acidic conditions^{2a} compelled us to check the feasibility of performing an acid mediated *N*-Boc deprotection. Gratifyingly, reacting **11** or **12** with TFA cleanly removed all the *N*-Boc groups (79% and 82% yields respectively) affording the corresponding TFA salts. Liberation of the free amine (Et₃N) from the TFA salt of *N*-deprotected **12** followed by ¹H-NMR analysis confirmed the Tröger base heterocycle to be intact. Testing the concept that additional α -amino acids can be readily appended, tetrapeptide **13** was synthesised in an excellent 89% yield (Scheme 2).

As part of our wider strategy towards utilising Tröger base as a scaffold we sought to append *two* differentially *N,C*-protected- α -amino acids onto *one* Tröger base. The possibility of chemoselectively cleaving, when desired, one of the four *N*- or *C*-protecting groups off one of the α -amino acids would significantly enhance the potential of the Tröger base scaffolds.

Disappointingly, utilising **6**, *one* equivalent of **7** and the catalyst/reaction conditions outlined in Scheme 1 failed to afford any significant amounts of the desired *mono*- α -amino acid appended adduct. A complex mixture of products comprising: starting material **6**, homocoupled *bis*-alkynyls, *bis*-2,8-alkynyl Tröger base adduct *i.e.* **9** as well as unidentifiable by-products resulted.

Buchwald *et al.*¹¹ and Wärnmark *et al.*⁷ have reported on an unusually reactive Sonogashira protocol for the efficient coupling of electron-rich aryl halides and alkynes that employs catalytic amounts of tri-*tert*-butylphosphine (10%). Utilising Wärnmark's






Fig. 3 *Bis*-2,8-*[N*-Boc-*(S)*-propargylglycine ethyl ester] Tröger base.



Scheme 2 Synthesis of Tröger base **13**.

procedure, **4** and *N*-Boc-*(S)*-propargylglycine ethyl ester (1 equivalent each) we isolated the desired *mono* α -amino acid coupled 8-iodo-Tröger base adduct in an unoptimised 56% yield. Subjecting this to a second Sonogashira coupling with TMSA [PdCl₂(PPh₃)₂, Et₃N, CuI, THF] and subsequent desilylation afforded **15** which was readily coupled to **14** affording the fully differentially *N,C*-protected α -amino acid derived Tröger base **16** (Scheme 3). With this important adduct in hand its chemoselective TMSE *C*-deprotection was undertaken using TBAF; the resulting carboxylic acid was coupled to *(S)*-valine methyl ester affording



Scheme 3 Synthesis of Tröger base derived tetrapeptides **18** and **19**.




Fig. 4 Calculated (using B3LYP/6-31G) conformation of **11** and *tert*-butyl Tröger base edge-to-face interactions reported by Wilcox *et al.*




Fig. 5 Calculated conformation using B3LYP/6-31G basis set.

tripeptide **17** in a 60% yield. Chemoselective cleavage of the *N*-Boc group off **17** (TFA, 83%) and subsequent appendage of (*N*)-Boc-(*S*)-histidine afforded Tröger base tetrapeptide **18**, which underwent hydrogenation affording **19**.

With the potential application of the α -amino acid Tröger base conjugates as scaffolds in mind we sought corroboration of their conformation. Despite an intensive effort we have been unable to grow crystals of **11–13** or **19** suitable for X-ray analysis, furthermore NOE experiments were unproductive.

Employing *ab initio* DFT calculations Stephens *et al.*¹² predicted the conformations of a series of Tröger base adducts and compared their results with X-ray crystal structures deposited in the CSD. The agreement of theory and experiment was excellent. Using Gaussian 98¹³ (B3LYP functional level and 6-31G basis set) an energy minimisation performed on the relatively simple adduct **11** revealed an unusual 'bite back' of one of the *N*-Boc groups such that the *tert*-butyl moiety undergoes edge-to-face aryl-alkyl interactions (Fig. 4). Interestingly, and in agreement with our calculations, Wilcox *et al.* observed similar molecular recognition forces in 'Tilted-T' Tröger base derivatives. Indeed Wilcox *et al.* reported that alkyl *tert*-butyl ester **20** had a 'strong preference, greater than any aryl ester, for the formation of edge-to-face interactions'.¹⁴

To test our concept that α -amino acid Tröger base conjugates may be capable of acting as conformationally restrictive scaffolds we subjected *N*-Cbz derived **19** to *ab initio* DFT calculations (B3LYP and 6-31G). Gratifyingly, the calculations indicate (Fig. 5) that the Tröger base scaffold does indeed constrain appended

α -amino acids, holding them to a near perpendicular angle (calculated aryl plane angle of 100.83°). Furthermore, and of importance for application as a scaffold, the DFT calculation suggests that the *N*- and *C*-terminated α -amino acid 'arms' are projected along the aryl planes of the Tröger base.

In summary, we have developed a novel strategy for the synthesis of structurally unique *bis*-(*N,C*-protected- α -amino acid) derived Tröger base adducts. Using chemoselective *N*- or *C*-amino acid deprotection strategies we have demonstrated the feasibility of synthesising non-symmetric Tröger base tetrapeptides. Our hypothesis that Tröger base adducts can be employed as new conformationally restricted scaffolds has been reinforced by high level *ab initio* calculations.

SPB would like to thank Pfizer, The University of East Anglia (VS), the Dovetrust (SVS), and the EPSRC for funding (LL) and Professor Floris Rutjers (University of Nijmegen) for a gift of (*S*)-propargylglycine.

Notes and references

- 1 M. Valík, R. M. Strongin and V. Král, *Supramol. Chem.*, 2005, **17**, 347.
- 2 M. Demeunynck and A. Tatibouët, *Prog. Heterocycl. Chem.*, 1999, **11**, 1; I. Sucholeiki, V. Lynch, L. Phan and C. S. Wilcox, *J. Org. Chem.*, 1988, **53**, 98.
- 3 N. Sewald and H.-D. Jakubke, *Peptides: Chemistry and Biology*, Wiley-VCH, Weinheim, 2002.
- 4 J. E. Chrencik, A. Brooun, M. I. Recht, M. L. Kraus, M. Koolpe, A. R. Kolatkar, R. H. Bruce, G. Martiny-Baron, H. Widmer, E. B. Pasquale and P. Kuhn, *Structure*, 2006, **14**, 321.
- 5 K. D. Stigers, M. J. Soth and J. S. Nowick, *Curr. Opin. Chem. Biol.*, 1999, **3**, 714; Y. J. Chung, B. R. Huck, L. A. Christianson, H. E. Stanger, S. Krauthauser, D. R. Powell and S. H. Gellman, *J. Am. Chem. Soc.*, 2000, **122**, 3995; S. Goswami, K. Ghosh and S. Dasgupta, *J. Org. Chem.*, 2000, **65**, 1907; C. Pardo, E. Sesnilo and E. Gutierrez-Puebla, *J. Org. Chem.*, 2001, **66**, 1607; S. Peluso, P. Dumy, L. Yokokana and M. Mutter, *J. Org. Chem.*, 1999, **64**, 7114.
- 6 F. Hof, M. Schar, D. M. Scofield, F. Fischer, F. Diederich and S. Sergeyev, *Helv. Chim. Acta*, 2005, **88**, 2333; C. Solano, D. Svensson, Z. Olomi, J. Jensen, O. F. Wendt and K. Wärnmark, *Eur. J. Org. Chem.*, 2005, **16**, 3510.
- 7 J. Jensen, M. Strozyk and K. Wärnmark, *Synthesis*, 2002, 2761.
- 8 S. H. Wilen, J. Z. Qi and P. G. Williard, *J. Org. Chem.*, 1991, **56**, 485.
- 9 See the ESI for the synthesis and analytical data associated with the compounds discussed within this manuscript.
- 10 U. Maitra and B. G. Bag, *J. Org. Chem.*, 1992, **57**, 6979.
- 11 T. Hundertmark, A. F. Littke, S. L. Buchwald and G. C. Fu, *Org. Lett.*, 2000, **2**, 1729.
- 12 A. Aamouche, F. J. Devlin and P. J. Stephens, *J. Am. Chem. Soc.*, 2000, **122**, 2346.
- 13 Gaussian 98; M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Montgomery, Jr., R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, C. Gonzalez, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, J. L. Andres, M. Head-Gordon, E. S. Replogle and J. A. Pople, Gaussian, Inc.: Pittsburgh, PA, 1998.
- 14 S. Paliwal, S. Geib and C. S. Wilcox, *J. Am. Chem. Soc.*, 1994, **116**, 4497; E.-I. Kim, S. Paliwal and C. S. Wilcox, *J. Am. Chem. Soc.*, 1998, **120**, 11192.