

Total Synthesis of (+)-Complanadine A Using an Iridium-Catalyzed Pyridine C–H Functionalization

Daniel F. Fischer and Richmond Sarpong*

Department of Chemistry, University of California, Berkeley, California 94720


Received March 5, 2010; E-mail: rsarpong@berkeley.edu

The *Lycopodium* alkaloids are among the most skeletally diverse families of natural products that have been isolated to date.¹ In addition to their complex architecture, many of the alkaloids in this family possess potentially useful biological activity, which includes acetylcholinesterase (AChE) inhibition.² The utility of known AChE inhibitors, such as the *Lycopodium* alkaloid huperzine A (**1**, Figure 1),³ to partially address the treatment of Alzheimer's disease is supported by the observation of increased memory and learning in rats upon administration of **1**.⁴ Furthermore, several AChE inhibitors, including galanthamine⁵ and Aricept,⁶ are commercial pharmaceuticals that are used to treat Alzheimer's disease indications. However, because AChE inhibition appears to treat only the symptoms of neurodegenerative diseases, there continues to be a need to identify other natural products that promote the growth of new neural networks. In this regard, *Lycopodium* alkaloids, such as the lyconadins (e.g., **3**)⁷ and complanadine A (**4**),⁸ are especially interesting because they enhance mRNA expression for nerve growth factor (NGF) and the production of NGF in human glial cells.⁹


As a first step in a program aimed at the biological evaluation of *Lycopodium* alkaloids and their analogues as neurotrophic factors, we have undertaken the synthesis of several of these compounds. We recently reported the total synthesis of lyconadin A (**3**).¹⁰ In this manuscript, we present the details of our total synthesis of complanadine A (**4**), which is a dimer of the well-known phlegmarine-derived alkaloid lycodine (**5**).¹¹ The synthetic challenge posed by **4** stems primarily from the fact that it is an *unsymmetrical* dimer of lycodine (C2–C3' linkage; see **4**) and therefore requires the introduction of position-control elements prior to merging of the two halves (see **6** and **7**, Scheme 1). A powerful simplification of the synthesis would entail formation of **6** and **7** from the same intermediate (e.g., tricyclic enamide **8**). Enamide **8** was in turn envisioned to arise from an acid-promoted formal cycloaddition of **9** and **10** following the precedent of Schumann for the synthesis of racemic *N*-desmethyl- α -obscurine (**17**, Scheme 2).

The synthesis of complanadine A commenced with the preparation of ketal **12** (eq 1), which is available from (+)-pulegone using

Scheme 1




literature procedures.¹³ Aminoketal **12** was used as a direct precursor to oxygen-sensitive α,β -unsaturated imine **10**.¹⁴




Enamide **9** was prepared from commercially available 5-keto-hexanenitrile¹⁵ via Zn(II)-catalyzed hydrolysis of the cyano group and subsequent dehydrative cyclization upon distillation.¹⁶ Interestingly, our empirical observations suggest that under the conditions identified by Schumann for the formation of **17** (Scheme 2) from **9** and **10** (70% HClO₄(aq), dioxane, 105 °C, 20 h), enamide **9** undergoes hydrolysis to **13**.¹⁷ Ketoamide **13** likely forms reactive enol tautomer **14**, which in turn reacts with protonated **10** (generated in situ from **12**) to yield *N*-desmethyl α -obscurine **17** following a series of cyclizations (i.e., via **15** and **16**).

Scheme 2




The synthetic sequence up to this point is easily amenable to scale-up, which has provided **17** on a multigram scale. At this stage, we proceeded to address the major challenge of the synthesis, which was to advance **17** to triflate **7** as well as boronic ester **6**. To set the stage for these tasks, selective Boc protection of **17** (Scheme 3) afforded **8** (65% yield over two steps).¹⁸ This was followed by oxidation of **8** with lead tetraacetate to provide pyridone **18** in 84% yield. Triflation of the pyridone using triflic anhydride in pyridine proved optimal, giving **7** in 72% yield.

Figure 1. Selected *Lycopodium* alkaloids.


With one of the desired coupling partners in hand, we looked to address the synthesis of the other (i.e., boronic ester **6**).¹⁹ The triflate group in **7** was removed under Pd-catalyzed reducing conditions to provide Boc-protected lycodine (**19**).

Scheme 3



The conversion of **19** to boronic ester **6** was achieved in a single step, as illustrated in Scheme 4. Our approach to the remarkable direct C–H functionalization of **19** rested on the uniquely effective Ir(I)-catalyzed borylation chemistry developed by Hartwig and Miyaura.²⁰ In the event, treatment of **19** with 4 mol % $[\text{Ir}(\text{OMe})(\text{cod})]_2$, 8 mol % di-*tert*-butylbipyridine (dtbpy), and diboron pinacolato ester (0.75 equiv) in THF at 80 °C gave **6** in 75% yield after 5.5 h. The observed site selectivity is consistent with that noted by Hartwig for pyridine functionalization and appears to be guided mainly by steric factors. Suzuki cross-coupling of boronic ester **6** and triflate **7** followed by cleavage of the Boc protecting groups gave complanadine A (**4**) in 42% yield over two steps.


Scheme 4



Synthetic complanadine A gave spectral data (^1H and ^{13}C NMR) consistent with that reported by Kobayashi and co-workers during its isolation.²¹

Access to boronic ester **6** should prove highly significant as a starting point for the synthesis of congeners and analogues of complanadine A and lycodine. For example, coupling of boronic ester **6** (Scheme 5) to vinyl bromide **20**²² followed by dihydroxylation using the Upjohn method²³ and periodate cleavage yields **22**, which is a direct precursor to lycopladienes G and F.²⁴

Scheme 5



In summary, we have reported the total synthesis of the unsymmetrical lycodine dimer complanadine A. Our synthetic sequence proceeds in eight steps from enamide **9** and acetal **12**. The synthetic strategy also provides potential access to the natural products lycodine, lycopladiene F, and lycopladiene G. Highlights of the complanadine synthesis include a Hartwig–Miyaura Ir(I)-catalyzed site-selective borylation and a late-stage Suzuki cross-coupling to form the C2–C3' bipyridine. The application of the strategy to the multigram scale production of **4** and congeners, as well as the synthesis of other *Lycopodium* alkaloids, is underway and will be reported in due course.

Acknowledgment. The authors are grateful to the NIGMS (RO1 GM086374-01), Eli Lilly, Johnson and Johnson, GlaxoSmithKline, Amgen, and AstraZeneca for financial support. D.F. is thankful for a DAAD postdoctoral fellowship.

Supporting Information Available: Experimental details and characterization data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For pertinent reviews, see: (a) Ma, X.; Gang, D. R. *Nat. Prod. Rep.* **2004**, *21*, 752. (b) Ayer, W. A.; Trofinov, L. S. *Lycopodium Alkaloids*; Academic Press: San Diego, CA, 1994. (c) Hudlicky, T.; Reed, J. W. *In The Way of Synthesis*, 1st ed.; Wiley-VCH: Weinheim, Germany, 2007; pp 573–602.
- (2) (a) Tang, X. C.; Han, Y. F.; Chen, X. P.; Zhu, X. D. *Acta Pharmacol. Sin.* **1986**, *7*, 507. (b) Tang, X. C.; De Sarno, P.; Sugaya, K.; Giacobini, E. *J. Neurosci. Res.* **1989**, *24*, 276.
- (3) Liang, Y. Q.; Tang, X. C. *Neurosci. Lett.* **2004**, *361*, 56.
- (4) Jiang, H. L.; Luo, X. M.; Bai, D. L. *Curr. Med. Chem.* **2003**, *10*, 2231.
- (5) Woodruff-Pak, D. S.; Vogel, R. W.; Wenk, G. L. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 2089.
- (6) For a leading review, see: Shigeta, M.; Homma, A. *CNS Drug Rev.* **2001**, *7*, 353.
- (7) For the isolation of **3**, see: Kobayashi, J.; Hirasawa, Y.; Yoshida, N.; Morita, H. *J. Org. Chem.* **2001**, *66*, 5901.
- (8) For the isolation of **4**, see: Kobayashi, J.; Hirasawa, Y.; Yoshida, N.; Morita, H. *Tetrahedron Lett.* **2000**, *41*, 9069.
- (9) (a) Ishiuchi, K.; Kubota, T.; Hoshino, T.; Obara, Y.; Nakahata, N.; Kobayashi, J. *Bioorg. Med. Chem.* **2006**, *14*, 5995. (b) Morita, H.; Ishiuchi, K.; Haganuma, A.; Hoshino, T.; Obara, Y.; Nakahata, N.; Kobayashi, J. *Tetrahedron* **2005**, *61*, 1955.
- (10) (a) Bisai, A.; West, S. P.; Sarpong, R. *J. Am. Chem. Soc.* **2008**, *130*, 7222. (b) West, S. P.; Bisai, A.; Lim, A. D.; Narayan, R. R.; Sarpong, R. *J. Am. Chem. Soc.* **2009**, *131*, 11187.
- (11) The group of Prof. D. Siegel (UT Austin) has also recently completed a synthesis of **4** (personal communication).
- (12) Schumann, D.; Naumann, A. *Liebigs Ann. Chem.* **1983**, 220.
- (13) (a) Schuster, E.; Jas, G.; Schumann, D. *Org. Prep. Proced. Int.* **1992**, *24*, 670. (b) Liu, K.-M.; Chau, C.-M.; Sha, C.-K. *Chem. Commun.* **2008**, 91. For more details, see the Supporting Information.
- (14) For ease of handling, air-sensitive ene-imine **10** was generated in situ under the reaction conditions for formation of **17**.
- (15) This reagent may also be prepared according to: Näslund, G.; Senning, A.; Lawesson, S.-O. *Acta Chem. Scand.* **1962**, *16*, 1324. For an improved procedure, see the Supporting Information.
- (16) This is an adaptation of a reported procedure: Kopylovich, M. N.; Kukushkin, V. Y.; Haukka, M.; Frausto da Silva, J. J. R.; Pombiero, A. J. L. *Inorg. Chem.* **2002**, *41*, 4798. For details, see the Supporting Information.
- (17) Preliminary results indicated that **13** may be used instead of **9** in the formation of **17**.
- (18) Because of its high polarity, **17** was taken directly into the next step without purification.
- (19) Although pyridone **18** may be brominated at the requisite position, attempts to advance the 3-bromopyridone were unsuccessful.
- (20) (a) Ishiyama, T.; Takagi, J.; Ishida, K.; Miyaura, N.; Anastasi, N. R.; Hartwig, J. F. *J. Am. Chem. Soc.* **2002**, *124*, 390. (b) For a related reaction, see: Cho, J.-Y.; Tse, M. K.; Holmes, D.; Maleczka, R. E., Jr.; Smith, M. R. *Science* **2002**, *295*, 305.
- (21) We are grateful to Prof. H. Morita (Hoshi University, Japan) for ^1H and ^{13}C NMR spectra of **4**. (b) Complanadine A was obtained along with small amounts of *N*-Boc-lycodine (**19**), which presumably arises from protodeborylation.
- (22) Mori, M.; Washioaka, Y.; Urayama, T.; Yoshiura, K.; Chiba, K.; Ban, Y. *J. Org. Chem.* **1983**, *48*, 4058.
- (23) (a) Schneider, W. P.; McIntosh, A. V. (Upjohn). U.S. Patent 2,769,824, 1956. (b) VanRheenen, V.; Kelly, R. C.; Cha, D. Y. *Tetrahedron Lett.* **1976**, *17*, 1973.
- (24) Ishiuchi, K.; Kubota, T.; Hayashi, S.; Shibata, T.; Kobayashi, J. *Tetrahedron Lett.* **2009**, *50*, 4221.

JA101893B