Formation of aluminacyclobutenes *via* carbon monoxide and isocyanide insertion

Xiaofei Li, Chengbao Ni, Haibing Song and Chunming Cui*

Received (in Cambridge, UK) 23rd January 2006, Accepted 3rd March 2006 First published as an Advance Article on the web 15th March 2006 DOI: 10.1039/b601056c

Reaction of $LAl[\eta^2-(CSiMe_3)_2]$ (L = HC[(CMe)(NAr)]_2, Ar = 2,6-*i*Pr₂C₆H₃) with carbon monoxide and *tert*-butyl isocyanide afforded unique AlC₃ aluminacyclobutenes *via* insertion into one of the aluminium–carbon bonds.

There is increasing interest in strained ring compounds that incorporate heavier main group elements because of their unique structures and high reactivity.¹ In this context, we are interested in the strained aluminium ring systems stabilized by sterically demanding ligands. Recently, several three-membered AlC₂ ring compounds have been reported.² It has been demonstrated that the AlC₂ ring compound, LAl[η^2 -(CSiMe₃)₂] (1, L = HC[(CMe)- $(NAr)_{2}$, Ar = 2,6-*i*Pr₂C₆H₃), can react with a range of small molecules such as carbon dioxide, ketones, nitriles, organic azides, dioxygen and CS₂, leading to either ring expansion or elimination of bis(trimethylacetylene).^{2b,3} These studies prompted us to explore the reaction of 1 with carbon monoxide and isocyanides since they are versatile synthons in organic and organometallic synthesis. Although the insertion of CO and isocyanides into transition metal-carbon bonds has been well established,⁴ similar studies on the insertion into aluminium-carbon bonds are much less common.⁵ To the best of our knowledge, there are only two reports concerning the insertion of CO and an isocyanide into aluminium-carbon bonds, namely the insertion of CO into the aluminium-carbon bond of tri-tert-butylaluminium⁶ and the double insertion of tert-butyl isocyanide into the Al-C bond of $Cp'_{3}Al (Cp' = C_{5}Me_{4}H)$.⁷ The facile insertion of CO and *tert*butyl isocyanide in these two cases may result from the threecoordinate environment of the aluminium centers. Herein we report on the syntheses and characterization of the AlC₃ aluminacyclobutene rings generated from the insertion reactions of CO and tert-butyl isocyanide with 1.

When a red black solution of **1** was exposed to a purified CO atmosphere (1 atm, moisture and oxygen should be strictly excluded) in *n*-hexane at room temperature, the colour change of the solution to yellow was observed within 2 min. Compound **2** was obtained in high yield after additional stirring for 20 min followed by a standard workup (Scheme 1). The reaction of **1** with *tert*-butyl isocyanide in *n*-hexane proceeds rapidly at -78 °C to give orange crystals of **3** after crystallization from diethyl ether. Compounds **2** and **3** have been characterized by ¹H and ¹³C NMR spectroscopy, IR spectroscopy and elemental analysis.† The ¹H NMR spectrum of **2** displays four doublets for the CH*Me*₂ groups on the Ar rings because of the non-symmetric AlC₃ ring. The ¹³C



Scheme 1 Reagents and conditions: i. CO, *n*-hexane, r.t.; ii. CNBu^t, *n*-hexane, -78 °C; iii. O₂, diethyl ether, -78 °C.

NMR spectrum of **2** shows a singlet at δ 234.8 ppm, attributed to the resonance of Al-C(O), which is downfield shifted from those of ketones (197-220 ppm)⁸ but significantly upfield from that reported for the bridging acyl group of $[t-Bu_2AlC(O)Bu-t]_2$,⁶ indicating terminal acyl coordination9 and the existence of an α,β -unsaturated acyl unit.^{2b} The infrared spectrum of **2** exhibits a band at 1677 cm⁻¹ as expected for a terminal acyl moiety.⁹ Unfortunately we were unable to obtain crystals suitable for single crystal X-ray analysis for further confirmation because of its extremely high sensitivity to the air and moisture although many attempts have been made. Interestingly, reaction of 2 with dioxygen at -78 °C in diethyl ether results in the selective insertion of one oxygen atom into the Al-C(O) bond with the formation of LAI[OC(O)C(SiMe₃)(CSiMe₃)] (4), the same product as reported for the reaction of 1 with CO_2 .^{2b} The formation of 4 has been confirmed by its ¹H NMR spectrum and single crystal X-ray analysis.[‡] 4 crystallises in a monoclinic space group, different from that reported previously, but the structural parameters are only marginally different from those reported.^{3a}

The NMR spectrum of **3** indicates the formation of two isomers because of the two possible orientations of the *tert*-butyl group (Scheme 1). The two isomers are stable, and cannot be interconverted in hot toluene. It should be noted that the ratio (*ca.* 2 : 1 estimated from the ¹H NMR spectrum) of the two

State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, P. R. China. E-mail: cmcui@nankai.edu.cn; Fax: +86-22-23503461; Tel: +86-22-23503461

isomers is independent on the reaction temperatures (-78 °C to room temperature). Attempts to separate 3a and 3b by repeated crystallisation failed because of their very similar physical properties (solubility and appearance). However, the ¹H NMR spectrum of the mixture could be partially distinguished despite the overlaps of the resonances of the aromatic protons and CHMe2 groups of the two molecules. The proton NMR spectrum of the major isomer shows three singlets at δ 0.48, 0.55 and 0.80 ppm with the same integration, attributed to the resonances of the two SiMe₃ groups and the *t*-butyl group while the spectrum of the minor one shows resonances at δ -0.14, 0.52 and 1.42 ppm. The down and upfield shifts of one of the SiMe₃ groups and the *t*-butyl group in the ¹H NMR spectrum of the major isomer may result from intramolecular H...H interactions among the two groups. The chemical shifts for the other protons and the resonances found in the ¹³C NMR spectrum of the two isomers are quite close to each other. Based on these observations, we, tentatively, assign 3a as the major isomer because of the existence of steric repulsion between the t-butyl and its neighboring SiMe3 group. The iminoacyl resonances in the ¹³C NMR spectrum of **3** appear at δ 197.5 and 194.6 ppm, which are in the reported range for those found in terminal iminoacyl complexes.9 The molecular structure of one isomer has been characterized by single crystal X-ray analysis, which disclosed that the molecular array corresponds to 3b.‡

The structure of **3b** is shown in Fig. 1. The most striking feature of **3b** is the almost planar AlC_3 ring (mean deviation from the plane: 0.0236 Å), which is arranged nearly perpendicular to the N1–Al1–N2 plane (the angle between the C30–Al1–C32 and N1–Al1–N2 planes: 91.6°). The aluminium atom adopts a distorted tetrahedral geometry because of the geometric constraints of the

Fig. 1 Thermal ellipsoid drawing of **3b** (30% probability). Hydrogen atoms and *i*Pr groups on the Ar rings have been omitted. Selected bond distances (Å) and angles (°): Al1–C30 1.9842(19), Al1–C32 1.989(2), Al1–N1 1.9020(16), Al1–N2 1.9141(15), C30–C31 1.519(2), C31–C32 1.374(3), N3–C30 1.285(2), N3–C39 1.480(2); N1–Al1–N2 95.88(7), C30–Al1–C32 72.94(8), N1–Al1–C30 125.36(7), N2–Al1–C32 117.81(7), C30–C31–C32 109.31(16), C30–N3–C39 119.71(16).

two fused rings around the aluminium atom. The iminoacyl group is coordinated to the aluminium atom in an η¹ fashion and the Al1–C30 bond length (1.9842(19) Å) is in line with a Al–C (sp² hybrid) single bond.^{2b} The N3–C30 bond length (1.285(2) Å) is in the range for iminoacyl groups. The C30–C31 bond length (1.519(2) Å) is consistent with a C–C single bond, and the C31– C32 bond length (1.374(3) Å) indicates double bonding. The *trans* α,β-unsaturated iminoacyl moiety N3–C30–C31–C32 is not planar (torsion angle: 10.8°) probably due to the sterically demanding ligands on the AlC₃ four-membered ring.

In summary, the facile insertion of CO and isocyanide into an Al–C bond of the four coordinate aluminium AlC_2 ring is remarkable, demonstrating that ring strain could induce new reactivity for higher coordinate aluminium species. The formed AlC_3 aluminacyclobutenes are unique and new additions to strained aluminium ring systems. Compounds 2 and 3 may undergo interesting insertion reactions with a number of unsaturated small molecules to generate novel aluminium ring systems because of the two distinct Al–C bonds and the ring strain. Investigation of new reactions of 2 and 3 is currently being undertaken.

We are grateful to the National Natural Science Foundation of China (Project No. 20572050 and 20421202) and the NCET for support of this work.

Notes and references

† 2: A solution of 1 (0.25 g, 0.4 mmol) in *n*-hexane at room temperature was exposed to dry carbon monoxide under normal pressure. A colour change to yellow was observed within 2 min. It was stirred at room temperature for an additional 20 min under a CO atmosphere. All volatiles were removed to give a pale yellow solid, which was washed with n-hexane to afford 2: yield 72%; mp 126 °C (dec.); anal. found for C₃₈H₅₉AlN₂OSi₂: C 70.63, H, 9.53. Calc. Ĉ 70.98, H, 9.41; ¹H NMR (C₆D₆, 400 MHz): δ -0.07 (s, 9H, SiMe₃), 0.43 (s, 9H, SiMe₃), 1.03 (d, 6H, J = 6.80 Hz, $CHMe_2$), 1.08 (d, 6H, J = 6.80 Hz, $CHMe_2$), 1.19 (d, 6H, J = 6.80 Hz, CHMe₂), 1.36 (d, 6H, J = 6.80 Hz, CHMe₂), 1.43 (s, 6H, β -CMe), 3.13 (sept, 2H, CHMe₂), 3.26 (sept, 2H, CHMe₂), 4.74 (s, 1H, γ -CH), 6.97 (m, 2H, Ar–H), 7.04 (m, 4H, Ar–H); ¹³C NMR (C₆D₆): δ 1.23 (SiMe₃), 2.66 (SiMe3), 22.13 (CHMe2), 23.02 (CHMe2), 25.86 (CHMe2), 26.80 (CHMe2), 30.21 (C*Me*), 31.07 (CHMe₂), 123.4, 126.8, 128.6, 129.6, 142.3, 145.5, 146.8, 148.9, 150.4 (AlCC, Ar–C), 234.8 (CO); IR v/cm⁻¹: 1677 (s, AlCO). **3**: To a solution of 1 (0.20 g, 0.32 mmol) in n-hexane (15 mL) was added neat tertbutyl isocyanide (0.027 g, 0.32 mmol) at -78 °C. The colour of the solution turned into orange immediately. The resulting mixture was allowed to warm up to room temperature and was stirred for 30 min. All volatiles were removed and the remaining solid was crystallised from diethyl ether at -25 °C to give orange crystals of 3: yield 57%; mp 145 °C (dec.); anal. found for C42H68AlN3Si2: C 72.26, H 9.96. Calc. C 72.25, H 9.82; ¹H NMR (C₆D₆): δ -0.14, 0.48 (s, 1 : 2, 9H, Si*Me*₃), 0.52, 0.55 (s, 1 : 2, 9H, SiMe₃), 0.80 (s 6H, CMe₃), 1.01 (d, 2H, J = 6.40 Hz, CHMe₂), 1.04 (d, 2H, J = 6.80 Hz, CHMe₂), 1.13 (d, 6H, J = 6.80 Hz, CHMe₂), 1.17 (d, 4H, J = 6.80 Hz, CHMe₂), 1.20 (d, 2H, J = 6.80 Hz, CHMe₂), 1.26 (d, 4H, J =6.80 Hz, CHMe₂), 1.39 (d, 2H, J = 6.80 Hz, CHMe₂), 1.42 (s, 3H, CMe₃), 1.47 (d, 6H, CMe), 2.29 (m, 4H, CHMe₂), 4.96, 5.09 (s, 1H, 1 : 2, γ-CH), 7.04–7.09 (m, 6H, Ar–H); ¹³C NMR (\tilde{C}_6D_6): δ 2.04, 2.73, 3.28 (SiMe₃), 23.90, 24.33, 24.54, 24.72, 24.85, 25.07, 25.15, 25.85, 26.64 (Me, CHMe2), 27.81, 29.03, 29.11, 29.24, 30.30 (CHMe2 and CMe3), 55.75, 55.48 (CMe3), 100.48, 101.42 (y-C), 124.23, 124.29, 125.12, 125.29, 127.41, 127.47, 139.00, 140.57, 140.82, 142.94, 144.91, 145.40 (År–*C*), 170.75, 171.27, 172.54 (N*C*Me and *C*SiMe₃), 194.62, 197.47 (Al*C*(N)). IR ν /cm⁻¹: 1587 (s, AlCN) \ddagger Crystal data for **3b**·Et₂O: C₄₆H₇₈AlN₃OSi₂, M = 772.27, triclinic, space group $P\overline{1}$, a = 10.672(3), b = 13.588(4), c = 18.468(6) Å, $\alpha = 72.022(4)$, $\beta = 13.588(4)$ 87.112(4), $\gamma = 80.639(4)^{\circ}$, $V = 2513.3(14) \text{ Å}^3$, Z = 2, $D_c = 1.020 \text{ g cm}^{-1}$ F(000) = 848, 13739 reflections measured (8763 unique). $R1 [I > 2\sigma(I)] =$ 0.0442, wR2 (all data) = 0.1361, GOF = 1.019 for 497 parameters and 0 restraints. CCDC 296206. For crystallographic data in CIF or other electronic format see DOI: 10.1039/b601056c 4: C38H59AlN2O2Si2,



M = 659.03, monoclinic, space group P_{2_1}/n , a = 16.1975(18), b = 11.3234(12), c = 22.709(2) Å, $\beta = 104.540(2)^\circ$, V = 4031.7(8) Å³, Z = 4, $D_c = 1.086$ g cm⁻³, F(000) = 1432, 22056 reflections measured (8091 unique). R1 [$I > 2\sigma(I)$] = 0.0537, wR2 (all data) = 0.1552, GOF = 0.995 for 422 parameters and 0 restraints The X-ray data were collected on a Bruker Smart-Apex II diffractometer using graphite monochromated Mo-Ka ($\lambda = 0.71073$ Å) at 294(2) K. The structure was solved by direct methods (SHELXS-97)¹⁰ and refined by full matrix least squares on F^2 . All non-hydrogen atoms were refined anisotropically and hydrogen atoms by a riding model (SHELXL-97).¹¹

- For reviews, see: (a) J. Escudié, C. Couret, H. Ranaivonjatovo and J. Satgé, *Chem. Rev.*, 1994, **130**, 427; (b) A. Sekiguchi, *Chem. Rev.*, 2003, **103**, 1429.
- W. Uhl, T. Spies, R. Koch and W. Saak, *Organometallics*, 1999, 18, 4598; (b) C. Cui, S. Köpke, R. Herbst-Irmer, H. W. Roesky, M. Noltemeyer, H.-G. Schmidt and B. Wrackmeyer, *J. Am. Chem. Soc.*, 2001, 123, 9091; (c) H.-J. Himmel, *Organometallics*, 2003, 22, 2679; (d) H. Zhu, J. Chai, H. Fan, H. W. Roesky, C. He, V. Jancik, H.-G. Schmidt, M. Noltemeyer, W. A. Merrill and P. P. Power, *Angew. Chem., Int. Ed.*, 2005, 44, 5090.
- 3 (a) H. Zhu, J. Chai, Q. Ma, V. Jancik, H. W. Roesky, H. Fan and R. Herbst-Irmer, J. Am. Chem. Soc., 2004, **126**, 10194; (b) X. Li,

H. Song, L. Duan, C. Cui and H. W. Roesky, *Inorg. Chem.*, 2006, 45, 1912.

- A. A. Wojcicki, Adv. Organomet. Chem., 1973, 11, 87; (b)
 E. J. Kuhlmann and J. J. Alexander, Coord. Chem. Rev., 1980, 33, 195; (c)
 E. Singleton and H. E. Oosthuizen, Adv. Organomet. Chem., 1983, 22, 209; (d)
 Y. Yamamoto, Coord. Chem. Rev., 1980, 32, 193; (e)
 P. M. Treichel, Adv. Organomet. Chem., 1973, 11, 21; (f)
 F. A. Cotton,
 G. Wilkinson, C. A. Murillo and M. Bochmann, Advanced Inorganic Chemistry, John Wiley & Sons, New York, 6th edn, 1999, p. 246.
- 5 For hydroalumination of isocyanides, see: (a) W. Zheng, A. Stasch, J. Prust, H. W. Roesky, F. Cimpoesu, M. Noltemeyer and H.-G. Schmidt, Angew. Chem., Int. Ed., 2001, 40, 3461; (b) R. J. Wehmschulte and P. P. Power, Inorg. Chem., 1998, 37, 6906.
- 6 M. R. Mason, B. Song and K. Kirschbaum, J. Am. Chem. Soc., 2004, 126, 11812.
- 7 P. J. Shapiro, A. Vij, G. P. A. Yap and A. L. Rheingold, *Polyhedron*, 1995, 14, 203.
- 8 R. M. Silverstein, G. C. Bassler and T. C. Morril, in *Spectrometric Identification of Organic Compounds*, Wiley, New York, 5th edn, 1991.
- 9 L. D. Durfee and I. P. Rothwell, *Chem. Rev.*, 1988, **88**, 1059.
- 10 SHELXS-90/96, Program for Structure Solution: G. M. Sheldrick, Acta Crystallogr., Sect. A, 1990, 46, 467.
- 11 G. M. Sheldrick, SHELXL-97, Program for Crystal Structure Refinement, University of Göttingen, 1997.