A switchable macrocycle–clip complex that functions as a NOR logic gate \dagger

Pin-Nan Cheng, Pinn-Tsong Chiang and Sheng-Hsien Chiu*

Received (in Cambridge, UK) 25th November 2004, Accepted 4th January 2005 First published as an Advance Article on the web 21st January 2005 DOI: 10.1039/b417823h

We have synthesized a new molecular switch—based on a macrocycle–clip complex—whose switching behavior not only can be controlled through the use of either K^+ –[2,2,2]cryptand or NH_4^+ –Et₃N systems but also provides color changes that are visible to the naked eye; consequently, this system operates as a two-input NOR functioning molecular logic gate.

Although many elegant pseudorotaxane-like¹ molecular switches² have been developed in recent years, the number of recognition motifs that can be exploited for the preparation of these complexes remains limited because of structural complexity and/or weak intermolecular interactions. Thus, we became interested in preparing new supramolecular assemblies by taking advantage of the increased binding affinities that clip-shaped guest molecules exert toward their macrocyclic hosts. Although many molecular tweezers³ and clips⁴ have been prepared as effective synthetic receptors, there are only a few examples of molecular clips that form complexes with macrocycles;5 of these macrocycle-clip complexes, we are unaware of any that can be switched between their threaded and unthreaded states by the application of external stimuli.6 The addition and removal of metal ions66,7 and protons^{6b,8} are efficient methods for switching supramolecular systems between different states; such systems that can be switched in two different ways and exhibit detectable gains or losses of the optical signals of their different states may be considered as molecular logic gates.9 In this paper, we report a new molecular switch-based on a macrocycle-clip complex-whose switching behavior not only can be controlled through the use of either K⁺-[2,2,2]cryptand or NH₄⁺-Et₃N systems but also provides color changes that are visible to the naked eye; consequently, this system operates as a two-input NOR¹⁰ functioning molecular logic gate.

We synthesized macrocycle $1.2PF_6$ in five steps from 2,4'-dibromoacetophenone. The structure of $1.2PF_6$ is reminiscent of a combination of a ring-expanded [18]crown-6 (18C6) unit—which we expected to bind to alkali metal ions—and a π -electrondeficient aromatic system, which we expected would bind π -electron-rich units, such as tetrathiafulvalene (TTF) (Scheme 1). Because of its flexible molecular structure, relative to those of 18C6 itself and other effective cyclophane-based π -electron acceptors,¹¹ we expected that $1.2PF_6$ would have quite low affinities for both alkali metal ions and π -electron-rich aromatic rings. Thus, we synthesized the glycoluril-based compound **2** in which two TTF molecules are positioned parallel to each other as the side walls of a molecular clip. We anticipated that the complementary electronic nature of the TTF and pyridinium motifs, the good separation (6.7 Å) for π -stacking of a pyridinium moiety between the two parallel aromatic side walls of the glycoluril-based molecular clip, and π -stacking interactions between the non-threaded "*exo*" TTF arm of clip **2** and the pyridinium rings of the macrocycle **1**·2PF₆ would all assist molecular clip **2** to complex with **1**·2PF₆ with reasonable affinity. The four triethylene glycol chains of clip **2** are present not only to increase its solubility but also to allow [C–H…O] hydrogen bonds to form between the pyridinium α -protons and the oxygen atoms of these triethylene glycol side chains, which are interactions that may partially decrease the free energy of binding during the complexation process.

Although clip 2 is only moderately soluble in MeCN, the addition of macrocycle $1.2PF_6$ to the solution helped to dissolve any remaining insoluble clip 2; more interestingly, this process



Scheme 1 Schematic representation of the complexation and switching behavior of the macrocycle-clip complex.

 $[\]dagger$ Electronic supplementary information (ESI) available: Experimental procedures for the preparation of $1\cdot 2PF_6$, **2**, and $3\cdot 2PF_6$ and their characterization data. See http://www.rsc.org/suppdata/cc/b4/b417823h/ *shchiu@ntu.edu.tw



Fig. 1 (A) A photograph depicting the color changes (MeCN, 3 mM, 298 K) that occur in the switching process and (B) the corresponding partial UV-vis spectra (MeCN, 1.5 mM, 298 K) of (a) an equimolar mixture of 1.2PF_6 and **2**; (b) the mixture obtained after adding KPF₆ (20 equiv.) to solution (a); (c) the mixture obtained after adding [2,2,2]cryptand (20 equiv.) to solution (b); (d) the mixture obtained after adding NH₄PF₆ (20 equiv.) to solution (a); (e) the mixture obtained after adding Et₃N (20 equiv.) to solution (d).

immediately changed the color of the solution from light-yellow to puce (Fig. 1A-a). The UV spectrum of the complex formed between $1.2PF_6$ and 2 in MeCN (Fig. 1B-a) displays a charge transfer band at 533 nm, which suggests the formation of the macrocycle-clip complex $[(1 \subset 2) \cdot 2PF_6]$. A Job plot based on this absorption in MeCN affords conclusive evidence for 1 : 1 complexation (see supporting information[†]). From a ¹H NMR spectroscopic dilution experiment, we determined the binding constant between macrocycle $1.2PF_6$ and molecular clip 2 in CD₃CN to be 4900 \pm 180 M^{-1.12} Adding 20 equiv. of potassium hexafluorophosphate (KPF₆) to the puce solution of a mixture of $1.2PF_6$ and 2 in MeCN switched the color of the solution back to its original yellow hue (Fig. 1A-b). The significant decrease in the intensity of the charge transfer band in the UV-vis spectrum (Fig. 1B-b) and the appearance of the characteristic absorptions of the free clip 2 (Fig. 2b) in the ¹H NMR spectrum both suggest that dissociation of the complex $[(1 \subset 2) \cdot 2PF_6]$ occurred. The macrocycle $3.2PF_6$, whose structure differs from that of $1.2PF_6$ by the absence of the carbonyl groups, also generates a puce solution when mixed with clip 2, but the color of the solution of $[(3 \subseteq 2) \cdot 2PF_6]$ did not switch back to light yellow upon adding an



Fig. 2 Partial ¹H NMR spectra (400 MHz, CD₃CN, 298 K) of (a) a mixture of $1 \cdot 2PF_6$ and **2** (10 mM each); (b) the mixture obtained after adding KPF₆ (20 equiv.) to the solution in (a); and (c) the mixture obtained after adding [2,2,2]cryptand (20 equiv.) to the solution in (b).

excess amount of KPF₆. This observation suggests that the 18C6like motif of macrocycle $1.2PF_6$ is likely to be the one that recognizes the K⁺ ions; the conformational changes that this binding event causes the aromatic units of macrocycle $1.2PF_6$ to undergo may be the source of the negative allosteric effect.¹³

To remove the complexed K⁺ ions from macrocycle 1·2PF₆, we added an excess of a very strong binder, [2,2,2]cryptand, to the solution. As we anticipated, the puce color of the solution returned instantly (Fig. 1A-c) upon the addition of [2,2,2]cryptand. The ¹H NMR spectrum (Fig. 2c) of this mixture appears to be similar to that of the original solution of macrocycle 1·2PF₆ and clip **2** (Fig. 2a); this observation implies that the original [(1 \subset 2)·2PF₆] complex has regenerated in the solution. This threading–unthreading process can be repeated several times by adding excess amounts of K⁺ ion and [2,2,2]cryptand sequentially; the reversible color changes allow us to monitor the switching process by the naked eye.

Alternatively, this [2]pseudorotaxane-like macrocycle-clip molecular switch can be switched by the sequential addition of NH₄⁺ ions and base. Adding ammonium hexafluorophosphate (NH_4PF_6) to the puce solution of macrocycle 1.2PF₆ and clip 2 in MeCN also causes the color of the solution to turn light-yellow (Fig. 1A-d). The similar sizes and binding affinities of NH_4^+ and K⁺ ions toward 18C6 may explain these phenomena.¹⁴ Adding an excess of Et₃N to the mixture switches the color of the solution back to puce immediately (Fig. 1A-e). The addition of the base leads to the formation of NH₃ and the Et₃NH⁺ ion, neither of which binds strongly to the 18C6 motif of $1.2PF_6$, and, thus, the threading of clip 2 through the macrocycle regenerates the complex $[(1 \subset 2) \cdot 2PF_6]$. Again, the switching process can be visualized by observing the color of the solution of the macrocycle-clip complex, which can also be operated reversibly when adding NH₄PF₆ and Et₃N sequentially.

As anticipated, the addition of both KPF₆ and NH₄PF₆ to the mixture of $1.2PF_6$ and 2 also switches the color of the solution from puce to light yellow. Thus, if we consider the intensity of the absorbance at 533 nm in the UV-vis spectrum of complex $[(1 \subset 2) \cdot 2PF_6]$ as the output and the K⁺ and NH₄⁺ ions as the inputs, the switching of the macrocycle–clip system reflects the operation of a two-input NOR logic gate.

We have prepared a new macrocycle–clip complex that functions as a molecular switch. Both K^+ –[2,2,2]cryptand and NH_4^+ –Et₃N stimuli control the movement of this molecular switch between its threaded and unthreaded states.¹⁵ The color changes, which are observable to the naked eye during switching by either of these two different methods, allows the macrocycle–clip complex system to function as a two-input molecular NOR logic gate.

This work was supported by the National Science Council, Taiwan (NSC-93-2113-M-002-020).

Pin-Nan Cheng, Pinn-Tsong Chiang and Sheng-Hsien Chiu*

Department of Chemistry, National Taiwan University, Taipei, Taiwan, 10617, ROC. E-mail: shchiu@ntu.edu.tw; Fax: +886 2 24980963; Tel: +886 2 223690152 ext. 150

Notes and references

 M. Asakawa, P. R. Ashton, V. Balzani, S. E. Boyd, A. Credi, G. Mattersteig, S. Menzer, M. Montalti, F. M. Raymo, C. Ruffilli, J. F. Stoddart, M. Venturi and D. J. Williams, *Eur. J. Org. Chem.*, 1999, 985; (*b*) F. Huang, L. N. Zakharov, A. L. Rheingold, J. W. Jones and H. W. Gibson, *Chem. Commun.*, 2003, 2122.

- 2 Molecular Switches, ed. B. L. Feringa, VCH-Wiley, Weinheim, 2001.
- (a) C.-W. Chen and H. W. Whitlock, J. Am. Chem. Soc., 1978, 100, 4921; (b) S. C. Zimmerman and K. W. Saionz, J. Am. Chem. Soc., 1995, 117, 1175; (c) H. Kurebayashi, T. Haino, S. Usui and Y. Fukazawa, Tetrahedron, 2001, 57, 8667.
- 4 A. E. Rowan, J. A. A. W. Elemans and R. J. M. Nolte, Acc. Chem. Res., 1999, 32, 995.
- 5 (a) O. A. Matthews, F. M. Raymo, J. F. Stoddart, A. J. P. White and D. J. Williams, *New J. Chem.*, 1998, **22**, 1131; (b) H. M. Colquhoun, Z. Zhu and D. J. Williams, *Org. Lett.*, 2003, **5**, 4353.
- 6 The external stimuli can be chemicals, electrons or photons, see: (a) J. D. Badjic, V. Balzani, A. Credi, S. Silvi and J. F. Stoddart, *Science*, 2004, **303**, 1845; (b) B. Armaroli, V. Balzani, J.-P. Collin, P. Gaviña, J.-P. Sauvage and B. Ventura, *J. Am. Chem. Soc.*, 1999, **121**, 4397; (c) F. G. Gatti, S. León, J. K. Y. Wong, G. Bottari, A. Allieri, M. A. F. Morales, S. J. Teat, C. Frochet, D. A. Leigh, A. M. Brouwer and F. Zerbetto, *Proc. Natl. Acad. Sci. USA*, 2003, **100**, 10.
- 7 (a) J.-P. Collin, C. Dietrich-Buchecker, P. Gavina, M. C. Jimenez-Molero and J.-P. Sauvage, Acc. Chem. Res., 2001, 34, 477; (b) G. Kaiser, G. T. Jarrosson, S. Otto, Y.-F. Ng, A. D. Bond and J. K. M. Sanders, Angew. Chem., Int. Ed., 2004, 43, 1959.

- 8 A. M. Elizarov, S.-H. Chiu and J. F. Stoddart, J. Org. Chem., 2002, 67, 9175.
- 9 A. P. de Silva and N. D. McClenaghan, Chem. Eur. J., 2004, 10, 574.
- 10 NOR gates (as well as NAND gates) have high value in electronics, since multiple copies of them can be wired up to emulate all other logic types, see: R. J. Mitchell, *Microprocessor Systems: An introduction*, Macmillan, London, 1995 and ref. 9.
- 11 M. Asakawa, P. R. Ashton, V. Balzani, A. Credi, C. Hamers, G. Mattersteig, M. Montalti, A. N. Shipway, N. Spencer, J. F. Stoddart, M. S. Tolley, M. Venturi, A. J. P. White and D. J. Williams, *Angew. Chem., Int. Ed.*, 1998, **37**, 333.
- 12 K. A. Connors, Binding Constants, Wiley, New York, 1987.
- 13 M. Takeuchi, M. Ikeda, A. Sugasaki and S. Shinkai, Acc. Chem. Res., 2001, 34, 865.
- 14 (a) R. M. Izatt, K. Pawlak, J. S. Bradshaw and R. L. Bruening, *Chem. Rev.*, 1991, **91**, 1721; (b) S. Inoue and G. W. Gokel, *Cation Binding by Macrocycles*, Marcel Dekker, New York, 1990.
- 15 There are other sorts of competitive decomplexation or dethreading supramolecular systems, see: (a) E. M. Seward, R. B. Hopkins, W. Sauerer, S.-W. Tam and F. Diederich, J. Am. Chem. Soc., 1990, 112, 1783; (b) A. Credi, M. Montalti, V. Balzani, S. J. Langford, F. M. Raymo and J. F. Stoddart, New J. Chem., 1998, 22, 1061.