

Letters

Synthesis and Evaluation of a Novel Indoledione Class of Long Chain Fatty Acid Elongase 6 (ELOVL6) Inhibitors

Toshiyuki Takahashi, Tsuyoshi Nagase, Takahide Sasaki, Akira Nagumo, Ken Shimamura, Yasuhisa Miyamoto, Hidefumi Kitazawa, Maki Kanesaka, Ryo Yoshimoto, Katsumi Aragane, Shigeru Tokita, and Nagaaki Sato*

Tsukuba Research Institute, Merck Research Laboratories, Banyu Pharmaceutical Co., Ltd, Okubo 3, Tsukuba, Ibaraki 300-2611, Japan


Received March 27, 2009

Abstract: Novel indoledione derivatives were synthesized and evaluated as long chain fatty acid elongase 6 (ELOVL6) inhibitors. Systematic optimization of an indole class of lead **1** led to the identification of potent ELOVL6 selective inhibitors. Representative inhibitor **37** showed sustained plasma exposure and good liver penetrability in mice. After oral administration, **37** potently inhibited ELOVL6 activity in the liver in mice.

The incidence of type 2 diabetes has dramatically increased over the past decade. Accumulated evidence suggests a strong correlation between insulin resistance and the development of type 2 diabetes mellitus. An increase in de novo lipid synthesis and fat storage in tissues such as liver leads to the dysfunction in those tissues, i.e., insulin resistance.¹ Although it is unclear how increased intracellular lipid content exacerbates tissue and whole body insulin sensitivity, it has been suggested that increased levels of long chain fatty acyl-CoA antagonize the metabolic actions of insulin. Interestingly, recent reports suggested that alternation of the specific fatty acid component (i.e., palmitoleate) has significant impact on the insulin sensitivity of liver and whole body.^{2,3}

With regard to de novo lipid synthesis, the microsomal enzymes are responsible for the elongation of long chain fatty acids with chain length >C16 while FAS,⁴ a cytosolic enzyme, is responsible for the de novo synthesis of the fatty acid with chain lengths up to C16.⁴ Fatty acid elongation at microsomal fractions requires four sequential steps: (1) condensation between fatty acyl-CoA and malonyl-CoA to generate β -ketoacyl-CoA, (2) reduction by β -ketoacyl-CoA reductase, (3) dehydrogenation by β -hydroxyacyl-CoA dehydrogenase, and (4) reduction by *trans*-2,3-enoyl-CoA reductase.^{4,5} ELOVL enzymes are responsible for the rate-limiting initial condensation reaction.^{6,7}


So far, seven ELOVL enzymes have been identified in mammals and are designated ELOVL1–7.^{6–11} Each ELOVL enzyme exhibits different fatty acid substrate preferences and tissue distributions, suggesting that they play different physiological roles in vivo.¹² Among ELOVL enzymes, ELOVL3 and ELOVL6 show the highest homology to each other (43.7%) and are expressed in the highly lipogenic tissues such as liver



1
Human ELOVL6, IC₅₀ = 290 nM
Human ELOVL3, IC₅₀ > 10 μ M


Figure 1. Potency and structure of lead **1**.

Scheme 1^a



^a Reagents and conditions: (a) CHCl₃, reflux, 2 h; (b) SOCl₂, toluene, reflux, 3 h; (c) CHCl₃, room temp, 14 h.

Scheme 2^a



^a Reagents and conditions: (a) neat, room temp, 14 h; (b) toluene, reflux, 2 h; (c) (i) SOCl₂, toluene, reflux, 3 h, (ii) 3-anilino-5,5-dimethylcyclohex-2-en-1-one, CHCl₃, room temp, 14 h.


and adipose that play important roles in the regulation of lipid metabolism.^{6,9} ELOVL6 regulates the synthesis of stearoyl-CoA (C18:0) and *cis*-vaccenoyl-CoA (C18:1).^{6,7} Regarding the regulatory mechanism for expression, liver ELOVL6 expression is up-regulated by refeeding after fasting, by a high carbohydrate diet, and in obese rodents.^{6,7,13} At the molecular level, ELOVL6 expression is directly and primarily regulated by sterol regulatory element-binding protein (SREBP)-1c.¹⁴








Recent investigations using gene-deleted mice for ELOVL6 suggest that ELOVL6 regulates the hepatic insulin sensitivity by modulating the fatty acid components.³ Although increasing information on the physiological and pathological roles of ELOVL6 has drawn much attention, a lack of useful chemical tools makes it difficult to address the pharmacological roles of ELOVLs and their therapeutic potentials.

We recently reported the establishment of a homogeneous enzyme assay for ELOVL6, which is applicable to UHTS using


* To whom correspondence should be addressed. Phone: 81-29-877-2004. Fax: 81-29-877-2029. E-mail: nagaaki_sato@merck.com.

^a Abbreviations: ELOVL, long chain fatty acid elongase; FAS, fatty acid synthase; UHTS, ultrahigh-throughput screen; SAR, structure–activity relationship; SCD-1, stearoyl-CoA desaturase 1.

Table 1. Human ELOVL6 Inhibitory Activity of **1–17**^a


| compd | R ¹ | human ELOVL6 ^b (IC ₅₀ , nM) |
|-----------|---|---|
| 1 | Ph | 290 ± 61 |
| 2 | H | 1710 ± 308 |
| 3 | Me | 520 ± 148 |
| 4 | Et | 550 ± 185 |
| 5 | <i>n</i> -Pr | 185 ± 38 |
| 6 | <i>i</i> -Pr | 600 ± 155 |
| 7 | cyclopropyl | 144 ± 48 |
| 8 | cyclobutyl | 271 ± 67 |
| 9 | cyclopentyl | 411 ± 24 |
| 10 | cyclohexyl | 210 ± 60 |
| 11 |  | 879 ± 74 |
| 12 |  | 6340 ± 1360 |
| 13 |  | 226 ± 19 |
| 14 |  | 566 ± 130 |
| 15 |  | 6520 ± 2420 |
| 16 |  | 245 ± 53 |
| 17 |  | 603 ± 132 |


^a The values represent the mean ± SE for *n* ≥ 3. ^b Inhibitory activity of compounds on human ELOVL6 for palmitoyl-CoA elongation.














Table 2. Human ELOVL6 Inhibitory Activity of **18–23**^a


| compd | R ² | R ³ , R ⁴ | human ELOVL6 ^b (IC ₅₀ , nM) |
|-----------|----------------|---|---|
| 18 | Et | R ³ = R ⁴ = Me | 99 ± 28 |
| 19 | <i>i</i> -Pr | R ³ = R ⁴ = Me | >10000 |
| 20 | Me | R ³ = R ⁴ = H | 4500 ± 190 |
| 21 | Me | R ³ = H, R ⁴ = Me | 704 ± 97 |
| 22 | Me | R ³ = H, R ⁴ = Ph | 6260 ± 428 |
| 23 | Me | -(CH ₂) ₃ - | 123 ± 18 |

^a The values represent the mean ± SE for *n* ≥ 3. ^b Inhibitory activity of compounds on human ELOVL6 for palmitoyl-CoA elongation.

the recombinant histidine-tagged ACBP as a molecular probe for the detection of radioactive products of ELOVL6.¹⁵ Our

Table 3. Human ELOVL3 and 6 Inhibitory Activity of **24–38**^a


| compd | R ⁵ | human ELVOL6 ^b (IC ₅₀ , nM) | human ELOVL3 ^c (IC ₅₀ , nM) |
|-----------|--|---|---|
| 24 | Me | 2020 ± 389 | |
| 25 | <i>i</i> -Pr | >10000 | |
| 26 |  | >10000 | |
| 27 |  | 4330 ± 2560 | |
| 28 |  | 44 ± 4 | |
| 29 |  | 12 ± 3 | 221 ± 15 |
| 30 |  | 58 ± 17 | |
| 31 |  | 43 ± 12 | |
| 32 |  | 17 ± 0.4 | 503 ± 69 |
| 33 |  | 44 ± 9 | |
| 34 |  | 14 ± 4 | 1100 ± 142 |
| 35 |  | 10 ± 2 | 59 ± 14 |
| 36 |  | 8.7 ± 3.0 | 375 ± 50 |
| 37 |  | 8.9 ± 2.7 | 337 ± 74 |
| 38 |  | 196 ± 88 | |

^a The values represent the mean ± SE for *n* ≥ 3. ^b Inhibitory activity of compounds on human ELOVL6 for palmitoyl-CoA elongation. ^c Inhibitory activity of compounds on human ELOVL3 for stearoyl-CoA elongation.

corporate chemical collection was screened against human ELOVL6, resulting in the identification of a novel indoleione class of lead **1**, which has an IC₅₀ of 290 nM for human ELOVL6 (Figure 1). Systematic SAR studies of the indoleione lead **1**, aiming at optimization of ELOVL6 activity, resulted in the discovery of potent ELOVL6 inhibitors. In this report, preliminary results of the SAR studies of the indoleione class of ELOVL6 inhibitors and the discovery of the potent ELVOL6 selective inhibitor **37** are reported.

The synthesis of the indoleione ELOVL6 inhibitors **1–38** reported herein is outlined in Schemes 1 and 2. The substituted

Table 4. Human ELOVL Subtype Selectivity and Mouse ELOVL3 and 6 Activity of **37**^a

| | enzyme | | | | | | |
|------------------------------|-------------|-------------|-----------------|-------------|------------------|---------------|-----------------|
| | hELOVL1 | hELOVL2 | hELOVL3 | hELOVL5 | hELOVL6 | mELOVL6 | mELOVL3 |
| activity (IC ₅₀) | >10 μ M | >10 μ M | 337 \pm 74 nM | >10 μ M | 8.9 \pm 2.7 nM | 31 \pm 4 nM | 194 \pm 26 nM |

^a Inhibitory activity of **37** on ELOVLs for their respective substrates. The values are the mean \pm SE for $n \geq 3$.

Table 5. Plasma Exposure of **37** in Mice

| | time | | | |
|--------------------------------------|------|-----|-----|------|
| | 2 h | 4 h | 8 h | 24 h |
| plasma level (μ M) ^a | 3.4 | 2.8 | 1.4 | 0.4 |

^a The plasma levels were determined after 1 mg/kg oral administration. The values represent the mean for $n = 3$ animals.

pyrazol-3-one **A** and methyl 3,3,3-trifluoro-2-oxopropanoate were coupled to afford **B**, which was dehydrated using thionyl chloride to give enone **C**. Cyclocondensation of the enone **C** with the substituted 3-aminocyclohex-2-en-1-one **D** provided the desired indoleione derivatives **1–36**. The derivatives **37** and **38** were prepared by a slightly modified method (Scheme 2). Ethyl 3-oxobutanoate and methyl 3,3,3-trifluoro-2-oxopropanoate were coupled to give methyl 3,5-dideoxy-3-(ethoxycarbonyl)-2-C-(trifluoromethyl)pent-4-ulosonate, which was condensed with the appropriate substituted hydrazine to give **B**. Compound **B** was dehydrated with thionyl chloride followed by cyclocondensation with 3-anilino-5,5-dimethylcyclohex-2-en-1-one to afford **37** or **38**.

The compounds were optimized based on human ELOVL6 inhibitory activity, and the potent human ELOVL6 inhibitors were tested for human ELOVL3 inhibitory activity due to the high sequence homology among the ELOVL family members.

Substituent effects of R¹ of the lead compound **1** were investigated initially (Table 1). The R¹ substituent was essential for potency, as shown by a 6-fold potency decrease of **2**. The methyl and ethyl derivatives (**3** and **4**) also showed decreased activities, while the *n*-propyl derivative **5** was equipotent to lead **1**. The branched isopropyl derivative **6** displayed a decrease in activity. A series of cycloalkyl derivatives (**7–10**) were prepared and evaluated. Of them, the cyclopropyl derivative **7** was slightly more potent than the parent **1**. Benzyl substitution as in **11** was not tolerated. In light of the substituted phenyl derivatives (**12–17**), positional scanning was carried out with electron withdrawing chloro and electron donating methoxy groups. However, no noticeable substituent effect was identified, and the substitution at the meta-position was found to retain potency.

Next, R²–R⁴ substituent effects were studied briefly (Table 2). The ethyl derivative **18** was slightly more potent than the parent **1**; however, the isopropyl derivative **19** showed a complete loss of potency. Because this portion of the molecule





Figure 2. Effects of **37** on elongase activity in the liver of C57BL/6J mice. Male C57BL/6J mice were orally administered 0.1, 0.3, and 1 mg/kg **37** (dissolved in 0.5% methylcellulose), and 1 h later [¹⁴C]palmitic acid was interperitoneally administered at 10 μ Ci/body. At 2 h postdosing of **37**, fatty acids were extracted and measured by radio-HPLC to calculate the elongation index (C16/C18 ratio).

seems to be sensitive to steric effects, further investigation for the R² substituent was not attempted. Regarding the R³ and R⁴ substituents, demethylation as in **20** was found to be detrimental to potency. The monomethyl derivative **21** was less potent than **1**, and the phenyl derivative **22** displayed a further decrease in potency. It was delightful to find that the cyclobutyl derivative **23** displayed an IC₅₀ of 123 nM, which is twice as potent as lead **1**.

Finally, the substituent effect of the R⁵ group was studied (Table 3). Methyl substitution as in **24** showed decreased activity, and the isopropyl and cyclohexyl derivatives (**25** and **26**) showed complete loss of potency. Hence, we decided to focus on substituted phenyl derivatives. Positional scanning with chloro and methoxy groups was carried out. As a result, substitution of the para-position appeared to be the most promising. A series of the para-substituted derivatives **33–38** were prepared and evaluated. With the exception of the phenoxy derivative **38**, all the para-substituted phenyl derivatives displayed potent ELOVL6 activities. The potent ELOVL6 inhibitors were tested for ELOVL3 inhibitory activity. The test compounds **29**, **32**, **34–37** showed moderate to good ELOVL6 selectivity over ELOVL3 as shown in Table 3. The isopropyl derivative **35** is the least selective where the selectivity (a ratio

Scheme 3. Enzymatic Reactions of Palmitoyl-CoA with ELOVL6 and SCD-1

of ELOVL3 to ELOVL6) is ~6-fold. Of them, compound **37** is a significantly more potent and selective inhibitor (Table 3). Compound **37** was shown to be selective over human ELOVL1, -2, and -5 enzymes and has a potent inhibitory activity for the mouse ELOVL6 enzyme as well (Table 4). The selectivity against mouse ELOVL3 is moderate where the ratio of mouse ELOVL3 to ELOVL6 is ~6-fold.

The plasma and liver levels 2 h following 10 mg/kg oral administration of **37** in mice were determined to be 30 and 50 μ M, respectively; therefore, **37** is demonstrated to be highly liver penetrable (a liver-to-plasma ratio of 1.7). After oral dosing at 1 mg/kg in mice, **37** exhibited sustained plasma exposure as shown in Table 5.

Having demonstrated potent ELOVL6 activity and good exposure, **37** was evaluated for its effects on the fatty acid profile in the liver in mice. ELOVL6 is mainly responsible for the elongation of palmitoyl-CoA. Enzymatic reactions of palmitoyl-CoA involving ELOVL6 and SCD-1 are depicted in Scheme 3. When ELOVL6 elongation takes place first, palmitoyl-CoA is elongated to give stearoyl-CoA followed by desaturation by SCD-1 to give oleoyl-CoA. When the desaturation process by SCD-1 takes place first, palmitoyl-CoA is converted to palmitoleoyl-CoA, which is elongated by ELOVL6 to yield *cis*-vaccenoyl-CoA. We defined the elongation index as follows: elongation index = C18 fatty acids/C16 fatty acids = [stearoyl acid + oleic acid + *cis*-vaccenic acid]/[palmitic acid + palmitoleic acid]. The elongation index was used as a surrogate readout for ELOVL6 inhibitory activity in the liver using [14 C]palmitic acid as a radiotracer. After oral administration, **37** potently and dose-proportionally suppressed the elongation index in the liver in mice (Figure 2).

In conclusion, new indoledione derivatives were synthesized and evaluated as ELOVL6 inhibitors. After systematic SAR studies of the R¹–R⁵ substituents of the screen hit **1**, several para-substituted phenyl derivatives were identified as potent ELOVL6 inhibitors. Representative potent inhibitor **37** was selective for human ELOVL6 over human ELOVL1, -2, -3, and -5 enzymes. Compound **37** displayed potent inhibitory activity for mouse ELOVL6 as well; however, selectivity against mouse ELOVL3 was moderate. Sustained plasma exposure and appreciable liver penetration were observed after oral administration of **37** in mice. After oral dosing, **37** potently and dose-proportionally suppressed the elongation index of fatty acids in the liver in mice. The potent and selective inhibitor **37** would be a powerful tool to probe the pharmacology of the ELOVL6 enzyme inhibition in vivo. The updated results will be reported in due course.

Acknowledgment. We thank Hirokazu Ohsawa for collecting the high-resolution mass spectral data. We also thank Dr. Peter T. Meinke (Merck Research Laboratories, Rahway, NJ) for the editing of this manuscript.

Supporting Information Available: Synthetic procedures for the preparation of **1**–**38**, biological methods, HPLC retention times,

and purity for the target compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Unger, R. H. Minireview: weapons of lean body mass destruction: the role of ectopic lipids in the metabolic syndrome. *Endocrinology* **2003**, *144*, 5159–5165.
- (2) Cao, H.; Gerhold, K.; Mayers, J. R.; Wiest, M. M.; Watkins, S. M.; Hotamisligil, G. S. Identification of a lipokine, a lipid hormone linking adipose tissue to systemic metabolism. *Cell* **2008**, *134*, 933–944.
- (3) Matsuzaka, T.; Shimano, H.; Yahagi, N.; Kato, T.; Atsumi, A.; Yamamoto, T.; Inoue, N.; Ishikawa, M.; Okada, S.; Ishigaki, N.; Iwasaki, H.; Iwasaki, Y.; Karasawa, T.; Kumadaki, S.; Matsui, T.; Sekiya, M.; Ohashi, K.; Hasty, A. H.; Nakagawa, Y.; Takahashi, A.; Suzuki, H.; Yatoh, S.; Sone, H.; Toyoshima, H.; Osuga, J.; Yamada, N. Crucial role of a long-chain fatty acid elongase, Elov6, in obesity-induced insulin resistance. *Nat. Med.* **2007**, *13*, 1193–1202.
- (4) Nugteren, D. H. The enzymic chain elongation of fatty acids by rat-liver microsomes. *Biochim. Biophys. Acta* **1965**, *106*, 280–290.
- (5) Barrett, P. B.; Harwood, J. L. Characterization of fatty acid elongase enzymes from germinating pea seeds. *Phytochemistry* **1998**, *48*, 1295–1304.
- (6) Matsuzaka, T.; Shimano, H.; Yahagi, N.; Yoshikawa, T.; Amemiya-Kudo, M.; Hasty, A. H.; Okazaki, H.; Tamura, Y.; Iizuka, Y.; Ohashi, K.; Osuga, J.; Takahashi, A.; Yato, S.; Sone, H.; Ishibashi, S.; Yamada, N. Cloning and characterization of a mammalian fatty acyl-CoA elongase as a lipogenic enzyme regulated by SREBPs. *J. Lipid Res.* **2002**, *43*, 911–920.
- (7) Moon, Y. A.; Shah, N. A.; Mohapatra, S.; Warrington, J. A.; Horton, J. D. Identification of a mammalian long chain fatty acyl elongase regulated by sterol regulatory element-binding proteins. *J. Biol. Chem.* **2001**, *276*, 45358–45366.
- (8) Tvrdik, P.; Westerberg, R.; Silve, S.; Asadi, A.; Jakobsson, A.; Cannon, B.; Loison, G.; Jacobsson, A. Role of a new mammalian gene family in the biosynthesis of very long chain fatty acids and sphingolipids. *J. Cell. Biol.* **2000**, *149*, 707–718.
- (9) Tvrdik, P.; Asadi, A.; Kozak, L. P.; Nedergaard, J.; Cannon, B.; Jacobsson, A. Cig30, a mouse member of a novel membrane protein gene family, is involved in the recruitment of brown adipose tissue. *J. Biol. Chem.* **1997**, *272*, 31738–31746.
- (10) Zhang, K.; Kniazeva, M.; Han, M.; Li, W.; Yu, Z.; Yang, Z.; Li, Y.; Metzker, M. L.; Allikmets, R.; Zack, D. J.; Kakuk, L. E.; Lagali, P. S.; Wong, P. W.; MacDonald, I. M.; Sieving, P. A.; Figueroa, D. J.; Austin, C. P.; Gould, R. J.; Ayyagari, R.; Petrukhin, K. A 5-bp deletion in ELOVL4 is associated with two related forms of autosomal dominant macular dystrophy. *Nat. Genet.* **2001**, *27*, 89–93.
- (11) Leonard, A. E.; Bobik, E. G.; Dorado, J.; Kroeger, P. E.; Chuang, L. T.; Thurmond, J. M.; Parker-Barnes, J. M.; Das, T.; Huang, Y. S.; Mukerji, P. Cloning of a human cDNA encoding a novel enzyme involved in the elongation of long-chain polyunsaturated fatty acids. *Biochem. J.* **2000**, *350* (Part 3), 765–770.
- (12) Jakobsson, A.; Westerberg, R.; Jacobsson, A. Fatty acid elongases in mammals: their regulation and roles in metabolism. *Prog. Lipid Res.* **2006**, *45*, 237–249.
- (13) Miyazaki, M.; Dobrzyn, A.; Man, W. C.; Chu, K.; Sampath, H.; Kim, H. J.; Ntambi, J. M. Stearoyl-CoA desaturase 1 gene expression is necessary for fructose-mediated induction of lipogenic gene expression by sterol regulatory element-binding protein-1c-dependent and -independent mechanisms. *J. Biol. Chem.* **2004**, *279*, 25164–25171.
- (14) Kumadaki, S.; Matsuzaka, T.; Kato, T.; Yahagi, N.; Yamamoto, T.; Okada, S.; Kobayashi, K.; Takahashi, A.; Yatoh, S.; Suzuki, H.; Yamada, N.; Shimano, H. Mouse Elov6 promoter is an SREBP target. *Biochem. Biophys. Res. Commun.* **2008**, *368*, 261–266.
- (15) Shimamura, K.; Miyamoto, Y.; Kobayashi, T.; Kotani, H.; Tokita, S. Establishment of a high throughput assay for long chain fatty acyl-CoA elongase using homogeneous scintillation proximity assay. *Assay Drug Dev. Technol.*, in press.

JM900391X