



Pergamon

Tetrahedron Letters 41 (2000) 2851–2854

TETRAHEDRON
LETTERS

Diastereoselective amination of vinylic sulfoximines: application to the asymmetric synthesis of functionalized β -substituted and β,β -disubstituted β -amino acids, and of γ -amino alcohols

Hans-Joachim Gais,* Ralf Loo, Parthasarathi Das and Gerhard Raabe

Institut für Organische Chemie der RWTH Aachen, Professor-Pirlet-Straße 1, 52056 Aachen, Germany

Received 2 February 2000; accepted 15 February 2000


Abstract

An asymmetric synthesis of protected β -substituted and β,β -disubstituted β -amino acids, which carry a hydroxyalkyl side chain, from sulfonimidoyl functionalized homoallylic alcohols is described. The method allows for an asymmetric synthesis of γ -amino alcohols as well. © 2000 Elsevier Science Ltd. All rights reserved.

β -Peptides are currently of high interest because of their structure and biological activities.¹ Recently, β,β -disubstituted β -peptides have attracted attention due to novel secondary structures hitherto not observed for β -peptides.² Besides the β,β -disubstituted β -peptides, those carrying a hydroxy group in the side chain would be of particular interest because of the synthesis of glyco- β -peptides as analogs of glyco- α -peptides which are of high biological relevance.³ Whereas numerous methods exist for the asymmetric synthesis of β -substituted β -amino acids,^{4,5} we are aware of only one method giving access to β,β -disubstituted β -amino acids.^{5c,5f,6} We now describe a method for the asymmetric synthesis of β -substituted and β,β -disubstituted β -amino acids of type **A**^{5a,5g} and **B**, which carry a hydroxyalkyl side chain, from sulfonimidoyl functionalized *anti*-homoallylic alcohols of type **C**⁷ and **D**⁷ (Scheme 1). The δ -hydroxy β -amino acids **A** and **B** may not only serve as starting materials for the synthesis of β -peptides and their glycosides but also for that of analogs of sperabillin and negamycin which are peptide-like natural products with interesting biological activities.⁸ An additional feature of the method described is that it also provides for an asymmetric synthesis of substituted γ -amino alcohols which have found much use in the synthesis of biologically active compounds.⁹


Treatment of the enantio- and diastereopure hydroxy sulfoximines **1a**⁷ and **1b**⁷ with trichloroacetyl isocyanate followed by the cleavage of the corresponding intermediate *N*-trichloroacetyl carbamates with aqueous NH_3 gave the carbamates **2a** (90%) and **2b** (85%),^{10a} respectively (Scheme 2). A similar successive treatment of the enantio- and diastereopure hydroxy sulfoximine **4**⁷ with trichloroacetyl isocyanate and NH_3 furnished the carbamate **5** (94%). Reaction of carbamates **2a** and **2b**^{10b} with

* Corresponding author.



Scheme 1.

*n*BuLi at low temperatures and warming the reaction mixtures to room temperature led to a smooth Hirama–Itô¹¹ cyclization to afford, after aqueous work-up, the cyclic carbamates **3a** (94%)¹² and **3b** (93%),¹² respectively, each as a single diastereomer ($\geq 98\%$ de).^{13,14} A similar cyclization of the β,β -disubstituted carbamate **5** also proceeded readily and gave after aqueous work-up the bicyclic carbamate **6** (92%)¹² as a single diastereomer ($\geq 98\%$ de). The configurations of carbamates **3a**, **3b** and **6** were determined by ¹H NMR spectroscopy in combination with NOE experiments and X-ray structure analysis of derivatives thereof (vide infra).




Scheme 2. (i) 1. Cl₃CCONCO, THF, 22°C, 2 h; 2. NH₃ (aq), 22°C, 10–18 h. (ii) 1. 1.1 equiv. of *n*BuLi, THF, –78°C–22°C, 10–18 h; 2. H₃O⁺

For the synthesis of amino acids from **3a**, **3b** and **6** the introduction of a carboxyl group at the α -position of the sulfonimidoyl group was required. Key to the realization of this transformation was the observation that the double lithiation of **3a**, **3b** and **6** with *n*BuLi proceeded readily to give quantitatively the dilithium salts Li₂-**3a**, Li₂-**3b** and Li₂-**6**, respectively, being stable at low temperatures in solution.


The reaction of Li₂-**3a** and Li₂-**6** with ClCOOMe¹⁵ gave the ester **7** (66%) as a mixture of diastereomers (84:16) and the ester **9** (70%)¹² as a single diastereomer, respectively (Scheme 3). The configurations of the new stereogenic center of esters **7** and **9** were provisionally assigned as depicted based on NMR spectroscopic investigations. The conversion of sulfoximines **7** and **9** to the protected amino acids **8** and **10**, respectively, called for a reductive removal of the sulfonimidoyl group, which was best accomplished with Raney nickel.¹⁶ Thus, reaction of **7** and **9** with Raney nickel furnished the protected β -substituted β -amino acid **8** (90%)¹² and the β,β -disubstituted β -amino acid **10** (86%),¹² respectively (Scheme 3). The configuration of the ester **10** was established by X-ray structure analysis.¹⁷

The above results prompted us to also investigate the synthesis of functionalized γ -amino alcohols from sulfoximines of type **3a** and **6**. Thus, reaction of Li₂-**3a** with MeCHO afforded the hydroxy sulfoximine **11** (67%, 28% de) as a mixture of only two diastereomers which are presumably epimers in regard to the C atom bearing the hydroxy group. Treatment of Li₂-**3a** with MeI and BnOCH₂Cl furnished with modest diastereoselectivities the substituted sulfoximines **12** (68% de) and **13** (de not



Scheme 3. (i) 1. 2.2 equiv. of *n*BuLi, THF, -78°C ; 2. ClCO_2Me , -78°C , 4 h. (ii) Ra-Ni, THF, H_2O , 22°C , 8 h

determined), respectively. The methylation of the dilithium salt $\text{Li}_2\text{-3b}$ with MeI proceeded with a higher diastereoselectivity and gave the methylated sulfoximine **14** (80% de). Sulfoximines **12–14**¹² were easily obtained diastereomerically pure by chromatography in 75%, 70% and 68% yield, respectively, based on the starting sulfoximines. The configurations of the newly generated stereogenic center of **11–14** were determined by NOE experiments. This assignment was verified in the case of **12** by X-ray structure analysis.¹⁷ Finally, treatment of **3a**, **12** and **6** with Raney nickel under similar conditions as used above afforded the protected γ -amino alcohols **15** (88%),¹² **16** (79%)¹² and **17** (90%),¹² respectively (Scheme 4).



Scheme 4. (i) MeCHO, MeI or BnOCH_2Cl , THF, -78°C . (ii) Ra-Ni, THF, H_2O , 22°C , 5–18 h

In summary, the sulfoximine route provides for a new access to enantio- and diastereopure β -substituted and β,β -disubstituted δ -hydroxy β -amino acids and to γ -amino alcohols. The starting hydroxy sulfoximines are readily available enantio- and diastereopure through hydroxyalkylation of the corresponding allylic sulfoximines which in turn are easily prepared from (+)- or (–)-*N,S*-dimethyl-*S*-phenylsulfoximine.⁷ Although in the final reduction step the chirality of the sulfonimidoyl group is lost, as it is the case in the other route to β,β -disubstituted β -amino acids,^{5c,5f} this disadvantage is compensated for by the ready accessibility of the enantiopure starting material either from commercial sources, by catalytic asymmetric synthesis or by an efficient resolution amenable to large scale.¹⁸ The synthesis of acyclic amino acids of type **B** is now underway in our laboratories.¹⁹

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft (Graduiertenkolleg 11 and Sonderforschungsbereich 380).

References

- For leading references, see: Seebach, D.; Matthews, J. L. *Chem. Commun.* **1997**, 2015. Gellman, S. H. *Acc. Chem. Res.* **1998**, *31*, 173.
- Seebach, D.; Abele, S.; Sifferlen, T.; Hänggi, M.; Gruner, S.; Seiler, P. *Helv. Chim. Acta* **1998**, *57*, 2218.
- Marcaurelle, L. A.; Bertozzi, C. R. *Chem. Eur. J.* **1999**, *5*, 1384 and references cited therein.
- Enantioselective Synthesis of β -Amino Acids*; Juaristi, E., Ed.; Wiley-VCH: Weinheim, 1997.
- For selected examples, see: (a) de Blas, J.; Carretero, J. C.; Dominguez, E. *Tetrahedron Lett.* **1994**, *35*, 4603. (b) Seebach, D.; Overhand, M.; Kühnle, F. N. M.; Martinoni, Oberer, L.; Hommel, U.; Widmer, H. *Helv. Chim. Acta* **1996**, *79*, 913. (c) Davis, F. A.; Zhou, P.; B.-C. Chen, *Chem. Soc. Rev.* **1998**, *27*, 13 and references cited therein. (d) Kim, B. J.; Park, Y. S.; Beak, P. *J. Org. Chem.* **1999**, *64*, 1705. (e) Ohtake, H.; Imada, Y.; Murahashi, S.-I. *J. Org. Chem.* **1999**, *64*, 3790. (f) Tang, T. P.; Ellman, J. A. *J. Org. Chem.* **1999**, *64*, 12. (g) Davies, S. G.; Ichihara, O. *Tetrahedron Lett.* **1999**, *40*, 9313.
- Abele, S.; Seebach, D. *Eur. J. Org. Chem.* **2000**, 1 and references cited therein.
- Gais, H.-J.; Müller, H.; Decker, J.; Hainz, R. *Tetrahedron Lett.* **1995**, *36*, 7433. Gais, H.-J.; Hainz, R.; Müller, H.; Bruns, P.; Giesen, N.; Nienstedt, S.; Raabe, G.; Runsink, J.; Decker, J.; Schleusner, M.; Hachtel, J.; Loo, R.; Woo, C.-W.; Das, P. *Eur. J. Org. Chem.*, submitted for publication.
- Sperabillin: (a) Hida, T.; Tsubotani, S.; Hori, A.; Murakami, M.; Natsugari, H.; Kozai, Y.; Harada, S. *Chem. Pharm. Bull.* **1993**, *41*, 889; (b) Ref 5g. Negamycin: (a) Korzybski, T.; Kowszyk-Gindifer, Z.; Kurytowicz, W. *Antibiotics* **1978**, *1*, 343; (b) Schmidt, U.; Stäbler, F.; Lieberknecht, A. *Synthesis* **1992**, 482.
- Gothelf, K. V.; Jorgensen, K. A. *Chem. Rev.* **1998**, *98*, 863.
- (a) Isolated as a mixture of **Z-2b** and **E-2b** in a ratio of 3:1. (b) A mixture of **Z-2b** and **E-2b** in a ratio of 3:1 was used.
- Hirama, M.; Itô, S. *Heterocycles* **1989**, *28*, 1229 and earlier work cited therein.
- Optical rotation ($[\alpha]_D^{22}$ in CH_2Cl_2): **3a**: +130.5 (c 0.55); **3b**: +179.1 (c 0.45); **6**: +89.3 (c 0.70); **8**: +51.4 (c 0.70); **9**: +49.3 (c 0.15); **10**: +23.3 (c 0.24); **12**: +140.4 (c 0.29); **13**: +81.2 (c 1.05); **14**: +134.0 (c 0.70); **15**: -44.7 (c 0.55); **16**: -28.0 (c 0.69); **17**: -37.3 (c 0.23).
- For the addition of *N*-nucleophiles to vinylic sulfoximines, see: (a) Annunziata, R.; Cinquini, M. *J. Chem. Soc., Perkin Trans. 1* **1979**, 1684. (b) Pyne, S. G. *J. Chem. Soc., Chem. Commun.* **1986**, 1686. (c) Reggelin, M.; Heinrich, T. *Angew. Chem.* **1998**, *110*, 3005; *Angew. Chem., Int. Ed. Engl.* **1998**, *37*, 2883. (d) Reggelin, M.; Zur, C. *Synthesis* **2000**, 1.
- For the asymmetric synthesis of functionalized pyrrolidines from sulfoximines of type **C** carrying an amino group at the β' -position and a chiral substituent at the N atom, see: Ref 13c.
- Pyne, S. G. *Tetrahedron Lett.* **1986**, *27*, 1691.
- Johnson, C. R.; Stark Jr., C. J. *J. Org. Chem.* **1982**, *47*, 1193.
- Crystallographic data for the structures of **12** and **10** have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication nos. CCDC-140587 and CCDC-140588. Copies of the data can be obtained free of charge on application, supplying the full journal reference, to: The Director, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK.
- Brandt, J.; Gais, H.-J. *Tetrahedron: Asymmetry* **1997**, *8*, 909 and references cited therein.
- All new compounds were fully characterized including elemental analysis.