## Lewis acid-promoted reactions of zirconacyclopentadienes with isocyanates. A one-pot three-component synthesis of multiply-substituted iminocyclopentadienes from one isocyanate and two alkynes<sup>†</sup>

Jiang Lu,<sup>a</sup> Guoliang Mao,<sup>a</sup> Wenxiong Zhang<sup>a</sup> and Zhenfeng Xi\*<sup>ab</sup>

Received (in Cambridge, UK) 30th June 2005, Accepted 2nd August 2005 First published as an Advance Article on the web 5th September 2005 DOI: 10.1039/b509142j

Multiply-substituted iminocyclopentadienes were formed from Lewis acid-promoted reactions of zirconacyclopentadienes and isocyanates *via* a one-pot three-component coupling process; the C=O double bond of the RN=C=O moiety in the isocyanate was cleaved, and the isocyanates behaved formally as a onecarbon unit with Lewis acid-dependent and substituentdependent reactions being realized.

We have recently been interested in Lewis acid-mediated reactions of organometallic compounds with organic substrates,<sup>1,2</sup> since the development of synthetic methods can be expected from a combination of transition metal-mediated C–C bond-forming reactions and Lewis acid-mediated organic transformations.<sup>1–10</sup> This strategy has been proved to be efficient and useful by us,<sup>1,2</sup> Tilley,<sup>3</sup> Negishi<sup>4</sup> and others.<sup>5–10</sup> Zirconacyclopentadienes **1**, which can be easily prepared *in situ* from two alkyne molecules and low valent zirconocene species, are generally inert towards most organic substrates.<sup>11</sup> Interestingly, as we have reported previously, zirconacyclopentadienes **1** could react smoothly with aldehydes to afford cyclopentadiene derivatives when a Lewis acid such as AlX<sub>3</sub> or BF<sub>3</sub> was present in the reaction mixture.<sup>1</sup>

Isocyanates (RN=C=O) are a class of important synthetic intermediates. The site selectivities of these unsymmetrical heterocumulenes toward nucleophiles are of great interest in heteroatom chemistry and controlled synthetic chemistry. As part of our continuous interest in the development of synthetically useful methods by combining transition metal-mediated C–C bond forming reactions with Lewis acid-mediated organic transformations, we investigated the reactions of **1** with unsymmetrical heterocumulenes such as isothiocyanates (RN=C=S) and isocyanates (RN=C=O). Iminocyclopentadiene derivatives composed of three different components were prepared in one pot in excellent yields *via* cleavage of the C=O double bond of the RN=C=O moiety (Scheme 1). Depending on the Lewis acids used and the reaction conditions, different types of products were formed. No reaction was observed when zirconacyclopentadienes 1 were treated with isocyanates. As demonstrated in Scheme 2, after addition of  $BF_3$ ·Et<sub>2</sub>O, reaction of zirconacyclopentadienes 1 with isocyanates immediately took place to afford cyclopentadienyl imines 2. Representative examples are given in Table 1.

2,3,4,5-Tetrasubstituted-1-zirconacyclopentadienes **1a** (with tetraalkyl substituents) and **1b** (with tetraaryl substituents),<sup>12</sup> and zirconaindene **1c**<sup>13</sup> all reacted with aryl isocyanates in the presence of BF<sub>3</sub>·Et<sub>2</sub>O to afford multiply-substituted iminocyclopentadiene derivatives **2a–f** and indenyl imines **2g–i**, respectively, in good to high isolated yields. This reaction is highly site selective; only the C=O double bond in the RN=C=O moiety is cleaved to form the iminocyclopentadienes. No cleavage of the C=N double bond is observed. Iminocyclopentadiene and indenimine derivatives are all interesting and useful intermediates for synthetic chemistry and organometallic chemistry.<sup>14,15</sup> The reaction reported here provides an alternative practical method for the preparation of such complex compounds, *via* a one-pot three-component procedure. It should be noted that reactions of aliphatic isocyanates with



<sup>&</sup>lt;sup>a</sup>Key Laboratory of Bioorganic Chemistry and Molecular Engineering of Ministry of Education, College of Chemistry, Peking University, Beijing, 100871, China. E-mail: zfxi@pku.edu.cn; Fax: +86-10-62751708; Tel: +86-10-62759728

<sup>&</sup>lt;sup>b</sup>State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai, 200032, China

 $<sup>\</sup>dagger$  Electronic Supplementary Information (ESI) available: Experimental details and copies of  $^1H$  and  $^{13}C$  NMR charts for all new compounds. See http://dx.doi.org/10.1039/b509142j

Run	Zirconacyclopentadiene 1		Isocyanate Ar group	Product 2		Yield of $2 (\%)^b$
1	Pr Pr Pr Pr Pr	1a	3,5-dichlorophenyl	Pr Pr Pr Pr	2a	74
2		1a	2,4-dichlorophenyl		2b	73
3		1a	4-bromophenyl		2c	81
4		1a	4-fluorophenyl		2d	77
5		1a	2-chlorophenyl		2e	75
6	Ph ZrCp <sub>2</sub> Ph Ph	1b	2-chlorophenyl	Ph C=NAr Ph Ph	2f	91
7	Bu Bu	1c	2-chlorophenyl	Bu Bu	2g	59
8		1c	4-methoxyphenyl		2h	55
9		1c	4-methylphenyl		2i	51
<sup><i>a</i></sup> Reaction <sup><i>b</i></sup> Isolated	conditions: 2 equiv. of ArNCO, 4 e yields.	equiv. of BF	$F_3$ ·Et <sub>2</sub> O; the reaction mixture	was stirred firstly at	0 °C for 1	h then at 80 °C for 1 h

Table 1 Preparation of multi-substituted iminocyclopentadienes from BF<sub>3</sub>·Et<sub>2</sub>O-mediated reaction of zirconacyclopentadienes with isocyanates<sup>a</sup>

zirconacyclopentadienes and zirconaindenes under the same reaction conditions described above generally gave messy mixtures of products. Messy mixtures of products were also obtained when isothiocyanates, either aryl or aliphatic, were applied under the

same reaction conditions. Other Lewis acids generally behaved differently from BF<sub>3</sub>. For example, CsF, InCl<sub>3</sub>, CoCl<sub>2</sub>, VCl<sub>3</sub>, TiCl<sub>4</sub> and CrCl<sub>3</sub> did not promote the above reactions. In the case of FeCl3 as the Lewis acid, a messy mixture of reaction products was observed. When AlCl<sub>3</sub> and AlBr<sub>3</sub> were used, the zirconacyclopentadienes 1 and isocyanates disappeared, but did not result in formation of isolable products. Very interestingly, when EtAlCl<sub>2</sub> was used as the Lewis acid, substituent-dependent and Lewis acid-dependent selectivities were observed. Scheme 3 shows a representative example. Results are given in Table 2. Treatment of 2,3,4,5-tetrapropyl-1-zirconacyclopentadiene 1a with ArNCO in the presence of BF3·Et2O afforded their iminocyclopentadiene derivatives in high yields (Table 1). However, on the contrary, different products were obtained only by changing the Lewis acid from BF3 to EtAlCl2 (Table 2). No iminocyclopentadiene derivatives were obtained when EtAlCl<sub>2</sub> was used as the Lewis acid. Furthermore, the substituent on the Ar ring played a very important role in



determining the type of products obtained. 2,4-Dichlorophenyl isocyanate (Table 2, run 1) and 2-chlorophenyl isocyanate (Table 2, run 2) afforded 1,6-bisamide derivatives in high isolated yields whilst 4-bromophenyl isocyanate (Table 2, run 3) gave the monoamide product.<sup>16</sup> Other aryl isocyanates such as 4-chloro-, 4-fluoro-, and 4-methyl- isocyanates afforded a mixture of bis- and monoamide derivatives. Such dramatic Lewis acid-dependent and substituent-dependent changes have not been observed in our previous work. Transmetallation of the zirconium species to an aluminium species is assumed to be essential for this reaction. Further investigation is needed to explain this novel selectivity.

Table 2EtAlCl2-mediated substituent-dependent reactions of zirco-<br/>nacyclopentadiene 1d with isocyanates $^{a}$ 

Run	Isocyanate	Yield of <b>2</b> (%)	Yield of <b>3</b> (%)	Yield of <b>4</b> (%)
1	CI NCO	0	<b>3a</b> : 89	trace
2	CI NCO	0	<b>3b</b> : 80	0
3	Br	0	0	<b>4a</b> : 54
4	CI NCO	0	<b>3c</b> : 23	<b>4b</b> : 30
5	F NCO	0	<b>3d</b> : 26	<b>4c</b> : 40
6	Me	0	<b>3e</b> : 34	<b>4d</b> : 21
<sup>a</sup> Isol	ated vields			







As given in Table 2, 4-bromophenyl isocyanate (run 3) afforded **4a** as the sole product after hydrolysis, which indicates that the intermediate can be further applied. Indeed, as demonstrated in Scheme 4, deuteriolysis of the reaction mixture with DCl/D<sub>2</sub>O afforded the deuterated product **4D** in 47% isolated yield. Addition of 2,4-dichlorophenyl isocyanate to the reaction mixture of **1a** with 4-bromophenyl isocyanate generated the mixed bisamide derivative **3f** in 39% isolated yield.

For the formation of iminocyclopentadienes 2 from the BF<sub>3</sub>mediated reaction of zirconacyclopentadienes 1 with isocyanates, a proposed reaction mechanism is given in Scheme 5. Metathesis in 6 is assumed to be a key process in the mechanism from 1 to 2.<sup>17</sup> Detailed investigation to understand the scope, application and reaction mechanisms is in progress.

This work was supported by the National Natural Science Foundation of China (29825105, 20172003, 20232010) and the Major State Basic Research Development Program (G2000077502-D). The Cheung Kong Scholars Programme and BASF are gratefully acknowledged.

## Notes and references

- Z. Xi and P. Li, Angew. Chem., Int. Ed., 2000, 39, 2950; C. Zhao, P. Li,
  X. Cao and Z. Xi, Chem.–Eur. J., 2002, 8, 4292.
- 2 C. Zhao, T. Yu and Z. Xi, *Chem. Commun.*, 2002, 142; C. Zhao, J. Yan and Z. Xi, *J. Org. Chem.*, 2003, **68**, 4355; C. Zhao, J. Lu, J. Yan and Z. Xi, *Tetrahedron Lett.*, 2003, **44**, 6895; C. Zhao, J. Lu, Z. Li and Z. Xi, *Tetrahedron*, 2004, **60**, 1417.
- B. Jiang and T. D. Tilley, J. Am. Chem. Soc., 1999, **121**, 9744; M. C. Suh,
  B. Jiang and T. D. Tilley, Angew. Chem., Int. Ed., 2000, **39**, 2870;
  M. Nakamoto and T. D. Tilley, Organometallics, 2001, **20**, 5515;
  S. A. Johnson, F. Liu, M. C. Suh, S. Zurcher, M. Haute, S. S. H. Mao and T. D. Tilley, J. Am. Chem. Soc., 2003, **125**, 4199.
- 4 E. Negishi, N. Okukado, A. O. King, D. E. van Horn and B. J. Spiegel, J. Am. Chem. Soc., 1978, 100, 2254; D. E. van Horn, L. F. Valente, M. J. Idacavage and E. Negishi, J. Organomet. Chem., 1978, 156, C20; E. Negishi, T. Takahashi, S. Baba, D. E. van Horn and N. Okukado, J. Am. Chem. Soc., 1987, 109, 2393; E. Negishi, Dalton Trans., 2005, 827.
- K. Suzuki, *Pure Appl. Chem.*, 1994, **66**, 1557; K. Suzuki, T. Hasegawa, T. Imai, H. Maeta and S. Ohba, *Tetrahedron*, 1995, **51**, 4483; K. Ohmori, K. Hatakeyama, H. Ohrui and K. Suzuki, *Tetrahedron*, 2004, **60**, 1365.
- P. Wipf and W. Xu, *Tetrahedron Lett.*, 1994, **35**, 5197; S. A. Hitchcock,
  D. R. Mayhugh and G. S. Gregory, *Tetrahedron Lett.*, 1995, **36**, 9085;
  P. Wipf, J. G. Pierle and N. Zhuang, *Org. Lett.*, 2005, **7**, 483.
- 7 T. Kamikawa, Y. Uozumi and T. Hayashi, *Tetrahedron Lett.*, 1996, 37, 3161.
- 8 G. J. Gordon, T. Luker, M. W. Tuckett and R. J. Whitby, *Tetrahedron*, 2000, 56, 2113.
- 9 B. Zheng and M. Srebnik, J. Org. Chem., 1995, 60, 3278. See also: L. Deloux and M. Srebnik, J. Org. Chem., 1994, 59, 6871.
- 10 T. Takahashi, Y. Li, T. Ito, F. Xu, K. Nakajima and Y. Liu, J. Am. Chem. Soc., 2002, **124**, 1144. See also: Y. Li, H. Matsumura, M. Yamanaka and T. Takahashi, *Tetrahedron*, 2004, **60**, 1393.
- 11 For recent reviews on the preparation and reaction chemistry of zirconacyclopentadienes see: Z. Xi and Z. Li, *Top. Organomet. Chem.*, 2004, **8**, 27; T. Takahashi, M. Kotora, R. Hara and Z. Xi, *Bull. Chem. Soc. Jpn.*, 1999, **72**, 2591; T. Takahashi, Z. Xi and M. Kotora, *Recent Res. Dev. Pure Appl. Chem.*, 1998, **2**, 515; M. Kotora, Z. Xi and T. Takahashi, *J. Synth. Org. Chem. Jpn.*, 1997, **55**, 958. See also: Z. Xi, H. Fan, S. Mito and T. Takahashi, *J. Organomet. Chem.*, 2003, **682**, 108.
- 12 E. Negishi, F. E. Cederbaum and T. Takahashi, *Tetrahedron Lett.*, 1986, 27, 2829.
- 13 S. L. Buchwald, B. T. Watson and J. C. Hoffman, J. Am. Chem. Soc., 1986, 108, 7411.
- 14 I. Nakamura and Y. Yamamoto, *Chem. Rev.*, 2004, **104**, 2127; A. A. Vizer, K. B. Yerzhanov, A. A. A. Al Quntar and V. M. Dembitsky, *Tetrahedron*, 2004, **60**, 5499; T. Shibata, K. Yamashita, E. Katayama and A. Takagi, *Tetrahedron*, 2002, **58**, 8661.
- 15 C. Wang, Q. Song and Z. Xi, *Tetrahedron*, 2004, **60**, 5207; T. Takahashi, F. Tsai, Y. Li and K. Nakajima, *Organometallics*, 2001, **20**, 4122; J. E. Hill, G. Balaich, P. E. Fanwick and I. P. Rothwell, *Organometallics*, 1993, **12**, 2911; K. Tamao, K. Kobayashi and Y. Ito, *Synlett*, 1992, 539; K. Tamao, K. Kobayashi and Y. Ito, *J. Org. Chem.*, 1989, **54**, 3517; J. J. Eisch, A. A. Aradi and K. I. Han, *Tetrahedron Lett.*, 1983, **24**, 2073; H. Yamazaki and Y. Wakatsuki, *Bull. Chem. Soc. Jpn.*, 1979, **52**, 1239; Y. Suzuki and T. Takizawa, *J. Chem. Soc., Chem. Commun.*, 1972, 837.
- 16 C. Wang, J. Lu, G. Mao and Z. Xi, J. Org. Chem., 2005, 70, 5150.
- 17 K. E. Meyer, P. J. Walsh and R. G. Bergman, J. Am. Chem. Soc., 1995, 117, 974; S. Y. Lee and R. G. Bergman, J. Am. Chem. Soc., 1996, 118, 6396; R. L. Zuckerman, S. W. Krska and R. G. Bergman, J. Am. Chem. Soc., 2000, 122, 751; R. L. Zuckerman and R. G. Bergman, Organometallics, 2000, 19, 4795; R. L. Zuckerman and R. G. Bergman, Organometallics, 2001, 20, 1792; A. W. Holland and R. G. Bergman, J. Am. Chem. Soc., 2002, 124, 9010; S. A. Blum and R. G. Bergman, Organometallics, 2004, 23, 4003.