## 2-Methylimidazole Does Not Bind to (Octaethylporphinato)iron(III) Chloride in the Presence of Methanol: A Resonance Raman Study

Tadayuki Uno,\*,1a Keiichiro Hatano,1b and Yoshifumi Nishimura<sup>1c</sup>

> Faculty of Pharmaceutical Sciences University of Tokushima Shomachi, Tokushima 770, Japan Faculty of Pharmaceutical Sciences Nagoya City University Tanabedori, Mizuho-ku, Nagoya 467, Japan Graduate School of Integrated Science Yokohama City University Seto, Kanazawa-ku, Yokohama 236, Japan

> Received October 4, 1993 Revised Manuscript Received March 7, 1994

In many heme proteins, the heme moiety is bound to the protein through the nitrogen atom of a histidyl imidazole. To elucidate the mode of action of the iron porphyrin center in the heme proteins, many studies have been done on the imidazole complexes of iron porphyrins.<sup>2-12</sup> The earlier work by Walker's group<sup>3</sup> has revealed that two imidazole bases coordinate to the iron simultaneously and/or in a stepwise manner and the bis-ligated complex is an ultimate product. In studying the binding property of 2-MeImH<sup>13</sup> to Fe(OEP)(Cl) in CH<sub>2</sub>Cl<sub>2</sub>, however, we have found a curious phenomenon: the product was not Fe(OEP)-(2-MeImH) nor Fe(OEP)(2-MeImH)<sub>2</sub>, but was the Fe(OEP)-(OMe) complex. The methoxide ligand originated from methanol, which is present as a stabilizer<sup>14</sup> in our commercial  $CH_2Cl_2$ . Employing visible-absorption and resonance Raman technique, here we report the formation of the Fe(OEP)(OMe) complex by titrating Fe(OEP)(Cl) with 2-MeImH in the presence of a trace amount of methanol.15

In Figure 1 are shown visible spectral changes upon increasing the amount of 2-MeImH. Since spectrum 1 is almost identical with the spectrum of Fe(OEP)(Cl), methanol does not bind

- (1) (a) University of Tokushima. (b) Nagoya City University. (c) Yokohama City University.
- (2) Abbot, E. H.; Rafson, P. A. J. Am. Chem. Soc. 1974, 96, 7378 (3) Walker, F. A.; Lo, M.-W.; Ree, M. T. J. Am. Chem. Soc. 1976, 98, 5552.
- (4) Nappa, M.; Valentine, J. S.; Snyder, P. A. J. Am. Chem. Soc. 1977, 99, 5799.
- (5) Pasternack, R. F.; Gillies, B. S.; Stahlbush, J. R. J. Am. Chem. Soc. 1978, 100, 2613
- (6) Wang, J.-T.; Yeh, H. J. C.; Johnson, D. F. J. Am. Chem. Soc. 1978, 100, 2400.
- (7) Satterlee, J. D.; La Mar, G. N.; Bold, T. J. J. Am. Chem. Soc. 1977, 99, 1088.
- (8) Jones, J. G.; Tondreau, G. A.; Edwards, J. O.; Sweigart, D. A. *Inorg. Chem.* 1985, 24, 296.
  (9) Byers, W.; Cossham, J. A.; Edwards, J. O.; Gordon, A. T.; Jones, J. G.; Kenny, E. T. P.; Mahmood, A.; McKnight, J.; Sweigart, D. A.; Tondreau, G. A.; Wright, T. *Inorg. Chem.* 1986, 25, 4767.

- (10) Mandon, D.; Ott-Woelfel, F.; Fischer, J.; Weiss, R.; Bill, E.; Trautwein, A. X. Inorg. Chem. 1990, 29, 2442
- (11) Hatano, K.; Safo, M. K.; Walker, F. A.; Scheidt, W. R. Inorg. Chem. 1991, 30, 1643.
- (12) Safo, M. K.; Gupta, G. P.; Watson, C. T.; Simonis, U.; Walker, F. A.; Scheidt, W. R. J. Am. Chem. Soc. 1992, 114, 7066.
- (13) Abbreviations used: ImH, imidazole; 2-MeImH, 2-methylimidazole; 1-MeIm, 1-methylimidazole; OEP, dianion of octaethylporphyrin; OMe,
- (14) Fidler, V.; Ogura, T.; Sato, S.; Aoyagi, K.; Kitagawa, T. Bull. Chem.
   Soc. Jpn. 1991, 64, 2315.
   (15) Fe(OEP)(Cl)<sup>16</sup> (Aldrich) and 2-MeImH<sup>17</sup> were recrystallized as

described previously. CH2Cl2 was doubly distilled over CaH2. Aliquots of 2-MeImH solution were added to about 0.1 mM Fe(OEP)(Cl) dissolved in CH<sub>2</sub>Cl<sub>2</sub> containing 1% CH<sub>3</sub>OH (Nakarai Chemicals, SP grade), CH<sub>3</sub>OD (99.5 atom % D, Aldrich), or CD<sub>3</sub>OD (99 atom % D, Merck), and visible<sup>17</sup> and resonance Raman<sup>18,19</sup> spectra were recorded.

(16) Uno, T., Hatano, K.; Nishimura, Y.; Arata, Y. Inorg. Chem. 1988, 27, 3215.

directly to the iron in the absence of 2-MeImH. On addition of 2-MeImH, absorption bands at 580 and 467 nm increased (spectra 2-6) with one set of isosbestic points. This indicates that the reaction proceeds with an equilibrium between Fe(OEP)(Cl) and the product species. It should be noted that spectrum 6 is quite similar to the spectrum of Fe(OEP)(OMe) we have reported.<sup>17,20</sup> The absorbance change was well explained by 1 mol of 2-MeImH equilibrating with 1 mol of Fe(OEP)(Cl). The apparent binding constant was estimated to be about 100 M<sup>-1</sup>.

In Figure 2 are shown resonance Raman spectral changes under parallel conditions with the absorption spectral experiment. The Raman lines in the 1300-1700-cm<sup>-1</sup> region are a sensitive probe for the iron states of porphyrin complexes.<sup>21-24</sup> In the absence of both methanol and 2-MeImH (A),  $\nu_4$ ,  $\nu_3$ ,  $\nu_2$ , and  $\nu_{10}$  modes<sup>25</sup> of the starting complex, Fe(OEP)(Cl), were observed at 1377, 1494, 1582, and 1629 cm<sup>-1</sup>, respectively. These frequencies are in good agreement with those reported<sup>20,26</sup> for 5-coordinate ferric high-spin complexes. The higher frequency modes were insensitive to the presence of methanol (B) and the successive addition of 2-MeImH (C-E). In the lower frequency region, however, a new line at 541 cm<sup>-1</sup> gains intensity on addition of 2-MeImH.

As shown in Figure 3, the 541-cm<sup>-1</sup> line clearly downshifted to 528 cm<sup>-1</sup> by substituting CH<sub>3</sub>OH with CD<sub>3</sub>OD, although it did not shift with CH<sub>3</sub>OD substitution (data not shown). The frequency well coincides with the  $\nu$ (Fe–OCH<sub>3</sub>) stretch in the Fe(OEP)(OMe) complex in  $CH_2Cl_2$  solution (541 cm<sup>-1</sup>)<sup>17</sup> and in the crystalline state  $(542 \text{ cm}^{-1})$ ,<sup>27</sup> the structure of which is confirmed by an X-ray study.<sup>27</sup> In both cases, the  $\nu$ (Fe–OCD<sub>3</sub>) stretch was detected at about 530 cm<sup>-1</sup>. Therefore, the present resonance Raman results have established unequivocally that, in the presence of traces of methanol, 2-MeImH does not coordinate to the iron but promotes the formation of the 5-coordinate ferric high-spin product, Fe(OEP)(OMe).

The possible acid-base chemistry may be drawn schematically as below:

$$HN \xrightarrow{N \cdots H-OMe}_{CH_3} + CI-Fe \xrightarrow{HN}_{CH_3} HN \xrightarrow{N \cdots H-CI}_{CH_3} + MeO-Fe$$

The added 2-MeImH may engage in hydrogen bonding with methanol and promote the formation of methoxide. The chloride ligand then will exchange with the methoxide to produce the Fe(OEP)(OMe) complex and 2-MeImH<sub>2</sub>+Cl<sup>-</sup>. The apparent binding constant may be the estimate of this equilibrium reaction. Since addition of ImH or 1-MeIm to Fe(OEP)(Cl)<sup>16</sup> and even to Fe(OEP)(OMe)<sup>17</sup> produced bis-coordinated complex, the methyl substituent at the 2-position may offer sterical hindrance to the binding of 2-MeImH.

It has been established by NMR<sup>28-30</sup> and X-ray<sup>31</sup> studies that  $[Fe(OEP)(2-MeImH)_2]^+$  is produced when Fe(OEP)(Cl) is

- (17) Uno, T.; Hatano, K.; Nawa, T.; Nakamura, K.; Nishimura, Y.; Arata, Y. Inorg. Chem. 1991, 30, 4322. (18) Uno, T.; Nishimura, Y.; Tsuboi, M.; Makino, R.; Iizuka, T.; Ishimura,
- Y. J. Biol. Chem. 1987, 262, 4549.
- (19) Makino, R.; Uno, T.; Nishimura, Y.; Iizuka, T.; Tsuboi, M.; Ishimura, Y. J. Biol. Chem. 1986, 261, 8376.
- (20) Uno, T.; Hatano, K.; Nishimura, Y.; Arata, Y. Inorg. Chem. 1990, 29, 2803
- (21) Spiro, T. G. Iron Porphyrins; Lever, A. B. P., Gray, H. B., Eds.; (21) Spiro, I. G. Iron Porphyrins; Lever, A. B. P., Gray, H. B., Eds.;
  Addison-Wesley: Reading, MA, 1983; Part II, p 89.
  (22) Spiro, T. G. Adv. Protein Chem. 1985, 37, 111.
  (23) Asher, S. A. Methods Enzymol. 1981, 76, 371.
  (24) Felton, R. H.; Yu, N.-T. Porphyrins 1978, 3, 347.
  (25) Abe, M.; Kitagawa, T.; Kyogoku, Y. J. Chem. Phys. 1978, 69, 4526.
  (26) Callahan, P. M.; Babcock, G. T. Biochemistry 1981, 20, 952.
  (27) Hatano, K.; Uno, T. Bull. Chem. Soc. Jpn. 1990, 63, 1825.
  (28) Satterlee, J. D.; La Mar, G. N. J. Am. Chem. Soc. 1976, 98, 2804.
  (29) Satterlee, I. D.; La Mar, G. N. Frye, I. S. J. M. Chem. Soc. 1976.
- (29) Satterlee, J. D.; La Mar, G. N.; Frye, J. S. J. Am. Chem. Soc. 1976, 98. 7275
- (30) Chacko, V. P.; La Mar, G. N. J. Am. Chem. Soc. 1982, 104, 7002.

© 1994 American Chemical Society



Figure 1. Visible spectral changes upon addition of 2-MeImH to Fe-(OEP)(Cl). Aliquots of 2-MeImH solutions were added to Fe(OEP)-(Cl) in CH<sub>2</sub>Cl<sub>2</sub> containing 1% CH<sub>3</sub>OH. Concentrations of 2-MeImH (mM): (1) 0; (2) 1.48; (3) 4.88; (4) 9.08; (5) 14.9; (6) 29.9.



Figure 2. Resonance Raman spectral changes of Fe(OEP)(Cl) upon addition of 2-MeImH. Spectra were obtained for Fe(OEP)(Cl) dissolved in  $CH_2Cl_2$  in the absence (A) or presence of 1%  $CH_3OH$  with (B) 0, (C) 5.62, (D) 11.1, (E) 18.9 mM 2-MeImH. The Raman lines with S denote those of the solvent. Excitation: 406.7 nm. Laser power: 10 mW. Slit width: 5 cm<sup>-1</sup>. Data acquisition time: 200 s.

reacted with 2-MeImH. Indeed, by the addition of 2-MeImH in great excess (about 1 M), an absorption spectrum of  $[Fe-(OEP)(2-MeImH)_2]^+$  was obtained even when methanol was present. Therefore, the  $[Fe(OEP)(2-MeImH)_2]^+$  complex is the ultimate product but the addition of 2-MeImH does not necessarily mean its binding, at least at the first stage. It should be noted that the addition of 2-MeImH in the presence of ethanol, which



Figure 3. Effect of methanol isotopes on the resonance Raman spectrum of Fe(OEP)(Cl) in the presence of 18.9 mM 2-MeImH. The solution of Fe(OEP)(Cl) was measured in  $CH_2Cl_2$  containing 1%  $CH_3OH$  (upper) or 1%  $CD_3OD$  (lower). Spectral conditions were the same as those in Figure 2.

is contained in chloroform as a stabilizer, also produced spectral changes similar to those in Figure 1.

Recently, photoreduction of iron porphyrin was reported<sup>14,32</sup> in a system quite similar to that in our present study. A trace amount of methanol was indispensable to the photoreduction,<sup>14</sup> as we have already reported.<sup>33</sup> The starting adduct, having a  $\nu_{10}$ line at 1630 cm<sup>-1</sup> before photoreduction, was assigned to a 6-coordinate ferric intermediate-spin complex,<sup>14</sup> [Fe(OEP)(2-MeImH)(CH<sub>3</sub>OH)]<sup>+</sup>. The  $\nu$ (Fe-methanol) stretching frequency (524 cm<sup>-1</sup>)<sup>14</sup> is interestingly close to the  $\nu$ (Fe-OMe) stretch in the presence of a higher amount (20%) of methanol (527 cm<sup>-1</sup>),<sup>17</sup> where the hydroxide moiety of bulk methanol hydrogen bonds to the methoxide ligand. Therefore, the starting adduct may have a methoxide ligand, which is in good agreement with the recent proposal that the primary process in the photoreduction consists of an electron transfer from the axially ligated alkoxide to iron.<sup>34</sup>

In the community of porphyrin chemists, imidazoles have been widely believed to coordinate directly to the iron. However, our present results clearly indicate that 2-MeImH does not necessarily bind to the Fe(OEP)(Cl) in the presence of a trace amount of methanol. Our finding will stimulate the reconsideration of the product species thus far reported for the reactions of iron porphyrins and imidazoles and may provide a clue for a more detailed understanding of porphyrin chemistry.

Acknowledgment. This work was supported in part by a grantin-aid (02771685,03771714,04771907 to T.U.) from the Ministry of Education, Science, Culture of Japan.

<sup>(31)</sup> Geiger, D. K.; Lee, Y. J.; Scheidt, W. R. J. Am. Chem. Soc. 1984, 106, 6339.

<sup>(32)</sup> Ozaki, Y.; Iriyama, K.; Ogoshi, H.; Kitagawa, T. J. Am. Chem. Soc. 1987, 109, 5583.

 <sup>(33)</sup> Hatano, K.; Usui, K.; Ishida, Y. Bull. Chem. Soc. Jpn. 1981, 54, 413.
 (34) Bartocci, C.; Maldotti, A.; Varani, G.; Battioni, P.; Carassiti, V. Inorg. Chem. 1991, 30, 1255.