This article was downloaded by: ["Queen's University Libraries, Kingston"] On: 03 August 2013, At: 11:52 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Anti-Alzheimer and Anti-Cox-2 Activities of the Newly Synthesized 2,3'-Bipyridine Derivatives (I)

Fawzy A. Attaby $^{\rm a}$, Azza M. Abdel-Fattah $^{\rm a}$, Labeeb M. Shaif $^{\rm a}$ & Mohamed M. Elsayed $^{\rm b}$

^a Chemistry Department, Faculty of Science, Cairo University, Giza, Egypt

^b Research Units, Hi-Care Pharmaceutical Co., Cairo, Egypt Published online: 28 Dec 2009.

To cite this article: Fawzy A. Attaby , Azza M. Abdel-Fattah , Labeeb M. Shaif & Mohamed M. Elsayed (2009) Anti-Alzheimer and Anti-Cox-2 Activities of the Newly Synthesized 2,3[']-Bipyridine Derivatives (I), Phosphorus, Sulfur, and Silicon and the Related Elements, 185:1, 129-139

To link to this article: http://dx.doi.org/10.1080/10426500902717333

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



Phosphorus, Sulfur, and Silicon, 185:129–139, 2010 Copyright © Taylor & Francis Group, LLC ISSN: 1042-6507 print / 1563-5325 online DOI: 10.1080/10426500902717333

ANTI-ALZHEIMER AND ANTI-COX-2 ACTIVITIES OF THE NEWLY SYNTHESIZED 2,3'-BIPYRIDINE DERIVATIVES (I)

Fawzy A. Attaby,¹ Azza M. Abdel-Fattah,¹ Labeeb M. Shaif,¹ and Mohamed M. Elsayed²

¹Chemistry Department, Faculty of Science, Cairo University, Giza, Egypt ²Research Units, Hi-Care Pharmaceutical Co., Cairo, Egypt

3-Aryl-1-pyridin-3-ylprop-2-en-1-ones 1a,b reacted with 2-cyanoethane-thioamide (2) to afford the corresponding 4-aryl-6-thioxo-1,6-dihydro-2,3'-bipyridine-5-carbonitriles 6a,b. The synthetic potentiality of compounds 6a,b were investigated in the present study via their reactions with several active-hydrogen containing compounds 8a-g aiming to synthesize 4-aryl-6-pyridin-3-ylthieno[2,3-b]pyridin-3-amines 10a-n via 6-(alkylthio)-4-aryl-2,3'-bipyridine-5-carbonitriles 9a-n. The structures of all newly synthesized heterocyclic compounds were elucidated by considering the data of IR, ¹H NMR, and mass spectra, as well as that of elemental analyses. Anti-Alzheimer and anti-COX-2 activities for all newly synthesized heterocyclic compounds were investigated.

Supplemental materials are available for this article. Go to the publisher's online edition of Phosphorus, Sulfur, and Silicon and the Related Elements to view the free supplemental file.

Keywords 6-(Alkylthio)-4-aryl-2, 3'-bipyridine-5-carbonitriles; 3-aryl-1-pyridin-3-ylprop-2en-1-ones; 2-cyanoethanethioamide; 6-pyridin-3-ylthieno-[2,3-*b*]pyridin-3-amines; 6-thioxo-1, 6-dihydro-2,3'-bipyridine-5-carbonitrile

INTRODUCTION

In conjunction with our previous recent work,^{1–19} and with the aim to investigate and evaluate the biological activities of the newly synthesized heterocyclic compounds, we here interested to use 3-aryl-1-pyridin-3-ylprop-2-en-1-ones as key compounds to synthesize 2,3'-bipyridine-5-carbonitriles required for several chemical transformations, as well as our medicinal chemistry programs.

RESULTS AND DISCUSSION

3-Phenyl-1-pyridin-3-ylprop-2-en-1-one²⁰ (1a) reacted with 2-cyanoethanethioamide (2) in absolute ethanol containing a catalytic amount of piperidine under reflux to afford a reaction product. The reaction product formed via a Michael addition of $-CH_2$ -

Received 8 July 2008; accepted 31 December 2008.

Address correspondence to Fawzy A. Attaby, Chemistry Department, Faculty of Science, Cairo University, Giza 12613, Egypt. E-mail: Fattaby@hotmail.com

in 2 on -CH=CH- of 1a to give the non-isolable products 3a, 4a, and 5a followed by cyclization via dehydration and dehydrogenation to give 6a. The IR (cm⁻¹) of this reaction product showed the bands of NH (3169) and CN (2220) groups. Its mass spectrum gave m/z = 289 (100%), which corresponded to the molecular weight of the formula $C_{17}H_{11}N_3S$ of the assigned structure, as well as m/z = 256 (12.5%), which corresponded to (M⁺ - SH).

In a similar manner, 3-(4-methoxyphenyl)-1-pyridin-3-ylprop-2-en-1-one (**1b**) reacted with 2-cyanoethanethioamide (**2**) under the same above-mentioned experimental conditions to give the finally isolated **6b** via the non-isolable intermediates **3b**, **4b**, and **5b**. The chemical structure of **6b** was elucidated by considering the data of IR, ¹H NMR, and mass spectra, as well as that of elemental analyses (see the Experimental section). A further confirmation of **6a,b** arose from their synthesis through another pathway via the reaction of each of **1a,b** and malononitrile (**7**) in a dispersed sulfur, morpholine, and ethanol under reflux for 2 h (Scheme 1).²¹ It important to note here that **6a,b** obtained by the two pathways are identical in all physical and chemical properties.

The synthetic potentiality of each of 6a,b was investigated through electrophilic substitution reactions using several electrophilic C-species. Thus, it has been found that 1a reacted with ethyl chloroacetate (8a) in stirred methanolic solution of sodium methoxide at room temperature for 15 min to give a reaction product. The IR (cm⁻¹) of this reaction product showed the bands of CN (2220) and CO (1734) of the newly introduced COOEt group. Its ¹H NMR (δ ppm) spectrum revealed the signals of $-SCH_2-$, $-COOCH_2CH_3$, and $-COOCH_2CH_3$ protons, and this confirm the good nucleophilicity of S in **6a** that facilitates the electrophilic attack of **8a** to afford **9a** in a very pure form and in a good yield. Furthermore, the structure of 9a was elucidated through its cyclization in ethanolic sodium ethoxide under reflux for 30 min to give a reaction product whose IR spectrum showed no bands of CN group, but instead bands of the newly formed NH₂ group were detected. Also, the ¹H NMR spectrum of this reaction product revealed signals of $-NH_2$ protons and did not reveal the signals of -CH2- protons. Considering the data of both IR and ¹H NMR, we concluded that both $-SCH_2-$ and CN functional groups in **9a** involved in the cyclization step to give the finally isolated 10a. A further confirmation of the structure of 10a was obtained through its preparation authentically via the reaction of 6a with **8a** in ethanolic sodium ethoxide under reflux for 2 h (see the Experimental section). Similarly, **6a** reacted with each of **8c**, **8e**, and **8g** in stirred methanolic solution of sodium methoxide at room temperature to give the corresponding 2-alkylthio derivatives 9c, 9e, and 9g, whose structures were elucidated by considering the data of IR and elemental analyses. Also, 9c and 9e cyclized in ethanolic sodium ethoxide under reflux for 30 min and gave the corresponding thieno[2,3-b]pyridine derivatives **10c** and **10e** respectively, which were also obtained via refluxing of **6a** with each of **8c** and **8e** in ethanolic sodium ethoxide for 2 h. Unexpectedly, 9g did not undergo a cyclization reaction under a variety of experimental conditions to give the corresponding thieno[2,3-b]pyridine derivative 10g.

In contrast to the behavior of **6a** towards each of **8a**, **8c**, **8e**, and **8g**, it has been found that **6a** reacted with each of **8b**, **8d**, and **8f** either in stirred methanolic sodium methoxide at room temperature or under reflux in ethanolic sodium ethoxide to give the corresponding thieno[2,3-b]pyridine derivatives **10b**, **10d**, and **10f**, respectively, whose structures were elucidated by considering the data of IR, ¹H NMR, and mass spectra, as well as that of elemental analyses. It is important to report here that all trials aimed at isolating compounds **9b**, **9d**, and **9f** failed under a variety of experimental





conditions. The structure of **10d** was confirmed via its preparation through another route by reaction of **6a** with or 3-chloropentan-2,4-dione (**8d**') in methanolic solution of sodium methoxide either under stirring at room temperature or under reflux for 2 h. The reaction appears to proceed via the intermediates I and II through removal of the acetic acid molecule to give 2-acetyl-3-amino-4-phenyl-6-(3-pyridyl)thieno[2,3-b]pyridine (**10d**) (see Equation 1).

In continuation to our effort to investigate the electrophilic substitution reaction along the SH group in each of **6a**,**b**, compound **6b** is a key structure for that goal. Thus, it has been found that **6b** reacted with each of **8a**, **8b**, **8c**, **8e**, **8f**, and **8g** in a stirred methanolic solution of sodium methoxide for 15 min to afford the corresponding 2-alkylthio derivatives **9h**, **9i**,



Equation 1

9j, **9l**, **9m**, and **8n** (Scheme 2). The structures of these reaction products were elucidated by considering the data of their elemental analyses, IR, and ¹H NMR (see the Experimental section). A further elucidation for these structures arose from their cyclization in methanolic solution of sodium methoxide under reflux for 30 min to give the corresponding thieno[2,3-b]pyridine derivatives 10h, 10i, 10j, 10l, 10m, and 10n, respectively. An authentic sample of each of **10h**, **10i**, **10j**, **10n**, **and 10n** obtained via a reflux mixture of **6b** and each of **8a**, **8b**, **8c**, **8e**, **8f**, and **8g** in ethanolic solution of sodium ethoxide for 2 h. In contrast to this behavior, **6b** reacted with 1-chloroacetone (**8d**) or 3-chloropentan-2,4-dione (**8d**') either in a methanolic solution of sodium methoxide at room temperature for 15 min under stirring or under reflux for 2 h in an ethanolic solution of sodium ethoxide to give directly the corresponding thieno[2,3-b]pyridine derivative **10k** (see Equation 1). The structure of **10n** was confirmed further via its preparation authentically through the hydrolysis of **10h** in ethanolic 10% KOH under reflux (see the Experimental section).

Biological Evaluation

Anti-Alzheimer activity. For compounds **1a**,**b**, their potency as anti-Alzheimer agents relative to Flurbiprofen is high enough, while after their reactions to afford the corresponding bipyridine-5-carbonitriles **6a**,**b**, their relative potency decreased. (See the Supplemental Materials online).

Structural activity relationship of anti-Alzheimer activity. Generally for compounds **9**, we note that the phenyl moiety provides the highest anti-Alzheimer activity compared with phenyl-*p*-methoxy. Thus, we can conclude that the *p*-methoxy group has no effect. The potency as anti-Alzheimer agents for S-substitution in all compounds **9** with

phenyl moiety relative to Flurbiprofen is arranged in descending order as follows: acid > ketone > ester > amide.

Anti-COX-2 activity. For the substituted pyridine derivatives 9, their potency as anti-COX-2 activities is arranged in descending order as follows: 9h, 9n, 9e, 9i, 9m, 9a, (9c, 6b), 9j, (9l, 9g). (See Figure 6 in the Supplemental Materials online).



10a, b, c, d, e, f, g, h, i, j, k, l, m, n

9a, [b], c, [d], e, [f], g, h, i, j, l, m, n

10	Ar	Z	9	Ar	Y
a	C_6H_5	COOEt	a	C_6H_5	COOEt
b	C_6H_5	CN	с	C_6H_5	CONH ₂
с	C_6H_5	CONH ₂	e	C_6H_5	COPh
d	C_6H_5	COMe	g	C ₆ H ₅	СООН
e	C_6H_5	COPh	h	C ₆ H ₄ -4-OCH ₃	COOEt
f	C_6H_5	COPh-p-Cl	i	C ₆ H ₄ -4-OCH ₃	CN
h	C ₆ H ₄ -4-OCH ₃	COOEt	j	C_6H_4 -4-OCH ₃	CONH ₂
i	C ₆ H ₄ -4-OCH ₃	CN	1	C ₆ H ₄ -4-OCH ₃	COPh
j	C ₆ H ₄ -4-OCH ₃	CONH ₂	m	C_6H_4 -4-OCH ₃	COPh-p-Cl
_					
k	C ₆ H ₄ -4-OCH ₃	COMe	n	C_6H_4 -4-OCH ₃	COOH
1	C ₆ H ₄ -4-OCH ₃	COPh			
m	C ₆ H ₄ -4-OCH ₃	COPh-p-Cl			
n	C ₆ H ₄ -4-OCH ₃	COOH			

8	X	Y	R
a	Cl	COOEt	H
b	Cl	CN	H
c	Cl	CONH ₂	H
d	Cl	COMe	H
ď	Cl	COMe	COMe
е	Br	COPh	H
f	Br	COPh-p-Cl	H
g	Cl	COOH	H

Scheme 2

F. A. ATTABY ET AL.

EXPERIMENTAL

All melting points were uncorrected. IR (KBr discs) spectra were recorded on a Shimadzu FTIR-8201PC Spectrophotometer. ¹H NMR spectra were recorded on a Varian Mercury 300 MHz, and a Varian Gemini 200 MHz spectrometers using TMS as an internal standard and CDCl₃, DMSO-d₆, and (CD₃)₂CO as solvents. Chemical shifts were expressed as δ (ppm) units. Mass spectra were recorded on Shimadzu GCMS-QP1000EX using an inlet type at 70 eV. The Microanalytical Center of Cairo University performed the microanalyses.

Synthesis of 6a,b (General Method)

Method A. A solution of each of **2** (2.60 g, 2.6 mmol) and each of **1a,b** (5.43 g and 5.8 g, 2.6 mmol) in absolute ethanol (30 mL) containing a catalytic amount of piperidine (0.4 mL) was heated under reflux for 5 h. The reaction mixture was evaporated, cooled, and triturated with ethanol. The products were formed collected by filtration, washed with cold ethanol, and then crystallized from the proper solvent to give the corresponding **6a,b**, respectively.

Method B. A mixture of dispersed sulfur (0.67 g, 1.9 mmol) and morpholine (1.7 mL, 1.9 mmol) in ethanol (50 mL) was refluxed for 20 min. Malononitrile (7) (1.30 g, 1.9 mmol) was added followed by **1a**,**b** (3.97 g and 4.54 g, 1.9 mmol), and the mixture was refluxed for 2 h. The mixture was cooled to \sim 20 °C, and 10% HCl was added to reach pH 5–6. The precipitates that formed were filtered off and washed with water and cooled ethanol, then crystallized from dioxane to give the corresponding **6a**,**b** respectively.

4-Phenyl-6-thioxo-1,6-dihydro-2,3'-bipyridine-5-carbonitrile (6a): As orange crystals, yielded by 71%, mp 240 °C, **IR** (ν cm⁻¹): 3169 (NH), 3053 (aromatic—CH) and 2220 (CN); **MS**: 289 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₁₇H₁₁N₃S of the assigned structure), 287 (M⁺–2H, 84%); 256 (M⁺–SH, 12.5%); ¹H NMR (DMSO-D₆) (δ ppm): 7.22–7.82(m, 5H, Ph, protons), 8.19–9.01 (m, 5H, pyridine H·s) and 14.10 (s, br, 1H, SH); Anal. for C₁₇H₁₁N₃S (289) Calcd./Found (%): C(70.56/70.60), H(3.83/3.90), N(14.52/14.58), S(11.08/11.16%).

4-(4-Methoxyphenyl)-6-thioxo-1,6-dihydro-2,3'-bipyridine-5-carbonitrile (6b): As orange crystals, yielded by 76%, mp 272 °C; **IR** (ν cm⁻¹): 3174 (NH), 3020 (aromatic-CH) and 2216 (CN); **MS**: 319 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₁₈H₁₃N₃OS of the assigned structure), 318 (M⁺-H, 30.6%), 287 (M⁺-S, 4.6%); ¹H NMR (DMSO-D₆) (δ ppm): 3.85 (s, 3H, OCH₃), 7.11–7.84(m, 4H, Ar), 8.17–9.22 (m, 5H, pyridine H·s) and 14.2 (s, br, 1H, SH); Anal, for C₁₈H₁₃N₃OS (319) Calcd./Found(%): C(67.69/67.72), H(4.10/4.14), N(13.16/13.19), and S (10.04/10.11%).

Synthesis of 9a, 9c, 9e, 9g-i, 9j, 9l-n: General Procedure

A solution of each of **6a**,**b** (0.29 g and 0.32 g, 1 mmol) and ethyl chloroacetate (**8a**), chloroacetonitrile (**8b**), 2-chloroacetamide (**8c**), 1-chloroacetone (**8d**), 3-chloropentane-2,4-dione (**8d**'), 2-bromo-1-phenyl-ethanone (**8e**), 2-bromo-1-*p*-chlorophenylethanone (**8f**), chloroacetic acid (**8g**) (0.12 g, 0.08 g, 0.09 g, 0.09 g, 0.13 g, 0.2 g, 0.23 g and 0.09 g 1 mmol, respectively) in sodium methoxide (prepared from 0.14 g of sodium and methanol 25 mL) was stirred at room temperature for 15 min. The formed precipitate was collected by filtration, washed with water, and crystallized from the proper solvent to give **9a**, **9c**, **9e**, **9g**, **9h**, **9i**, **9j**, **9l**, **9m**, and **9n**, respectively.

Ethyl [(5-cyano-4-phenyl-2,3'-bipyridin-6-yl)thio]acetate (9a): As pale yellow crystals (87%), mp 196 °C; **IR** υ (cm⁻¹): 3066 (C–H, aromatic), 2220 (CN), 1734 (ester CO); ¹H NMR (DMSO-D₆) (δ ppm): 1.16 (t, 3H, CH₂<u>CH₃</u>), 2.49 (s, 2H, S<u>CH₂</u>), 4.30 (q, 2H, <u>CH₂</u>CH₃), 7.56–7.79(m, 5H, phenyl H·s), 8.07–9.40 (m, 5H, pyridine H·s); Anal., for C₂₁H₁₇N₃O₂S (375), Calcd./Found (%): C(67.18/67.23), H(4.56/4.61), N(11.19/11.23), S(8.54/8.61%).

2-[(5-Cyano-4-phenyl-2,3'-bipyridin-6-yl)thio]acetamide (**9c):** As pale yellow crystals (84%), mp 265 °C; **IR** υ (cm⁻¹): 3432, 3128 (NH₂), 2215 (CN), 1684 (CO amide); Anal., for C₁₉H₁₄N₄OS (346), Calcd./Found (%): C(65.88/65.90%), H(4.07/4.11%), N(16.17/16.22%), S(9.26/9.30%).

6-[(2-Oxo-2-phenylethyl)thio]-4-phenyl-2,3'-bipyridine-5-carbonitrile (9e): As orange crystals (91%); mp 210 °C; **IR** $v(cm^{-1})$: 3058 (C—H, aromatic), 2213 (CN), 1690 (CO); Anal. for C₂₅H₁₇N₃OS (407), Calcd./Found(%): C(73.69/73.72), H(4.21/4.26), N(10.31/10.40), S(7.87/7.90%).

[(5-Cyano-4-phenyl-2,3'-bipyridin-6-yl)thio]acetic acid (9g): As orange crystals (80%); mp = 300–302 °C; **IR** v(cm⁻¹): 2927–3387 acid (OH), 3062 (CH, aromatic), 2214 (CN); Anal. for C₁₉H₁₃N₃O₂S (347), Calcd./Found (%): C(65.69/65.73), H(3.77/3.85), N(12.10/1219), S(9.23/9.30).

Ethyl{[**5-cyano-4-(4-methoxyphenyl)-2,3'-bipyridin-6-yl]thio**}acetate (**9h**): As pale yellow crystals (80%); mp 182 °C; **IR** υ (cm⁻¹): 3045 (C–H aromatic), 2214 (CN), 1736 (ester CO). ¹H NMR (DMSO-D₆) (δ ppm): 2.50 (s, 2H, –SCH₂), 3.18–3.18 (overlapped, 3H, –CH₂<u>CH₃</u>), 3.86 (s, 3H, OCH₃), 4.28 (overlapped q, 2H, –<u>CH₂</u>CH₃), 7.79 (m, 4H, Ar–H·s), 8.03–9.39 (m, 5H, pyridine); Anal. for C₂₂H₁₉N₃O₃S (405), Calcd./Found (%): C(65.17/65.21), H(4.72/4.80), N(10.36/10.43), S(7.91/8.00).

6-[(Cyanomethyl)thio]-4-(4-methoxyphenyl)-2,3'-bipyridine-5-carbonitrile (9i): As yellow crystals (84%); mp 185–157 °C; **IR** $v(cm^{-1})$: 3051 (C—H aromatic), 2214 (CN); Anal. for C₂₀H₁₄N₄OS (358) Calcd./Found (%): C(67.02/67.10), H(3.94/4.03), N(15.63/15.70), S(8.95/9.06).

2-{[5-Cyano-4-(4-methoxyphenyl)-2,3'-bipyridin-6-yl]thio}acetamide (9j): As yellow crystals (84%); mp 212 °C; **IR** $v(cm^{-1})$: 3391, 3204 (NH₂), 3079 (C—H aromatic), 2216 (CN), 1666 (CO); Anal. for C₂₀H₁₆N₄O₂S (376) Calcd./Found(%):C(63.81/63.90), H(4.28/4.33), N(14.88/14.95), S(8.52/8.61).

6-[(2-Oxo-2-phenylethyl)thio]-4-(4-methoxyphenyl)-2,3'-bipyridine-5-carbonit rile (91): As orange crystals (91%); mp 182 °C; **IR** υ (cm⁻¹): 3046 (C–H aromatic), 2213 (CN); Anal. for C₂₆H₁₉N₃O₂S (437), Calcd./Found (%): C(71.38/71.42), H(4.38/4.44), N(9.60/9.64), S(7.33/7.41).

6-{[**2**-(**4**-Chlorophenyl)-**2**-oxoethyl]thio}-**4**-(**4**-methoxyphenyl)-**2**,**3**'-bipyrid-ine-**5**-carbonitrile (**9m**): As yellow crystals (89%); mp 225 °C; **IR** $v(\text{cm}^{-1})$: 3064 (C–H aromatic), 2206 (CN), 1697 (CO); Anal. for C₂₆H₁₈ClN₃O₂S (471), Calcd./Found (%): C(66.17/66.22), H(3.84/3.90), Cl(7.51/7.60), N(8.90/9.00), S(6.79/6.85).

{[5-Cyano-4-(4-methoxyphenyl)-2,3'-bipyridin-6-yl]thio}acetic acid (9n): As orange crystals (80%); mp 278 °C; **IR** $v(cm^{-1})$: 3336–3529 (acidic OH), 3040 (C–H aromatic), 2209 (CN), 1653 (CO); ¹H NMR (DMSO-D₆) (δ ppm): 2.49 (s, 3H, –SCH₂–), 3.85 (s, 3H, OCH₃), 7.11–7.83 (m, 4H, Ar H's), 8.16–9.21 (m, 5H, pyridine H's) and 14.3 (s, br., 1H, COOH); Anal. for C₂₀H₁₅N₃O₃S (377) Calcd./Found (%): C(63.65/63.70), H(4.01/4.10), N(11.13/11.20), S(8.50/8.58).

The Synthesis of 10a-n

Method A. A solution of each of **9a**, **9c**, **9e**, **9g**, **9h**, **9i**, **9j**, **9l**, **9m**, and **9n** (0.38 g, 0.35 g, 0.41 g, 0.35 g, 0.41 g, 0.36 g, 0.38 g, 0.44 g, 0.47 g, and 0.38 g, respectively 1 mmol) in sodium ethoxide solution (prepared from 0.10 g of sodium and 25 mL ethanol) was heated under reflux for 30 min. The solid that formed after cooling was collected by filtration, washed with water and ethanol, then crystallized from the proper solvent to afford **10a–n**, respectively.

Method B. A solution of each of **6a,b** (0.29 g and 0.32 g, respectively, 1 mmole) and ethyl-chloroacetate (**8a**), chloroacetonitrile (**8b**), 2-chloroacetamide (**8c**), 1-chloroacetone (**8d**), 3-chloropentane-2,4-dione (**8d**'), 2-bromo-1-phenylethanone (**8e**), 2-bromo-1-(4-chlorophenyl)ethanone (**8f**), and chloroacetic acid (**8g**) (0.12 g, 0.08 g, 0.09 g, 0.09 g, 0.13 g, 0.20 g, 0.23 g, and 0.09 g, respectively, 1 mmole) in sodium methoxide (prepared from 0.10 g of sodium and 25 mL ethanol) heated under reflux for 2 h. The solid products so formed after cooling, collected by filtration, washed with water and ethanol, and dried, then crystallized from the proper solvent to afford **10a–n**, respectively.

Ethyl 3-amino-4-phenyl-6-pyridin-3-ylthieno[2,3-*b***]pyridine-2-carboxylate** (**10a**): As yellow crystals crystallized from ethanol (90%); mp 180 °C; **IR** υ (cm⁻¹): 3478, 3333 (NH₂), 3057 (C—H aromatic), 1666 (CO); ¹H NMR (DMSO-D₆) (δ ppm): 1.05 (t, 2H, CH₂CH₃), 3.44 (q, 2H, CH₂CH₃), 6.46 (s, 2H, NH₂), 7.48–7.89 (m, 5H, phenyl H's) and 8.57–9.38 (m, 5H, pyridine H's); Anal, for C₂₁H₁₇N₃O₂S (375), Calcd./Found (%): C(67.18/67.23), H(4.56/4.61), N(11.19/11.23), and S (8.54/8.60).

3-Amino-4-phenyl-6-pyridin-3-ylthieno[**2**,**3-***b*]**pyridine-2-carbonitrile** (**10b**): As yellow crystals crystallized from ethanol (85%); mp 274 °C; **IR** v(cm⁻¹): 3448, 3298, 3180 (NH₂), 3052(C–H aromatic), 2183 (CN); **MS** (m/z): 328 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₁₉H₁₂N₄S of the assigned structure), 254 (M⁺–Pyridyl, 17%), 166 (Pyridine–CN, CN–C=C–NH₂, 5.3%), 140 (Pyridine, CN–C=C–NH₂, 7.0%), 100 (Pyridine–CN, 5.3%) and 66 (CN–C=C–NH₂, 10.5%); Anal. for C₁₉H₁₂N₄S (328), Calcd./Found (%): C(69.49/69.52), H(3.68/3.72), N(17.06/17.12), S(9.76/9.82).

3-Amino-4-phenyl-6-pyridin-3-ylthieno[**2**,**3-***b*]**pyridine-2-carboxamide** (**10c**): As orange crystals crystallized from dioxane (80%); mp 225 °C; **IR** $v(\text{cm}^{-1})$: 3475, 3433, 3300, 3119 (NH₂ and amidic NH₂), 1645 (amidic CO); **MS** (m/z): 346 (M⁺, 100% which corresponds the molecular weight of the molecular formula C₁₉H₁₄N₄OS of the assigned structure), 330 (M⁺-NH₂, 7.5%), 329 (M⁺-NH₂-H, 50.7%), 328 (M⁺-NH₂-2H, 11.0%), 300 (M⁺-CONH₂.2H, 84.9%), 256 (M⁺-PhCN-NH₂, 35.6%), 228 (M⁺-PhCN-CONH₂, 26.0%); Anal. for C₁₉H₁₄N₄OS (346), Calcd./Found (%): C(65.88/65.94), H(4.07/4.11), N(16.17/16.21), S(9.26/9.31%).

1-(3-Amino-4-phenyl-6-pyridin-3-ylthieno[2,3-*b***]pyridin-2-yl)ethanone** (10d): As orange crystals crystallized from acetic acid (83%); mp 183 °C; **IR** ν (cm⁻¹): 3485, 3310 (NH₂), 3054 (C–H aromatic), 1620 (CO); **MS** (m/z): 345 (M+, 100% which corresponds to the molecular weight of the molecular formula C₂₀H₁₅N₃OS of the assigned structure), 344 (M⁺–H, 43.4%), 330 (M⁺–CH₃, 56.2%), 302 (M⁺–COCH₃, 36.9%);¹**H NMR** (DMSO-D₆) (*δ*ppm): 2.39 (s, 3H, COCH₃), 6.50 (s, br., 2H, NH₂) 7.48–7.92 (m, 5H, phenyl H's) and 8.57–9.39 (m, 5H, pyridine H's); Anal. for C₂₀H₁₅N₃OS (345), Calcd./Found (%): C(69.54/69.62), H(4.38/4.42), N(12.17/12.22), S(9.28/9.30).

(3-Amino-4-phenyl-6-pyridin-3-ylthieno[2,3-*b*]pyridin-2-yl)(phenyl)-methanone (10e): As orange crystals crystallized from dioxane (79%); mp = 210 °C; IR ν (cm⁻¹): 3469, 3382, 3305 (NH₂), 3057 (C–H aromatic), 1678 (CO); **MS** (m/z): 407 (M+, 84.3% which corresponds to the molecular weight of the molecular formula $C_{25}H_{17}N_3OS$ of the assigned structure), 406 (M⁺–H, 100%), 302 (M⁺–COPh, 28.2%), 105 (PhCO, 68.8%); Anal. for $C_{25}H_{17}N_3OS$ (407), Calcd./Found (%): C(73.69/73.75), H(4.21/428), N(10.31/10.38), S(7.87/7.94).

(3-Amino-4-phenyl-6-pyridin-3-ylthieno[2,3-*b*]pyridin-2-yl)(4-chlorophenyl)ethanone (10f): As yellow crystals crystallized from dioxane (86%); mp 226 °C; IR $v(cm^{-1})$: 3461, 3284 (NH₂), 3053 (C–H aromatic), 1685 (CO); MS (m/z): 442 (M⁺, 60.6% which corresponds to the molecular weight of the molecular formula C₂₅H₁₆ClN₃OS of the assigned structure), 443 (M+1, 29.9%), 440 (M⁺–2H, 100%), 301 (M⁺–COC₆H₄–p–Cl, 10.3%); Anal. for C₂₅H₁₆ClN₃OS (441.5), Calcd./Found (%): C(67.94/67.99), H(3.65/3.71), Cl(8.02/8.11), N(9.51/9.59), S(7.26/7.33).

Ethyl 3-Amino-4-(4-methoxyphenyl)-6-pyridin-3-ylthieno[2,3-*b***]pyrid-ine-2carboxylate (10h): As yellow crystals crystallized from ethanol (88%); mp 219 °C; IR** $v(cm^{-1})$: 3497, 3379 (NH₂), 3062 (C–H aromatic), 1665 (CO); **MS** (m/z): 405 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₂₂H₁₉N₃O₃S of the assigned structure), 358 (M⁺-C₂H₅, -H, 66.4%), 332 (M⁺-COOEt, 4.3%), 330 (M⁺-COOEt, -2H, 27.6%), 139 (-COC₆H₄Cl 31%); ¹**H** NMR (DMSO-D₆) (δppm): 1.28 (t, 3H, CH₂<u>CH₃</u>), 3.86 (s, 3H, OCH₃), 4.27 (q, 2H, <u>CH₂</u>CH₃), 5.90 (s, 2H, NH₂), 7.14–7.48 (m, 4H, Ar H·s) and 8.53–9.35 (m, 5H, pyridine H·s); Anal. for C₂₂H₁₉N₃O₃S (405) Calcd./Found (%): C(65.17/65.22), H(4.72/4.80), N(10.36/10.41), S(7.91/7.99).

3-Amino-4-(4-methoxyphenyl)-6-pyridin-3-ylthieno[2,3-*b***]pyridine-2-carbonit rile (10i):** As orange crystals crystallized from dioxane (90%); mp 210 °C; **IR** v(cm⁻¹): 3460, 3382, 3296 (NH₂), 3065 (C–H aromatic), 2188 (CN); **MS** (m/z): 358 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₂₀H₁₄N₄OS of the assigned structure), 357 (M⁺–H, 34.6%), 327 (M⁺–OCH₃, 17.4%), 316 (M⁺–CN–NH₂, 5.3%), 313 (M⁺–NH₂, –CN, –3H 19%); Anal. for C₂₀H₁₄N₄OS (358), Calcd./Found (%): C(67.02/6711), H(3.94/4.00), N(15.63/15.70), S(8.95/9.02).

3-Amino-4-(4-methoxyphenyl)-6-pyridin-3-ylthieno[2,3-*b***]pyridine-2-carboxa mide (10j):** As yellow crystals crystallized from acetic acid (81%); mp 268 °C; **IR** $v(\text{cm}^{-1})$: 1656 (CO), 3467, 3289 (NH₂), 3407, 3126 (NH₂); **MS** (m/z): 376 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₂₀H₁₆N₄O₂S of the assigned structure), 360 (M⁺-NH₂, 12%), 358 (M⁺-NH₂, -2H, 80.8%), 332 (M⁺-CONH₂, 5.2%), 330 (M⁺-CONH₂, -2H, 44.3%); Anal. for C₂₀H₁₆N₄O₂S (376), Calcd./Found (%): C(63.81/63.90), H(4.28/4.33), N(14.88/1494), S(8.52/8.60).

1-[3-Amino-4-(4-methoxyphenyl)-6-pyridin-3-ylthieno[2,3-*b***]pyridin-2-yl]etha none (10k):** As yellow crystals crystallized from acetic acid (86%); mp = 233–236 °C; **IR** $v(\text{cm}^{-1})$: 3445, 3288 (NH₂), 3033 (C—H aromatic), 1620 (CO); **MS** (m/z): 375 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₂₁H₁₇N₃O₂S of the assigned structure), 374 (M⁺—H, 61.7%), 360 (M⁺—CH₃ 40.5%), 332 (M⁺—COCH₃ 18.8%); Anal. for C₂₁H₁₇N₃O₂S (375), Calcd./Found (%): C(67.18/67.22), H(4.56/4.62), N(11.19/11.23), S(8.54/8.62).

(3-Amino-4-(4-methoxyphenyl)-6-pyridin-3-ylthieno[2,3-*b*]pyridin-2-yl)(phen yl)methanone (10l): As orange crystals crystallized from dioxane (72%); mp 206 °C; IR $v(\text{cm}^{-1})$: 3481, 3301 (NH₂), 3045 (C–H aromatic), 1656 (CO); MS (m/z): 437 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₂₆H₁₉N₃O₂S of the assigned structure), 436 (M⁺–H, 93%), 105 (COPh, 20.9%); Anal. for C₂₆H₁₉N₃O₂S (437), Calcd./Found (%): C(71.38//71.42), H(4.38/4.42), N(9.60/9.68), S(7.33/7.42).

(3-Amino-4-(4-methoxyphenyl)-6-pyridin-3-ylthieno[2,3-*b*]pyridin-2-yl)-(4-chl orophenyl)methanone (10m): As orange crystals crystallized from dioxane (90%); mp 205 °C; **IR** υ (cm⁻¹): 3479, 3306 (NH₂), 3040 (C—H aromatic), 1654 (CO); **MS** (m/z): 471 (M⁺, 100% which corresponds to the molecular weight of the molecular formula C₂₆H₁₈ClN₃O₂S of the assigned structure), 473 (M⁺2, 37.0%), 472 (M⁺1, 50.3%), 470 (M⁺-H, 99.2%), 469 (M⁺-2H, 11.7%), 139 (-COC₆H₄Cl, 31%); Anal. for C₂₆H₁₈ClN₃O₂S (471), Calcd./Found (%): C(66.17/66.22), H(3.84/3.90), Cl(7.51/7.60), N(8.90/8.98), S(6.79/6.82).

3-Amino-4-(4-methoxyphenyl)-6-pyridin-3-ylthieno[2,3-*b***]pyridine-2-carboxy lic acid (10n): As orange crystals crystallized from acetic acid (71%), mp 320–322 °C; IR v(cm^{-1}): 3467 (NH₂), 3230–3467 (acid OH); MS (m/z): 332 (M⁺-COOH, 64.5%), 316 (M⁺-COOH–NH₂, 43.5%), 303 (M⁺-pyridyl, 9.7%), 289 (M⁺-CO₂–CS, 64.5%);¹H NMR (DMSO-D₆) (\deltappm): 3.85 (s, 3H, OCH₃), 5.57 (s, 2H, NH₂), 7.10–7.68 (m, 4H, Ar H·s), 8.50–932 (m, 5H, pyridine H·s) and 14.3 (s, br., 1H, COO<u>H</u>); Anal. for C₂₀H₁₅N₃O₃S (377), Calcd./Found (%): C(63.65/63.71), H(4.01/4.11), N(11.13/11.19), S(8.50/8.59).**

REFERENCES

- 1. F. A. Attaby, S. M. Eldin, and M. A. Razik, Phosphorus, Sulfur, and Silicon, 106, 21 (1995).
- F. A. Attaby, S. M. Eldin, W. M. Bassouni, and M. A. A. Elneairy, *Phosphorus, Sulfur, and Silicon*, **108**, 31 (1996).
- 3. F. A. Attaby and A. M. Abdel-Fattah, Phosphorus, Sulfur, and Silicon, 119, 257 (1996).
- 4. F. A. Attaby, S. M. Eldin, W. M. Bassouni, and M. A. A. Elneairy, *Phosphorus, Sulfur, and Silicon*, **119**, 1 (1996).
- 5. F. A. Attaby, Phosphorus, Sulfur, and Silicon, 126, 27 (1997).
- 6. F. A. Attaby, Phosphorus, Sulfur, and Silicon, 139, 1 (1998).
- 7. F. A. Attaby, S. M. Eldin, and M. A. A. Elneairy, Heteroatom Chem., 9, 571 (1998).
- F. A. Attaby, S. M. Eldin, and M A. A. Elneairy, J. Chem. Res., (M), 10, 2754 (1998); (S), 10, 632 (1998).
- F. A. Attaby, M. A. A. Elneairy, and M. S. Elsayed, *Phosphorus, Sulfur, and Silicon*, 149, 49 (1999).
- 10. F. A. Attaby and A. M. Abdel-Fattah, Phosphorus, Sulfur, and Silicon, 155, 253 (1999).
- F. A. Attaby, M. A. A. Elneairy, S. M. Eldin, and A. K. K. El-Louh, J. Chin. Chem. Soc., 48, 893 (2001).
- F. A. Attaby, H. M. Mostafa, A. H. H. Elghandour, and Y. M. Ibrahem, *Phosphorus, Sulfur, and Silicon*, **177**, 2753 (2002).
- F. A. Attaby, A. H. H. Elghandour, H. M. Mustafa, and Y. M. Ibrahem, J. Chin. Chem. Soc., 49, 561 (2002).
- F. A. Attaby, S. M. Eldin, M. A. A. Elneairy, and A. K. K. Elouh, *Phosphorus, Sulfur, and Silicon*, **179**, 2205 (2004).
- F. A. Attaby, M. A. Ali, A. H. H. Elghandour, and Y. M. Ibrahem, *Phosphorus, Sulfur, and Silicon*, 181, 1 (2006).
- F. A. Attaby, A. H. H. Elghandour, Ali M. A, and Y. M. Ibrahem, *Phosphorus, Sulfur, and Silicon*, 181, 1087 (2006).
- F. A. Attaby, A. H. Elghandour, M. A. Ali, and Y. M. Ibrahem, *Phosphorus, Sulfur, and Silicon*, 182, 133 (2007).
- F. A. Attaby, A. H. H. Elghandour, M. A. Ali, and Y. M. Ibrahem, *Phosphorus, Sulfur, and Silicon*, 182, 695 (2007).
- 19. F. A. Attaby, M. M. RamLa, and E. M. Gouda, Phosphorus, Sulfur, and Silicon, 182, 517 (2007).
- 20. C. Engler and A. Engler, Berichte der Deutsehen Chemischem, xxxv, 4061 (1902).

- A. H. Shestopalov, K. G. Nikishin, A.V. Gromova, and L. A. Rodinovskaya, *Russ. Chem. Bull.*, 52, 2203 (2003).
- K. Glaser, M.-L. Sung, K. O'Neill, M. Belfast, D. Hartman, and R. Carlson, *Eur. J. Pharmacol.*, 281, 107 (1995).
- 23. K. Hsiao, P. Chapman, S. Nilsen, C. Eckman, Y. Harigaya, and A. Younkin, *Science*, **274**, 99 (1996).
- 24. S. Weggen, J. L. Eriksen, P. Das, S. Sagi, R. Wang, and C. U. Pietrzik, Nature, 414, 212 (2001).
- 25. T. Morihara, T. Chu, O. Ubeda, W. Beech, and G. M. Cole, J. Neurochem., 83, 1009 (2002).
- J. L. Eriksen, S. A. Sagi, T. E. Smith, S. Weggen, P. Das, and D. C.McLendon, J. Clin. Invest., 112, 440 (2003).
- 27. G. P. Lim, F. Yang, T. Chu, P. Chen, W. Beech, and B. Teter, J. Neurosci., 20, 5709 (2000).
- G. P. Lim, F. Yang, T. Chu, E. Gahtan, O. Ubeda, and W. Beech, *Neurobiol. Aging*, 22, 983 (2001).
- P. T. Jantzen, K. E. Condor, G. Di Carlo, G. L. Wenk, J. L. Wallace, and A. M. Rojiani, J. Neurosci., 22, 2246 (2002).
- S. Weggen, J. L. Eriksen, S. A. Sagi, C. U. Pietrzik, T. E. Golde, and E. H. Koo, *J. Biol. Chem.*, 278, 30748 (2003).