This article was downloaded by: [Michigan State University] On: 09 January 2015, At: 09:38 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Nucleosides and Nucleotides

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/lncn19

The Synthesis of 2-Fluoropurine Nucleosides

K. Krolikiewicz^a & H. Vorbrüggen^a

^a Research Laboratories, Schering AG, 13342, Berlin, Germany Published online: 23 Sep 2006.

To cite this article: K. Krolikiewicz & H. Vorbrüggen (1994) The Synthesis of 2-Fluoropurine Nucleosides, Nucleosides and Nucleotides, 13:1-3, 673-678, DOI: <u>10.1080/15257779408013271</u>

To link to this article: <u>http://dx.doi.org/10.1080/15257779408013271</u>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

THE SYNTHESIS OF 2-FLUOROPURINE NUCLEOSIDES

K. Krolikiewicz and H. Vorbrüggen*

Research Laboratories, Schering AG, 13342 Berlin, Germany

Abstract: 2-Aminoadenosine, obtained by silylation-amination from guanosine, is readily converted by $KNO_2/$ HF/Pyridine in up to 80% yield into 2-fluoradenosine, which is a convenient starting material for the preparation of 9(β -D-arabinofuranosyl)-2-fluoroadenine 5'-phosphate (Fludara). N⁶,N⁶-Pentamethylene-2-aminoadenosine and guanosine afford likewise the corresponding 2-fluoropurine nucleosides in high yields.

The potent antileukemia drug Fludarabine Phosphate (Fludara) 6^{1} has been synthesized by coupling of 2,6-dichloropurine **2** with 2,3,5-tri-O-benzyl-D-arabinosyl chloride **1** to 9-(2,3,5-tri-O-benzyl- β -D-arabinofuranosyl)-2,6-dichloropurine **3**, conversion to the protected 2,6-diaminopurine nucleoside **4** and finally Schiemann reaction to the corresponding 2-fluoro-6-amino derivative **5**. Reductive removal of the O-benzyl groups²⁾ and 5'-O-phosphorylation¹⁾ gave finally Fludarabine Phosphate (Fludara) **6** (Scheme I). Since this route as well as later routes e. g. employing 2,6-diaminopurine³⁾⁽⁴⁾ instead of **2** all involve high cost nucleoside coupling of purine-bases with the relatively expensive 2,3,5-tri-O-benzyl-D-arabinofuranosyl chloride, we investigated alternative routes.

Starting from the inexpensive guanosine **7**, silylation-amination⁵⁾⁶⁾ (Scheme II) can be readily performed on a 100-250 kg scale to give 2-aminoadenosine **8** in ca. 85-95% yield. Since the 2',3',5'-tri-O-acetate **9** of 2-aminoadenosine **8** had been converted by Robins et al.⁷⁾ in 85% yield into 2-fluoroadenosine-2',3',5'-tri-O-acetate **10** employing tert.-butyl nitrite / 60% HF-pyridine, we tried a variety of acylation conditions to transform **8** into **9**. But in accordance with Montgomery et al.²⁾ we invariably obtained mixtures of the corresponding tetra- and pentaacetates. Since the described routes of Montgomery⁸⁾⁹⁾ and Robins¹⁰⁾ to con-

Dedicated to the memory of the late R. K. Robins.







Scheme II

vert guanosine **6** via its corresponding 6-chloro- and 6-azido-2',3',5'-tri-O-acetyl-derivatives into **9** involved many steps, including some awkward and dangerous ones from a technical point of view, we wondered whether the O-acetyl-protecting groups in **9** were necessary for the Schiemann reaction. Although Montgomery et al.⁸⁾ had only obtained a 16% yield of 2-fluoroadenosine **11** on Schiemann reaction of 2-aminoadenosine in aqueous solution, we reacted unprotected 2-aminoadenosine **8** in 60% HF/pyridine with tert.-butyl nitrite and obtained after workup with an icecold suspension of CaCO₃ and filtration of the CaF₂/CaCO₃ slurry 2-fluoroadenosine **11** in ca. 80% yield besides traces of isoguanosine¹¹) **12** and unreacted 2-aminoadenosine **8**, which were readily removed on crystallization from ethanol/ water.



Scheme III

To test the generality of this method, we converted also 2-amino-N⁶,N⁶-pentamethyleneadenosine 13^{5} as well as guanosine 7 into 2-fluoro-N⁶,N⁶-pentamethylene-adenosine 14 and 2-fluoroinosine 15 in ca. 70-80% yield. On attempted recrystallization from ethanol-H₂O, 2-fluoroinosine 15 was partially saponified to xanthosine so that no analytical sample of 15 could as yet be obtained. 2-fluoroinosine 15 has previously been prepared by Schiemann reaction of the O⁶-benzyl derivative of guanosine in aqueous solution followed by hydrogenation of the O⁶-benzyl group in ca. 19% overall yield¹² (Scheme III).

Since tert.-butyl nitrite turned out to be a potential explosive¹³), we subsequently replaced tert.-butyl nitrite by a concentrated aqueous solution of potassium nitrite without any decrease in yield of 2-fluoroadenosine **11**. The use of solid sodium or potassium nitrite is also feasable but the addition of solid salts is, however, more difficult to control and leads to increased foaming.

We anticipate that heterocyclic amino groups in other types of unprotected nucleosides can likewise be converted into the corresponding fluoronucleosides without cleavage of the nucleoside bond.

The further conversion of 2-fluoroadenosine 11 e. g. via its corresponding crystalline¹⁴⁾ 3',5'-O-(1,1,3,3-tetraisopropyldisiloxane-1,3-diyl)-derivative¹⁵⁾⁻²⁰⁾ followed by inversion of the 2'-hydroxy group²¹⁾⁻²⁴⁾, desilylation and 5'-phosphorylation into Fludarabinephosphate (Fludara) 6 is straightforward and is therefore not dealt with here.

EXPERIMENTAL

¹H-NMR spectra were recorded on a Bruker WH 300 instrument. Thin layer (TLC) and column-chromatography were carried out on silica plates (E. Merck) and silica gel No. 10757 E. Merck, containing 40% H₂O using the systems: A: upper phase of n-butanol-acetic acid - H₂O = 4:1:5, System B: upper phase of ethyl-acetate-2-methoxyethanol-H₂O = 4:1:2, System C: ethanol-25% NH₃.

2-Fluoroadenosine 11

To a stirred solution of 15 g (53,14 mmol) 2-aminoadenosine 8 in 50 ml 56% HF/pyridine, which was cooled from +14°C to -11°C, a concentrated solution of 4.9 g KNO₂ (57.6 mmol) in 3 ml H₂O was added dropwise at $-11^{\circ} \rightarrow -6^{\circ}$ C within 1h, whereupon gas evolution was observed. After continued stirring for 1h at $-6^{\circ} \rightarrow -8^{\circ}C$ the solution became turbid and TLC (system A) showed that practically all of the 2-aminoadenosine 8 had reacted. After a further hour at $+3^{\circ}$ C, the reaction mixture was poured on a stirred ice cold slurry of 100 g of powdered CaCO₃ in 200 ml H₂O, containing ca. 0.5 ml n-butanol to suppress any excess foaming. After 3h at +6°C and 16h at +24°C, the slightly basic solution ($P_{\rm H} = 7.5$ -8) was filtered, the CaF₂/CaCO₃ cake washed with 120 ml H_2O and 150 ml ethanol - H_2O (1:1) and the combined filtrates concentrated at 45°C/35-40 mbar, whereupon a voluminous yellowish precipitate of 2-fluoroadenosine 11 formed on standing overnight. After filtration, washing with 50 ml H₂O followed by 50 ml 95% ethanol and finally by 30 ml methyl-tert.-butyl ether, the pink precipitate was dried at 60°C in vacuo to give a first crop of 7.98 g of 11. On concentration of the combined mother liquors three further crops of 1.7 g, 1.7 g and 0.8 g of 11 were obtained (combined yield = 12.18 g = 80.3%), which were homogeneous on TLC (System A, $R_f = 0.63$). TLC of the mother liquor (System A) demonstrated the presence of minor amounts of the starting material 2-aminoadenosine 8 and isoguanosine 12 besides 2-fluoroadenosine 4 as the major spot. The 2-fluoroadenosine 11 gave on recrystallization from ethanol-H₂O with added charcoal a gel-like product, which solidified on washing with ethanol followed by methyl-tert.-butyl ether to give the analytical sample of 11. ¹H-NMR (DMSO-D₆) & 3.50-3.70 (m, 2H, H-5') 3.93 (m, 1 H, H-4'), 4.13 (m, 1H, H-3'), 4.52 (m, 1H, H-2'), 5.78 (d, J = 6 Hz, H-1'), 8.34 (s, 1H, H-8) Anal. calcd for $C_{10}H_{12}N_5O_4F$ (285.25) C, 42.11; H, 4.24; N, 24.55; F, 6.66. Found: C, 41.94; H 4.38; N 24.35; F, 6.77

2-Fluoro-N⁶,N⁶-pentamethyleneadenosine 14

To a solution of 0.7 g (2mmol) of finely powdered 2-amino-N⁶,N⁶-pentamethyleneadenosine **13** in 7 ml 50% HF/pyridine was added 0.412 g (2 mmol) tert-butyl nitrite at -25°C within 5 min with stirring. After 30 min at -25° the reaction mixture was poured on an ice cold and stirred suspension of 25 g CaCO₃. After filtering and washing the powdered CaF₂/CO₃ cake with ethanol-water, evaporation of the filtrate in vacuo gave 0.89 g crude product. This material dissolved in 20 ml methanol and evaporated with 15 g silica gel (40%

SYNTHESIS OF 2-FLUOROPURINE NUCLEOSIDES

H₂O). The resulting mixture was placed on top of a column containing additional 15 g silica gel (40% H₂O) that was then eluted with the upper phase of System B: ethyl acetate-2-methoxyethanol-H₂O = 4:1:2 to give after a 25 ml forerun with the subsequent 125 ml 0.64 g (90%) of yellowish, crystalline 2-fluoro-N⁶,N⁶-pentamethyleneadenosine 14, which was recrystallized from ethyl acetate to afford the analytical sample, mp 164°C. ¹H-NMR (DMSO-D₆) δ : 1.52-1.75 (m, 6H, (CH₂)₃), 3.91-3.98 (m, 1H, H-4'), 4.10-4.18 (m, 1H, H-3'), 4.45-4.52 (m, 1H, H-2'), 5.82 (d, J = 6 Hz, 1H, H-1'), 8.38 (s, 1H, H-8). Anal. calcd for C₁₅H₂₀N₅O₄F (353.37) C 50.99; H 5.7; N 19.82; F 5.38. Found: C 50.79; H 5.85; N 19.67; F 5.30

2-Fluoroinosine 15

1.41 g (5 mmol) guanosine-dihydrate 7 were dissolved in 20 ml 50% HF/pyridine, treated at -25°C with 1.108 g (5 mmol) tert.-butyl nitrite, kept for an additional 60 min at -30°C and worked up with ice cold CaCO₃. After washing the CaF₂/CaCO₃ cake with 150 ml H₂O and evaporating the filtrate in vacuo, the residue (0,9 g), which sometimes crystallizes spontaneously from ethanol-H₂O, was dissolved in 10 ml H₂O and evaporated with 10 g silica gel (40% H₂O). The resulting mixture was placed on top of a column containing additional 20 g silica gel (40% H₂O) that was then eluted with the upper phase of TLC system A n-butanol-acetic acid - H₂O = 4:1:5 the first 50 ml gave rise to impurities. Further elution with 50 ml afforded 0.46 g (80,4%) of <u>15</u>, which was homogenous on TLC (System A, R_f=0.37) after washing with 96% ethanol. After dissolving the substance in ca 4-5 ml H₂O pure 2-fluoroinosine **15** crystallized out, mp 250°C (dec.), and TLC (System C) of the mother liquor indicated besides 2-fluoroinosine (R_f=0.42) the presence of xanthosine (R_f=0.26). ¹H-NMR (DMSO-D₆) δ : 3.91-3.98 (br, H-4'), 4.1-4.21(br, H-3'), 4.48-4,6 (br, H-2'), 5.76-5.85 (br, H-1'), 8.15-8.45 (br, H-8); ¹⁹F-NMR (DMSO-D₆) δ : 110.5 ppm (br) (C₆F₆ as internal standard = 0 ppm); UV (H₂O) λ_{max} 253 nm (11.900)

Acknowledgment: We thank Dr. G. Michl for the measurement and interpretation of the ¹Hand ¹⁹F-NMR-Spectra.

REFERENCES

- 1) Drugs of the Future (1993) 18, 80 and ibid. (1991) 16, 78
- 2) Montgomery, J.A.; Hewson, K.; J. Med. Chem. (1969) 12, 498
- Montgomery, J.A.; Clayton, S.D.; Shortnacy, A.T.; in: Nucleic Acid Chemistry III (1986) 156; edited by L. B. Townsend and R. S. Tipson, John Wiley/New York
- Markovac, A.; LaMontagne, M.P.; Eur. Pat. Appl. EP 461.407, *Chem. Abstr.* (1992) 116, 194798,
- 5) Vorbrüggen, H.; Krolikiewicz, K.; Liebigs Ann. Chem. (1976) 745
- 6) Robins, M.J.; Hansske, F.; Bernier, S.E.; Can. J. Chem. (1981) 59, 3360

- 7) Robins, M.J.; Uznański, B.; Can. J. Chem. (1981) 59, 2608
- 8) Montgomery, J.A.; Hewson, K.; J. Amer. Chem. Soc. (1960) 82, 463
- 9) Montgomery, J.A.; Hewson, K.; J. Org. Chem. (1968) 33, 432
- 10) Robins, M.J.; Uznański, B.; Can. J. Chem. (1981) 59, 2601
- 11) Davoll, J.; J. Amer. Chem. Soc. (1951) 73, 3174
- Gerster, J.F.; Robins, R.K.; J. Amer. Chem. Soc. (1965) 87, 3752; J. Org. Chem. (1966) 31, 3258
- 13) Lopez, F.; Chem. Eng. News (1992), Dec. 21, 2
- 14) 2-Fluoroadenosine 11 was reacted with TIPS-Cl in pyridine to give after evaporation and filtration of the crude product in CH₂Cl₂ over a short column of SiO₂ the crystalline TIPS-derivative of 11 in 80% yield, mp 210°C (from methanol).
- 15) Markiewicz, W.T.; J. Chem. Res. (1979), (S) 24
- 16) Fukukawa, K.; Uedao, T.; Hirano, T.; Chem. Pharm. Bull. (1983) 31 (5), 1582
- Biggadike, K. Borthwick. A.D.; Evans, D.; Exall, A.M.; Kirk, B.E.; Roberts, S.M.; Stephenson, L.; Youds, P.; Slawin, A.M.Z.; Williams, D.J.; J. Chem. Soc., Chem. Comm. (1987) 251
- 18) Marriott, J.H.; Mottahedeh, M.; Reese, C.B.; Carbohyd. Res. (1991) 216, 257
- Matsuda, A.; Takenuki, K.; Tanaka, M.; Sasaki, T.; Ueda, T.; J. Med. Chem. (1991)
 34, 812
- Robins, M.J.; Hawrelak. S.D.; Hernández, A.E.; Wnuk, S.F.; Nucleosides & Nucleotides (1992) 11, 821
- 21) Fukukawa, K; Ueda, T.; Hirano, T.; Chem. Pharm. Bull. (1981) 29, 597
- 22) Samano, V.; Robins, M.J.; J. Org. Chem. (1990) 55, 5186
- 23) Lubineau, A.; Bienaymé, H.; Carbohyd. Res. (1991) 212, 267
- 24) Binkley, R.W.; J. Org. Chem. (1991) 56, 3892

Received 9/7/93 Accepted 10/26/93