

A CONVENIENT METHOD FOR THE CONVERSION OF
trans to cis-CINNAMIC ACIDS

Vilmos Galamb and Howard Alper*

Ottawa-Carleton Institute for Research and Graduate Studies
in Chemistry, Department of Chemistry, University of Ottawa, Ottawa, Ontario, Canada K1N 9B4

Summary: trans-Cinnamic acids react in sequence with bromine, base, and carbon monoxide [palladium(0) and phase transfer catalysis] to give cis-acids in reasonable yields.


The conversion of trans to cis-cinnamic acids has attracted attention over many years. For example, irradiation of trans (or cis)-cinnamic acid for 2.5 - 120 hr. affords an equilibrium mixture from which the cis-isomer can be obtained in 10-40% yields^{1,2}. Another approach involves esterification of the E-acid followed by bromination, base-induced elimination to the arylpropionic acid and subsequent hydrogenation affording the Z-cinnamic acid in 28-31% overall yield from the ester (the yield from the acid was not reported)³. Other unattractive routes involving ester substrates include the use of derivatives of phenylpropionic acid^{4a} and phenylalanine (i.e., diazo esters)^{4b}.

The use of vinylic bromides as intermediates has also been examined. Bromination of the trans-cinnamic acid followed by treatment with sodium bicarbonate in acetone gave the cis-bromostyrene. Grignard formation from the latter, followed by conventional carbonation and protonation afforded cinnamic acid in a 2:1 cis/trans ratio. Lithiation of cis-bromostyrene and subsequent carbonation and work-up gave a 4:1 mixture of trans-cinnamic acid and phenylpropionic acid⁵.

Phase transfer catalysis is an excellent means of effecting metal catalyzed processes under gentle conditions^{6,7}. For instance, benzylic halides are carbonylated to acids, at room temperature and atmospheric pressure in the presence of cobalt carbonyl^{6,7} or tetrakis(tri-phenylphosphine)palladium⁸, sodium hydroxide, methylene chloride, and a phase transfer agent. We now wish to report that cis-bromostyrenes (3), obtained from trans-cinnamic acids (1) as illustrated in Scheme 1, can be carbonylated to cis-acids (4) using palladium (0) and phase transfer catalysis conditions.

Bromination of a variety of trans-acids, 1 [R=H, p-Cl, p-CH₃, p-CF₃, m-Br, o-CH₃] in carbon tetrachloride gave the dibromoacids (2) in 90-97% yields (Table 1). The cis-bromides (3) were formed in 86-91% yield by exposure of 2 to bicarbonate ion in acetone⁹. When 3 was reacted with carbon monoxide, 5N sodium hydroxide, t-amyl alcohol as the organic phase (and some methylene chloride to solubilize Pd(PPh₃)₄), benzyltriethylammonium chloride as the phase transfer catalyst and tetrakis(triphenylphosphine)palladium(0) as the metal catalyst [25-40 / 1 mole ratio of 3 / Pd(PPh₃)₄], the pure cis-cinnamic acids (4) were obtained in 35-56%

Scheme 1



yields (not optimized). By-products of some of these reactions were the *cis,cis*-1,4-diaryl-1,3-butadienes (**5**) and very small amounts of the *t*-amyl ester of **4** were also sometimes isolated.

Note that, in contrast to benzylic halides⁸, the nature of the palladium(0) catalyst does not change the reaction course as bis[bis(1,2-diphenylphosphino)ethane]palladium(0) [$\text{Pd}(\text{diphos})_2$] behaved in the same manner as tetrakis(triphenylphosphine)palladium(0) [for **3**, $\text{R}=\text{p-Cl}$]. A phase transfer agent is required for these reactions, since the yield of **4** is much lower in the absence of the quaternary ammonium salt. No reaction occurs in the absence of the palladium catalyst.

TABLE 1
Product yields^a

<u>1</u> , R=	<u>2</u> , %	<u>3</u> , %	<u>4</u> , % ^b	Overall yield, %	M.p., °C	lit. m.p. °C	Other products
H	93	89	56	47	66-67	68 ¹	<u>5</u> , trace
p-Cl ^c	94	91	47	41	114-115	111-111.5 ²	<u>5</u> , 16.6%
m-Br	91	89	35	28	75.5-77.0	75-77 ²	<u>5</u> , 17.1%
p-CH ₃	97	87	49	41	73-75	77.5-79.0 ¹	<u>5</u> , 8.4%
p-CF ₃	90	91	48	40	53.0-53.5 ^d	-	<u>5</u> , 21.4%
o-CH ₃	94	86	54	44	91-92	89-90 ⁴	-
p-CH ₃ O	90	- ^e	-	-	-	-	-

^aYields are of pure materials. ^b Using Pd(PPh₃)₄ as the catalyst. ^cYield of 4, R=p-Cl, was 51% using Pd(diphos)₂ as the catalyst. ^dNew compound. Anal. calcd. for C₁₀H₇F₃O₂: C, 55.56; H, 3.26. Found: C, 55.95; H, 3.16. ^eOnly the trans-isomer of 3, R=p-CH₃O was formed.

The reaction is apparently insensitive to substituent effects since essentially the same yields of 4 were realized using 3, R=p-CF₃, or 3, R=p-CH₃ as reactants. The use of the less polar solvent, 4-methyl-2-pentanone, as the organic phase give superior yields of 4. However, the reaction is not useful, since a mixture of cis and trans acids are produced (e.g., 3, R=H afforded cinnamic acid in 79% yield, with the cis/trans ratio being 62:38). Isomeric mixtures of acids, in high yield, were also isolated when benzene was employed as the organic phase.

The overall yield for the described conversion of trans to pure cis-cinnamic acids are reasonable (28-47%). In addition, this reaction sequence is simple both in execution and work-up. The palladium catalyzed reaction is regiospecific, in terms of the formation of either 4 or 5 [i.e., no geometric isomers].

The following general procedure was used: a mixture of 5N sodium hydroxide [15 ml.] and benzyltriethylammonium chloride [0.5-0.6 mmol] was stirred for 30 minutes under nitrogen. The palladium(0) catalyst [0.09-0.12 mmol] in t-amyl alcohol (15 ml.) containing methylene chloride (4-6 ml.), was added and the reaction atmosphere was changed to carbon monoxide. After stirring for 3.5-4.0 hours, 3 [2.75-3.75 mmol] in t-amyl alcohol (5 ml.) was added dropwise and then the mixture was vigorously stirred overnight (CO atmosphere). The phases were separated, the aqueous phase was extracted with ether (2 x 15 ml.), and the organic phase was washed with water (20 ml.). The combined aqueous phases were cooled, acidified (c. HCl), extracted with ether (4 x 50 ml.), and the ether extracts were dried (MgSO₄) and concentrated. Analytically pure acid (4) was then obtained by thin-layer chromatography.

Acknowledgement

We are indebted to the Natural Sciences and Engineering Research Council for support of this work.

References

1. T.A. Wittstruck and E.N. Trachtenberg, J. Am. Chem. Soc., 89, 3803 (1967).
2. E.N. Ugochukwu and R.L. Wain, Ann. Appl. Biol., 61, 121 (1968).
3. E.R. Trumbull, R.T. Finn, K.M. Ibne-Rasa and C.K. Sauers, J. Org. Chem., 27, 2339 (1962).
4. (a) J. Basters, C.J. Groenenboom, H. van Bekkum and L.L. van Reijen, Rec. Trav. Chim., 92, 219 (1973); (b) N. Takamura, T. Mizoguchi and S. Yamada, Chem. Pharm. Bull., 23, 299 (1975).
5. G.F. Wright, J. Org. Chem., 1, 457 (1936).
6. H. Alper, Adv. Organometal. Chem., 19, 183 (1981).
7. H. Alper, Fundamental Research in Homogeneous Catalysis, in press.
8. H. Alper, K. Hashem, and J. Heveling, Organometallics, 1, 775 (1982).
9. S.J. Cristol and W.P. Norris, J. Am. Chem. Soc., 75, 2645 (1953).

(Received in USA 8 April 1983)