Tetrahedron Letters 50 (2009) 3693-3697

Contents lists available at ScienceDirect

Tetrahedron Letters

journal homepage: www.elsevier.com/locate/tetlet



Negligible diradical character for the ultralong C–C bond in 1,1,2,2-tetraarylpyracene derivatives at room temperature

Takashi Takeda^a, Hidetoshi Kawai^{a,b}, Rainer Herges^c, Eva Mucke^c, Yoshitaka Sawai^a, Kei Murakoshi^a, Kenshu Fujiwara^a, Takanori Suzuki^{a,*}

^a Department of Chemistry, Faculty of Science, Hokkaido University, Sapporo 060-0810, Japan

^b PRESTO, Japan Science and Technology Agency, Saitama, Japan

^c Insitut für Organische Chemie, Universität Kiel, Otto-Hanh-Platz 4, D-24098 Kiel, Germany

ARTICLE INFO

Article history: Received 15 January 2009 Revised 23 March 2009 Accepted 26 March 2009 Available online 2 April 2009

In memory of Professor Fumio Toda (1933-2008)

Keywords: Ultralong bond Strained molecule Steric repulsion Thermochromism Diradical X-ray analysis

ABSTRACT

There is evidence that the crystalline-state thermochromic behavior of dispiropyracene **acr-1** with an ultralong C–C bond [1.791(3) Å at 413 K, 1.771(3) Å at 93 K] might be due to thermal generation of a bond-dissociated triplet diradical. The C_1 – C_2 bond lengths (*d*) in the newly prepared 1,1,2,2-tetraarylpyracenes **1a–e** [1.717(4)–1.761(4) Å at 113/123/153 K] are also much larger than the standard value for $C(sp^3)$ – $C(sp^3)$ (1.54 Å). The fact that there is no correlation between *d* and the radical-stabilizing parameter (σ^{\bullet}) provides evidence that there is no contribution from a crystallographic artifact caused by contamination of the diradical in the crystal of **1** at room temperature or below. Further indication for the covalent nature of the ultralong C–C bond in tetraphenylpyracene **1c** was attained by Raman spectroscopy, showing the far red-shifted C_1 – C_2 stretching vibration (638 cm⁻¹) and by the theoretical prediction of very large vertical singlet–triplet energy gap (58.3 kcal mol⁻¹ at the UB3LYP/6-31G^{*} level). However, at elevated temperatures bond dissociation might occur forming a triplet diradical.

© 2009 Elsevier Ltd. All rights reserved.

The C–C bond length (*d*) is one of the most fundamental parameters in organic chemistry,¹ and a *d* value of 1.54 Å is assumed as the standard length of a C(sp³)–C(sp³) single bond. This explains why the experimentally determined *d* can be used to assess the π -bond order in combination with other standard values [1.33 Å for C(sp²)=C(sp²); 1.20 Å for C(sp)=C(sp)]. On the other hand, there have been several reports in which the C–C single bond is shortened or elongated independent of the bond order,² and such elongation is often accompanied by a strained molecular structure and/or severe steric hindrance.³

Hexaphenylethane (HPE) is a good example of this phenomenon; its central C–C bond is elongated $[1.67(3) Å]^4$ primarily due to steric repulsion among six aryl substituents. According to the proposed linear relationship between *d* and bond-dissociation energy (BDE),⁵ a smaller BDE is expected for a longer C–C bond. In fact, some isolable HPE derivatives exist as equilibrated mixtures containing bond-dissociated trityl radicals in solution. On the other hand, 'cross-clamped' or 'condensed' derivatives of HPE show higher thermodynamic stability because dimerization is switched from an intermolecular to an intramolecular process. Thus, the HPEs with an even longer C–C bond (d > 1.70 Å) could be isolated [1.712(5)-1.734(5) Å for 3,8-diha-lo-1,1,2,2-tetraphenylnaphthocyclobutenes by Toda's group⁶ and 1.713(2) Å for a multi-benzannulated caged hydrocarbon by one of us⁷].

Recently, we also reported^{8a,b} that a pyracene-based HPE derivative **acr-1** with two spiro(10-methylacridan) units has an ultralong C–C bond [d = 1.771(3)Å at 93 K], which is one of the longest ever reported⁹ [the d values for other crystallographically independent molecules of acr-1 are 1.758(3), 1.712(2), and 1.707(2) Å.]. The above X-ray analysis was carried out with special care, however, there still remains some possibility that the concomitant presence of a bond-dissociated diradical might have led to the assignment of a larger d than the intrinsic value.^{10,13} Our major concern in this Letter is to shed light on the following points: (a) does a bond-dissociated diradical exist or can it be generated in the crystal of HPEs with an ultralong C-C bond; and (b) is the crystallographically determined very large d (> 1.75 Å) be intrinsic or an artifact by the concomitant presence of a bond-dissociated diradical (or diradicaloid) in the crystal.



^{*} Corresponding author. Tel./fax: +81 11 706 2714. E-mail address: tak@sci.hokudai.ac.jp (T. Suzuki).

^{0040-4039/\$ -} see front matter \odot 2009 Elsevier Ltd. All rights reserved. doi:10.1016/j.tetlet.2009.03.202



Here we found the reversible thermochromic behavior¹⁴ of **acr**-1 in the single-crystalline state, which might be related to the bond dissociation of the ultralong bond forming the corresponding diradical in the crystal, as shown in the first part of this Letter. Next, we have conducted the X-ray analyses of acr-1 at several elevated temperatures to see if generation/increased contribution of the diradical at higher temperatures expands the experimentally determined *d* value of **acr-1**. Since the marginal expansion of 0.020 Å over a temperature difference of 320° does not provide decisive support for the diradical contribution, we have then turned our attention to the studies on a new series of HPEs, 1,1,2,2-tetraarylpyracenes **1**. The experimentally assigned d value should change with the attachment of radical-stabilizing/non-stabilizing substituents when the coexistence of a bond-dissociated diradicals affects the apparent bond lengths determined by X-ray analyses (Scheme 1). If the corresponding diradicals exist in equilibrium with HPEs in the crystal,¹⁴ compounds with radical-stabilizing substituents would exhibit a larger value of *d* than those with non-stabilizing or radical-destabilizing substituents. The low-temperature X-ray structures of the corresponding compounds do not reveal such a relationship. The details of our results will be shown in the last part of this Letter.

First, we noticed a color change of crystalline samples of acr-1 when they were heated in an ESR tube sealed with Ar. Similar thermochromic behavior was observed by heating the single crystalline sample on a glass fiber (Fig. 1), where temperature control was carried out by a N₂ gas flow. At 273 K, the sample is pale yellow, whose color gradually turned to dark red (413 K) via orange (333 K). This observation is not accompanied by decomposition of **acr-1** because the crystal returned to pale yellow upon cooling to room temperature or 123 K, which still gave X-ray diffraction data identical to those obtained before the heat treatment. A reversible phase transition or bond fission to generate a certain population of diradicals upon heating might be responsible for the observed thermochromic behavior. Thus, the X-ray analysis of acr-1 was carried out at several temperatures above 273 K.^{15,16} No phase transition occurs in the range between 273 K and 413 K, thus the observed thermochromic behavior is not related to a phase transition.

The X-ray structure at 413 K is quite similar to those obtained at lower temperatures except for the enlarged thermal ellipsoids at



Figure 1. Single-crystalline thermochromic behavior of acr-1.

most atoms. It is noteworthy that those for C_1 and C_2 are still small and round shaped (Fig. 2). The difference electron density map indicates the presence of bonding electron between C_1 and C_2 , (Fig. S2) as in the case of the X-ray analysis at 93 K.^{8a} The *d* value for the bond in question is 1.791(3) Å, which is only 0.020 Å longer than that measured at 93 K.¹⁷ The X-ray analyses at 333 K and 373 K gave the results with the intermediary structures with the *d* value of 1.776(3) Å and 1.786(3) Å, respectively. As shown by Toda's and Siegel's groups, 3,8-dichloro-1,1,2,2-tetraphenylnaphthocyclobutene exhibits a similar degree of bond expansion [1.710(2) Å at 90 K,^{6e} 1.720(4) at 298 K^{6a}].

The vertical singlet-triplet energy gap (ΔE_{S-T}) is a frequently used indicator to evaluate the diradical character of molecules.¹⁸ We performed calculations at the UB3LYP/6-31G⁻-level of DFT based on the X-ray geometry of **acr-1** at 413 K (fixed d of 1.7922 Å) and obtained the ΔE_{S-T} value of 58.99 kcal mol⁻¹. A quite similar value (59.04 kcal mol⁻¹) was obtained for the crystallographically independent molecule with the shortest C_1-C_2 bond measured at 93 K (d = 1.7062 Å). The very large ΔE_{S-T} values indicate that the diradical contribution to the overall wavefunction of **acr-1** is negligible at lower temperatures (compare with the ΔE_{S-T} of o-, m-, and p-benzyne: 37.5, 21.0 3.8 kcal/mol).¹⁹ On the other hand, we found a triplet minimum structure with a very large bond distance d of 3.110 Å which energetically is only 8.14 kcal mol⁻¹ above the singlet structure (UB3LYP/6-31G⁻¹, Fig. S3). This species could explain the observed thermochromism on a theoretical basis. According to TD DFT calculations the longest wavelength absorption of the singlet state is 362 nm which is in agreement with the fact that the crystals are almost colorless at low temperatures. By contrast the triplet state exhibits several



X = radical stabilizing / non-stabilizing substituent

observed as longer bond?



Figure 2. ORTEP drawings of acr-1 (top view and side view) at 413 K. Note the small and round-shaped ellipsoids for C_1 and C_2 .

absorptions within the visible range (strongest bands at 450 and 475 nm; Table S1) which explain the red color of the crystals at higher temperatures at which the triplet state will be increasingly populated. Thus, it is clear that the ultralong C–C bond in **acr-1** at room temperature and below is intrinsic and that the concomitant presence of bond-dissociated triplet diradical becomes important only at higher temperatures.

To gain further insight into the nature of the ultralong bonds we set out to crystallographically measure the bond length *d* in HPEs other than **acr-1**. For this purpose, we designed 1,1,2,2-tetraaryl-pyracene derivatives **1a–e** as suitable HPE derivatives from the following viewpoints: first, the pyracene skeleton should act as a scaffold to induce enough 'front strain'⁴ to ensure the presence of an extremely long C_1-C_2 bond (d > 1.70 Å), as shown by **acr-1**; second, the system should be designed in such a way that the bond-dissociated species is a diradical and not (as, e.g., in the case of the benzocyclobutenes-type HPEs) a closed shell structure (*o*-quinodimethane),²⁰ and third, a series of compounds with various substituents on aryl groups should be accessible via the corresponding dicationic precursors.²¹

We have prepared here the parent tetraphenylpyracene $(1c)^{22}$ from 5,6-dibromoacenaphthene and benzophenone in three steps via pyran **3c**, as shown in Scheme 2. By starting with the substituted benzophenones, HPEs with benzyl radical-stabilizing substituents on phenyl rings such as 4-^tBu (1a), and 4-Me (1b) could be similarly prepared along with those with non-stabilizing [3,5-Me₂ (1d)] and destabilizing [4-F (1e)] substituents. The desired HPEs (1) were isolated as surprisingly stable colorless/pale yellow solids despite the very large *d* value for the C_1 – C_2 bond (vide infra). Another series of HPEs **2** with an acenaphthene π -framework was also designed to further support the conclusion obtained from **1**. The structurally related acenaphthene derivative **acr-2**^{8a,8c} has a shorter bond length [1.696(3)Å] than **acr-1** because it adopts a skewed arvlene framework to relieve steric repulsion between two acridan units. Yet, the skeleton might be still suitable enough to test the validity of our approach adopted here. So. 1.1.2.2-tetraarvlacenaphthenes $2a(4-^{t}Bu).b(4-Me).d(3.5-Me_{2}).e(4-F)$ were prepared over three steps from 1,8-diiodonaphthalene and the corresponding benzophenones via pyrans **4** (Scheme 3).²² The preparation and X-ray structure of tetraphenylacenaphthene (2c) were reported previously by Gabbaï and coworker.23

After considerable effort, we finally obtained a single-crystal specimen of high quality for all nine kinds of HPEs 1a-e/ **2a,b,d,e**.¹⁶ Low-temperature X-ray analyses revealed that all of them have an extremely long C_1-C_2 bond [1.717(4)-1.761(4) Å]for **1** and 1.694(3)–1.708(4) Å for **2**] (Tables 1 and 2).²⁴ No structural disorder around the C_1 - C_2 bond was found in any case. As represented by 4-Me derivative **1b** [d = 1.717(4)Å: the smallest among **1a–e**] and 4-F derivative **1e** [d = 1.761(4)Å: the largest] (Fig. 3), the ORTEP drawings show no anomalies in thermal ellipsoids of the long C_1 - C_2 bond or in electron density maps. The radical-stabilizing parameters $(\sigma^{\bullet})^{25}$ are also summarized in Table 1, which quantify the thermodynamic stability of the benzyl radicals attached with a various substituent on the benzene nucleus. As shown by these data, HPEs with radical-stabilizing groups (1a,b/ **2a.b**: $\sigma^{\bullet} > 0$) do not necessarily have larger d values than the parent compound (**1c/2c**) or those with non-stabilizing/destabilizing groups (**1d.e**/**2d.e**: $\sigma^{\bullet} < 0$). Thus, the radical-stabilizing substituents do not increase the apparent bond length determined by X-ray, which leads us to conclude that the contribution from the bond-dissociated diradical is negligible for the present HPEs 1/2



a : Ar = $4^{-t}BuC_6H_4$; **b** : Ar = $4^{-}MeC_6H_4$; **c** : Ar = C_6H_5 ; **d** : Ar = $3,5^{-}Me_2C_6H_3$; **e** : Ar = $4^{-}FC_6H_4$

Scheme 2. Preparation of tetraarylpyracenes 1a-e. Reagents: (a) (1) *n*-BuLi, then Ar₂CO, ether; (2) trifluoroacetic acid, CH₂Cl₂; (b) TMSClO₄, 1,1,1,3,3,3-hexafluoro-2-propanol; (c) Zn, MeCN.



a : Ar = $4^{-t}BuC_6H_4$; **b** : Ar = $4-MeC_6H_4$; **c** : Ar = C_6H_5 ; **d** : Ar = $3,5-Me_2C_6H_3$; **e** : Ar = $4-FC_6H_4$

Table 1

Structural parameters of pyracene-type HPEs 1 determined by low-temperature X-ray analyses and the radical-stabilizing parameters σ^{\bullet}

Compound	1a	1b	1c	1d	1e
X	4- ^t Bu	4-Me	H	3,5-Me ₂	4-F
d (Å)	1.749(4)	1.717(4)	1.754(2)	1.743(2)	1.761(4)
σ•	0.036	0.015	0.000	-0.002	-0.011

Table 2

Structural parameters of acenaphthene-type HPEs **2** determined by low-temperature X-ray analyses

Compound	2a	2b	2c ^a	2d	2e ^b
X	4- ^t Bu	4-Me	H	3,5-Me ₂	4-F
d (Å)	1.708(4)	1.694(3)	1.701(3)	1.701(2)	1.707(2) 1.705(2)

^a Ref. 23.

^b Two crystallographically independent molecules.



Figure 3. ORTEP drawings of 1b (left) and 1e (right) at 123 K.

in the crystal state even for derivatives with an ultralong C–C bond with a d value as large as 1.76 Å.

For further validation of the crystallographically determined extremely long C-C bond, Raman spectroscopy was applied on the crystalline samples of tetraphenyl derivatives (1c/2c). The stretching vibration of the long C_1 - C_2 bond studied by Raman spectroscopy at 298 K gives direct information on the force constant of the bond. The absorption in pyracene derivative 1c with the highest intensity (expt. 638 cm⁻¹) in Figure 4a corresponds to the normal mode with the largest amplitude along the trajectory of the stretching vibration as verified by DFT calculation (B3LYP/6-31G^{*}: calcd 650 cm⁻¹). There are other absorptions assigned as normal modes that include C_1-C_2 stretching vibration (470, 621, 718, 812 cm⁻¹), for which the calculation well reproduced the frequencies (478, 637, 720, 825 cm⁻¹) (Fig. 4b). The observed redshifted value compared to the C-C stretching vibration of ethane itself (995 cm⁻¹) clearly shows that the ultralong C–C bond $[d_{\text{expt}} = 1.754(2) \text{ Å}; d_{\text{calcd}} = 1.759 \text{ Å}]$ in **1c** has a very small force constant.

Not only the frequencies of the C_1-C_2 stretching bands but also the other experimental spectral lines were well reproduced by the calculation. In contrast to the X-ray analysis and NMR, Raman is a very fast spectroscopic method. In a hypothetical equilibrium of a closed shell species with a large *d* value and a diradical, both species should appear with separate absorption bands and not as a time-averaged spectrum like in NMR. The agreement of the experimental and theoretical Raman spectrum is very good, and there are no additional lines in the experimental spectrum, which could



Figure 4. Experimental (a) and calculated (b) Raman spectra of 1c.

not be assigned to the closed shell structure. Therefore, at room temperature or below a diradical structure can either be completely excluded or, if there is an equilibrium, it must be present in very little concentration in the crystalline sample of **1c**.²⁶ Raman investigations at higher temperatures are in progress.

In conclusion, we have no reason to assume that the extremely long (d > 1.70 Å) or ultralong (d > 1.75 Å) C–C bond in the present HPEs gives rise to a noticeable diradical character of the corresponding compounds. However, there is indication that there is a triplet diradical in a thermal equilibrium with the closed shell structure at higher temperatures in crystalline **acr-1**. Despite the large expansion that may lower the BDE, the present HPEs exhibit surprising chemical stability: For example, the parent **1c** [d = 1.754(2) Å] can be purified by chromatography using aerated solvents and can be stored infinitely as solids. It remains intact upon treatment with Et₃SiH or Bu₃SnH. Based on our results, studies to achieve further elongated C–C bonds are warranted with a d value even longer than the shortest nonbonded C–C contact (1.80 Å).²⁷

Acknowledgments

This work was partly supported by the Global COE Program (Project No. B01: Catalysis as the Basis for Innovation in Materials Science) from MEXT, Japan. T.T. thanks JSPS research fellowship for young scientists (18-4425).

Supplementary data

Crystallographic data (excluding structure factors) for the structures in this Letter has been deposited with the Cambridge Crystallographic Data Centre as supplementary publication numbers CCDC 704958–704964, 715129, 715130, 715701–715703). Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK [fax: +44 (0)-1223-336033 or e-mail: deposit@ccdc.cam.ac.uk]. Spectral data of new HPEs (**1a–e/2a,b,d,e**), ORTEP drawings of **1a,c,d/2a,b,d,e** (Fig. S1), the *D* map of **acr-1** at 413 K (Fig. S2), plots of singlet (d = 1.776 Å) and triplet (d = 3.110 Å) species obtained by DT-DFT calculation on **acr-1** (Fig. S3), and calculated absorption bands of both singlet and triplet species of **acr-1** (Table S1). Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.tetlet.2009.03.202.

3696

References and notes

- Allen, F. H.; Kennard, O.; Watson, D. G.; Brammer, L.; Orpen, A. G.; Taylor, R. J. Chem. Soc., Perkin Trans 2 1987, S1.
- (a) Kaupp, G.; Boy, J. Angew. Chem., Int. Ed. **1997**, 36, 48–49; (b) Siegel, J. S. Nature **2006**, 439, 801–802; (c) Tanaka, M.; Sekiguchi, A. Angew. Chem., Int. Ed. **2005**, 44, 5821–5823; (d) Huntley, D. R.; Markopoulos, G.; Donovan, P. M.; Scott, L. T.; Hoffmann, R. Angew. Chem., Int. Ed. **2005**, 44, 7549–7553.
- (a) Suzuki, T.; Takeda, T.; Kawai, H.; Fujiwara, K. Pure Appl. Chem. 2008, 80, 547– 553; (b) Suzuki, T.; Takeda, T.; Kawai, H.; Fujiwara, K. In Strained Hydrocarbon: Ultralong C-C Bond; Dodziuk, H., Ed.; Wiley-VCH, 2009, pp 70–82.
- 4. Kahr, B.; Engen, D. V.; Mislow, K. J. Am. Chem. Soc. 1986, 108, 8305-8307.
- 5. Zavitsas, A. A. J. Phys. Chem. A 2003, 107, 897-898.
- (a) Toda, F.; Tanaka, K.; Stein, Z.; Goldberg, I. Acta Crystallogr., Sect. C 1996, 52, 177–180;
 (b) Toda, F.; Tanaka, K.; Watanabe, M.; Tamura, K.; Miyahara, I.; Nakai, T.; Hirotsu, K. J. Org. Chem. 1999, 64, 3102–3105;
 (c) Tanaka, K.; Takamoto, N.; Tezuka, Y.; Kato, M.; Toda, F. Tetrahedron 2001, 57, 3761–3767;
 (d) Toda, F.; Tanaka, K.; Takamoto, N. Tetrahedron Lett. 2001, 42, 7979–7982;
 (e) Baldridge, K. K.; Kasahara, Y.; Ogawa, K.; Siegel, J. S.; Tanaka, K.; Toda, F. J. Am. Chem. Soc. 1998, 120, 6167–6168.
- Kammermeier, S.; Jones, P. G.; Herges, R. Angew. Chem., Int. Ed. 1997, 36, 1757– 1760.
- (a) Kawai, H.; Takeda, T.; Fujiwara, K.; Wakeshima, M.; Hinatsu, Y.; Suzuki, T. *Chem. Eur. J.* 2008, 14, 5780–5793; (b) Kawai, H.; Takeda, T.; Fujiwara, K.; Inabe, T.; Suzuki, T. *Cryst. Growth Des.* 2005, *5*, 2256–2260; (c) Kawai, H.; Takeda, T.; Fujiwara, K.; Suzuki, T. *Tetrahedron Lett.* 2004, *45*, 8289–8293.
- The bond of 1.781(15) Å in a silabicyclobutane derivative had been the longest among the non-ionic organic molecules before this Letter: Fritz, G.; Wartanessian, S.; Matern, E.; Hönle, W.; Schnering, H. G. v. Z. Anorg. Allg. Chem. 1981, 475, 87–108.
- 10. In some earlier examples, the coexistence of a bond-dissociated valence isomer caused the misassignment of a too-large *d* value (Ref. 11), since X-ray structure only gives the time- and position-averaged geometry of molecules in the crystal. The true value of those cases was later proven to be much smaller by low-temperature data collection and the careful examination of thermal ellipsoids (Ref. 12).
- 11. Ehrenberg, M. Acta Crystallogr. 1966, 20, 182–186.
- (a) Battersby, T. R.; Gantzel, P.; Baldridge, K. K.; Siegel, J. S. Tetrahedron Lett. 1995, 36, 845–848; (b) Harada, J.; Ogawa, K.; Tomoda, S. Chem. Lett. 1995, 24, 751–752.
- 13. One of the reviewers commented that more than 10% of diradical content is necessary to observe the apparent bond expansion. Another reviewer, in contrast, assumes that only few % of diradical contamination is enough to produce the crystallographic artifact. The discrepancy of these experts' claims suggests that there has been no established discussion on this matter, for which this Letter can provide the important experimental results.

- Morita, Y.; Suzuki, S.; Fukui, K.; Nakazawa, S.; Kitagawa, H.; Kishida, H.; Okamoto, H.; Naito, A.; Sekine, A.; Ohashi, Y.; Shiro, M.; Sasaki, K.; Shiomi, D.; Sato, K.; Takui, T.; Nakasuji, K. *Nat. Mater.* **2008**, *7*, 48–51.
- 15. Variable temperature X-ray analysis of acr-1 was previously carried out for the temperature range between 93 K and 233 K. (Ref. 8b). Those measurements revealed that the phase transition actually occurred at 233 K, whereby four crystallographically independent molecules at lower temperatures are merged into two crystallographically independent molecules. No further phase transition occurred at higher temperature up to 413 K.
- Crystal data and details on crystallographic studies of 1a-e/2a,b,d,e (at 113/ 123/153 K) were given in Supplementary data.
- 17. In another crystallographically independent molecule, the C_1-C_2 bond length is 1.719(3) Å; whose expansion degree (0.012 Å) over 320° is similar to the one with the ultralong C–C bond.
- 18. Li, X.; Paldus, J. J. Chem. Phys. 2008, 129, 174101.
- Wenthold, P. G.; Squires, R. R.; Lineberger, W. C. J. Am. Chem. Soc. 1998, 120, 5279–5290.
- 20. Cycloreversion of 1,1,2,2-tetraphenylbenzocyclobutene to 7,7,8,8,tetraphenylo-quinodimethane at low temperature shows that the conversion to the thermodynamically stable bond-dissociated valence isomer is a very rapid process: Quinkert, G.; Wiesdorff, W.-W.; Finke, M.; Opitz, K.; von der Haar, F.-G. *Chem. Ber.* **1968**, *101*, 2302–2325; Tetraaryl-9,10-phenanthraquinodimathane was isolated as a stable compound which no longer isomerises into the corresponding HPE derivative (tetraarylphenanthracyclobutene): Iwashita, S.; Ohta, E.; Higuchi, H.; Kawai, H.; Fujiwara, K.; Ono, K.; Takenaka, M.; Suzuki, T. *Chem. Commun.* **2004**, 2076–2077.
- (a) Suzuki, T.; Nishida, J.; Tsuji, T. Angew. Chem., Int. Ed. **1997**, 36, 1329–1331;
 (b) Suzuki, T.; Ohta, E.; Kawai, H.; Fujiwara, K.; Fukushima, T. Synlett **2007**, 851–869.
- Spectral data of new compounds 1a-e/2a,b,d,e were given in Supplementary data.
- 23. Wang, H.; Gabbaï, F. P. Org. Lett. 2005, 7, 283-285.
- 24. Some variation in *d* was observed in the HPE series of both **1** and **2**. The larger *d* value for the parent (**1c**) than the 3,5-Me₂ derivative (**1d**) clearly shows that the steric bulkiness of substituents on the aryl groups have nothing to do with the C_1-C_2 bond. We are currently examining the electronic effects of the substituents affecting the C_1-C_2 bond length (*d*) through the high-level theoretical calculations, whose results will be reported in due course.
- 25. Dust, J. M.; Arnold, D. R. J. Am. Chem. Soc. 1983, 105, 1221-1227.
- 26. This is also the case for acenaphthene derivative **2c**, which has a shorter C_1-C_2 bond [d = 1.701(3) Å (Ref. 23)]. Its Raman spectrum measured on the crystalline sample has an intense band at 655 cm⁻¹ (calcd 665 cm⁻¹ by B3LYP/6-31G⁻), which is less red shifted than in **1c**. This is clear evidence that **1c** has a bond with a smaller force constant than **2c**, which is consistent with the crystallographic data shown above.
- Adcock, J. L.; Gakh, A. A.; Pollitte, J. L.; Woods, C. J. Am. Chem. Soc. 1992, 114, 3980–3981.