

Total Synthesis of (–)-Lasonolide A

Barry M. Trost, Craig E. Stivala, Daniel R Fandrick, Kami L. Hull, Audris Huang, Caroline Poock, and Rainer Kalkofen J. Am. Chem. Soc., Just Accepted Manuscript • Publication Date (Web): 22 Aug 2016 Downloaded from http://pubs.acs.org on August 22, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Total Synthesis of (-)-Lasonolide A

Barry M. Trost,* Craig E. Stivala, Daniel R. Fandrick, Kami L. Hull, Audris Huang, Caroline Poock, Rainer Kalkofen

Department of Chemistry, Stanford University, Stanford, California 94305-5080

ABSTRACT: The lasonolides are novel polyketides that have displayed remarkable biological activity in vitro against a variety of cancer cell lines. Herein we describe our first generation approach to the formal synthesis of lasonolide A. The key findings from these studies ultimately allowed us to go on and complete a total synthesis of lasonolide A. The convergent approach unites two highly complex fragments utilizing a Ru-catalyzed alkene-alkyne coupling. This type of coupling typically generates branched products, however through a detailed investigation we are now able to demonstrate that subtle structural changes to the substrates can alter the selectivity to favor the formation of the linear product. The synthesis of the fragments features a number of atom economical transformations which are highlighted by the discovery of an engineered enzyme to perform a dynamic kinetic reduction of a b-ketoester to establish the absolute stereochemistry of the southern tetrahydropyran ring with high levels of enantioselectivity.

INTRODUCTION

Lasonolide A was discovered in 1994 by McConnell and coworkers in an effort to identify new and diverse antitumor agents from marine organisms.¹ The connectivity of the lasonolides and relative stereochemistry of each tetrahydropyran were determined by NMR correlation spectroscopy from isolated materials. In 2002, the correct fully elucidated structure of lasonolide A was disclosed in Lee's total synthesis.² This seminal work established the unknown relative stereochemistry of the C28 stereocenter and corrected the C17-C18 and C25-C26 olefin geometries, which had been incorrectly assigned (Figure 1). Importantly, the synthesis also established the absolute stereochemistry of the natural product and revealed that the levorotatory or (-)-lasonolide A was the biologically active enantiomer, contrary to what had been reported in the isolation paper. Despite the remarkable activity lasonolide A displayed in the NCI's 60-cell line screen,³ very few analogs have been prepared.4,5

RESULTS AND DISCUSSION

Our pursuit of the target molecule was not only inspired by its unique biological activity, but also its novel molecular architecture that presented us with the opportunity to execute and expand on many of the organic transformations that had previously been developed in our laboratory. We aimed at developing a more concise and efficient synthesis compared to those previously known. ^{2,6,7} A preliminary report of our efforts has been communicated.^{7a}

We recognized that the main point of diversification between lasonolides A-F occurred at the C28 and C30 positions (Figure 1. Subtle structural differences had dramatic effects on the biological profiles between each of the six natural analogs. While a comprehensive understanding of the structure activity relationships remains unclear, we felt that it was important to devise a synthesis plan (Scheme 1) that could allow for the installation of a range of diverse structural elements at these positions. A late stage Wittig olefination could install the required Z-olefin geometry present in all lasonolides and enable a flexible strategy for analog synthesis. The macrocycle can then be disconnected into two equally complex



Figure 1. Reassigned Lasonopyran Skeleton – Lasonolides A – G

tetrahydropyrans, which could be joined together by an esterification/macrolactonization and a Ru-catalyzed alkene-alkyne coupling. These disconnections of the lasonopyran skeleton allowed us to identify three sub-targets **9**, **10**, and **11** for synthesis.

Scheme 1. Synthesis Plan for (-)-Lasonolide A

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37 38 39

40

41 42

43

44

57 58 59

60



The hallmark of our synthetic plan, the Ru-catalyzed alkenealkyne coupling, was unique in the fact that we aimed to obtain a linear 1,4-diene product. Previously we demonstrated that branched selectivity is generally observed, as shown in eq.1.8 However, in a few cases, the linear product dominated (eq. 2). We have put forth a mechanistic rationale to understand this behavior. The proposed mechanism for the transformation, via ruthenacyclopentene formation, is depicted in Figure 2. The initial oxidative cycloisomerization with the Ru-catalyst is believed to be reversible and can form one of two possible regioisomers (A or B). A subsequent β -hydride and reductive elimination, from the metallacycles, can afford the linear or branched isomers. Steric factors suggest that intermediate B is more stable than intermediate A which accounts for the preferential formation of the branched product. However, the tautomerization of the initial alkene-alkyne complex C is thought to be faster than that of complex D then forms the new C-C bond between the sterically less-hindered terminus of each unsaturated partner. If the rate of β -hydride elimination of complex A can outcompete its cycloreversion to complex C, a linear product would result. Indeed, introduction

of a tetrasubstituted propargylic center inhibits cycloisomerization of complex \mathbf{D} to the point that the reaction is now favored via complex \mathbf{A} .

In all our prior total syntheses involving terminal alkyne partners, only branched products have been formed – alternaric acid,⁹ amphidinolides A^{10} and P,¹¹ and laulimalide.¹² The particular efficiency of the Ru catalyzed macrocyclization in the laulimalide synthesis stimulates exploration of the Ru catalyzed process making linear rather than branched products. To be applied to the total synthesis of lasonolide A, there are two possible approaches that could afford the desired product (Scheme 2). The first would feature an intramolecular reaction to generate the macrocycle (14) of lasonolide A. The second would be an intermolecular approach to form 19. Theoretically, in each scenario the linear isomer can be formed using either functional group orientation (i.e. 12 vs. 13 and 15/16 vs. 17/18).

We, therefore, initiated studies on representative model systems. Tetrahydropyrans 20^{13} and 21 were subjected to several reaction conditions to test the feasibility of the intermolecular coupling and to gain insight concerning the levels of linear:branched selectivity. We screened several solvents that have been previously utilized in alkene-alkyne couplings. No reaction was observed in non-coordinating solvents such as CH₂Cl₂ and DCE (Table 1, entries 1 and 2), likely due to catalyst decomposition. Next we screened coordinating solvents (acetone and DMF) (Table 1, entries 3-5) in an effort to stabilize the coordinatively unsaturated Ru-catalyst. To our delight, reactions run in DMF and acetone delivered the desired coupled products. The linear:branched ratios were much higher in acetone (3.5:1) than in DMF (1:1), suggesting that DMF promotes a reversible cycloisomerization and therefore Curtin-Hammett conditions. Increasing the amount of alkene 21 (5 equiv.) delivered the desired 1,4-diene (22) as a 4:1 mixture of linear:branched isomers in 56% yield.¹⁴ Importantly, 74% of the excess alkene was recovered at the end of the reaction.





Scheme 2. Proposed Ru-catalyzed Alkene-Alkyne Couplings



The alternative substitution pattern was also explored on a similar model system using the optimal conditions discovered in Table 1 - entry 5 (Scheme 3). In this case, the branched isomer (27) was exclusively formed in 33% yield.

Table 1. Model Studies for Alkene-Alkyne Coupling



Entry ^a	Solvent (Conc.)	21 (equiv) ^b	Conv ^c	22:23 ^c
1	CH_2Cl_2 (5 M)	1.2	0%	-:-
2	DCE (5 M)	1.2	0%	-:-
3	DMF (5 M)	1.2	33%	1:1
4	Acetone (5 M)	1.2	37%	3.5 : 1
5	Acetone (1 M)	5.0	56% ^d	4.0:1

^aAll reactions were run using 0.1 mmol (**20**), and 10 mol% [CpRu(MeCN)₃]PF₆ at rt for 14 h. ^b**21** was added as a solution drop-wise over 1h using a syringe pump. ^cDetermined by ¹H NMR. ^dIsolated yield.

The synthesis plan for alkyne **28** is presented in Scheme 4. A dinuclear Zn-catalyzed aldol reaction between ynone **32** and aldehyde **31** is envisioned to establish the absolute stereochemistry of fragment **30**. It is important that high enantioselectivity is attained, since the C21 hydroxyl group will dictate the formation of each subsequent stereocenter. The C22 quaternary stereocenter will be created from the thermodynamic formation of benzylidene acetal **29** via transacetilization. An analogous approach was previously reported by the Kang group in their total synthesis of (+)-lasonolide $A.^{7d}$ Finally, Horner-Wadsworth-Emmons (HWE) olefination with subsequent oxy-Michael addition should favor formation of the desired equatorial C23 stereocenter to complete the synthesis of alkyne **28**. Scheme 3. Additional Model Studies for Alkene-Alkyne Coupling



The synthesis of alkyne **28** commenced with an asymmetric aldol reaction between ynone **32** and aldehyde **31**.¹⁵ Under our standard set of conditions significant elimination was observed when reactions were run at room temperature (Table S2, entry 1), and at 4°C (Table S2, entry 2).¹⁶ The elimination could be suppressed by running the reaction at -20°C (Table S2, entry 3) for 72 h. This affords the desired β -hydroxy ketone (**30**) in a moderate 54% yield but with high levels of enantioselectivity (99% ee). Elimination of the β -hydroxy group could be entirely suppressed by modifying the work-up procedure. Switching from an aqueous work-up to a direct filtration through celite improved the isolated yield of **30** to 78% (Table S2, entry 4).

Scheme 4. Synthesis Plan for Alkyne 28



With an optimized protocol in hand, we then investigated a 1,3-*syn* reduction of β -hydroxy ketone **30** (Scheme 5). Under standard conditions (Et₂BOMe and NaBH₄)¹⁷ the reduction of **30** only afforded a ~4:1 mixture of diastereomers favoring the *syn*-diol. Switching to Kiyooka's conditions (DIBAL-H, - 78°C) increased the diastereoselectivity to 17:1 and the desired diol (**33**) was isolated in 98% yield.¹⁸ Selective protection of

the less sterically hindered secondary propargyl alcohol with TBDPSCl gave **34**.



1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21 22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46 47

48

49

50

51

52

53

54

55

56

57 58

59 60



We then aimed to establish the C22 quaternary stereocenter via transacetalization of 34 to form a thermodynamically favored benzylidene acetal.^{7d} Utilization of Kang's conditions (PhCHO, TFA, PhMe -20°C to rt, 3.5 h), led primarily to the formation of isomers 35c and 35d (Figure 3). Only trace amounts of **35a** and **35b** were detected in the crude ¹H NMR spectrum. Although the cyclization favored the incorrect isomer, we were encouraged that these products were isolated in near quantitative yield with little evidence of decomposition that may have arisen from the acidic conditions. Assuming that the observed products were not the thermodynamically preferred ones, we decided to conduct the acetal formation in a more polar solvent (CHCl₃) for an extended period of time. Under these new conditions the desired acetal (35a) was strongly favored in good yield, 5:1 chemoselectivity and 10:1 diastereoselectivity. Importantly, the undesired isomers (35b, 35c and 35d) could be separated from the predominant diasteromer (35a) by silica gel chromatography and re-subjected to the reaction conditions. After 2 rounds of recycling the undesired diastereomers, acetal 35a, containing the newly formed C22 guaternary stereocenter, was isolated in 93% yield.

With **35a** in hand we pushed forward towards the completion of the alkyne **28** (Scheme 6). The sequence began with exploring an oxidation of the primary neopentylic alcohol. Surprisingly, **35a** was resistant to oxidation using Dess-Martin periodinane, PDC, TEMPO, and under Moffatt-Swern conditions. Fortunately, Ley's catalytic TPAP/NMO oxidation reliably delivered aldehyde **36** in 76% yield.¹⁹ TBAF mediated deprotection of the TBDPS and TES protecting groups followed by concomitant cyclization produced lactol **38** in 59% yield. In addition to lactol **38**, carboxylic acid **37** was also isolated in 20% yield. We postulated the undesired side product was arising from a Cannizzaro reaction promoted by the hydroxide typically present in TBAF solutions. Buffering the reaction with acetic acid²⁰ suppressed formation of the carboxylic acid and improved the isolated yield to 94%.

Horner-Wadsworth-Emmons olefination and concomitant intramolecular oxy-Michael addition generated tetrahydropyran **39** as a single diastereomer detectable by ¹H NMR in 94% isolated yield.²¹ The stereochemistry of the tetrahydropyran was assigned by ROESY correlation.

Scheme 6. Synthesis of Alkyne 28



The completion of the alkyne 28 involved hydrolysis of the benzylidene acetal and cyclization of the resulting primary alcohol. Toward this end, a variety of acids were screened to effect the desired hydrolysis/cyclization sequence. The use of *p*-toluenesulfonic acid (MeOH)²² or acetic acid (THF, 50 $^{\circ}$ C)²³ gave only recovered starting material. Reactions involving BCl₃ (DCM, -78 °C to rt),²⁴ 2M HCl (MeOH, 65 °C),²⁵ or Amberlyst A-15 (4 Å MS, MeCN)²⁶ led to complicated mixtures of partially hydrolyzed products and alkyne 28 in <40%isolated yield. This partial conversion to the lactone was not unexpected since the equilibrium to remove the benzylidene requires the presence of both acid and water, whereas the equilibrium to form the lactone requires the presence of acid, but the exclusion of water/alcohol. Along these lines, we discovered that the desired lactol could be obtained in 53-82% isolated yield by utilizing a two-step procedure, which involved hydrolysis of the acetal with HCl in THF at 80 °C and then ring closure using catalytic *p*-TsOH in refluxing toluene.²⁷ We later discovered that formation of alkyne 28 could be accomplished in a single step using LiBF₄ in aq. MeCN in 96% yield.28



Figure 3. Selective Formation of Benzylidene Acetal 35a

The synthesis plan for alkene 11 is presented in Scheme 7. A dvnamic kinetic asymmetric reduction of β-ketoester 41 was envisioned to establish the absolute stereochemistry. The C7 stereocenter would be formed through a Michael addition, which could occur after the installation of an appropriate acceptor. Several possible routes were considered for the formation of the dienoate fragment. Our first tier of experiments would focus on incorporating this side chain via Horner-Wadsworth-Emmons olefination. Alternatively, if complications arose from the proposed HWE reaction, we also developed a convenient new strategy to form 2,4-dienoates that evolved from the reaction of terminal alkynes with ethyl diazoacetate²⁹ followed by subsequent phosphine-catalyzed isomerization.³⁰ A series of model studies, as illustrated in the conversion of alkyne 43 to dienoate 46 via intermediate 45, validated this latter approach (see Scheme 7).¹¹

Scheme 7. Synthesis Plans for Alkene 11



Initially, we were intrigued by the prospect of utilizing a transition metal catalyzed hydrogenation reaction to establish the absolute stereochemistry of fragment **11**.³¹ Although these highly atom economical³² methods proceed with high enantioand diastereoselectivities for a range of substrates, the chiral catalysts developed so far exhibit poor diastereoselectivities for methyl substituted β -ketoesters such as **41**.³³ As a result, we believed it would be worthwhile to investigate a microbial transformation that could establish the necessary absolute and relative stereochemistry via dynamic reduction of a racemic β ketoester. After surveying the literature, we discovered that Baker's yeast reductions of methyl-substituted β -ketoesters could provide the requisite products.^{16,34}

Scheme 8. Relative Stereochemical Assignment of 49



Encouraged by this precedent, we embarked on our synthesis of fragment **11** first by preparing the necessary starting material (Scheme 8). β -ketoester **41** was generated through a Blaise reaction with commercially available α -bromoester **47** and allyl cyanide.³⁵ Interestingly, when we investigated the reduction of β -ketoester **41** with Bakers' yeast (Sigma-Aldrich, YSC2, batch 035K0169) the undesired *anti*-product was formed with 10:1 diastereoselectivity (30% ee). Without heat treatment or additives, the diastereoselectivity for the reduc-

tion decreased to 4:1. The relative stereochemistry was further confirmed by nOe and *J*-coupling analysis on an advanced intermediate.³⁶These results were in direct contrast to the trends observed from the literature, therefore we decided to screen a number of isolated enzymes,³⁷ generously donated to us by Codexis Inc., to determine their performance in the dynamic kinetic asymmetric reduction of β -ketoester **41**. Two types of enzymes were examined in the reaction; the first was derived from Bakers' yeast and utilized GDH (glucose dehydrogenase) and NADP with glucose as the stoichiometric reductant.¹⁶ The pH of these reactions must be carefully controlled due to the accumulation of gluconic acid. The second type of enzyme utilizes NADPH as the cofactor and 2-propanol as the reductant. Table S5 of the SI provides a comprehensive screening of enzymatic methods.

NADP dependent ketoreductases, in general, showed enhanced selectivity for the *syn* product. However, under the standard conditions (Conditions A) using triethanolamine as a buffer, significant olefin isomerization of β -ketoester **41** was observed. After an exhaustive investigation of various buffers (pH 4.5 – 10 screened), we found that a pH 4.5 phosphate buffer completely suppressed the isomerization. CDX-024 demonstrated the highest diastereoselectivity favoring the *syn*-diastereomer (**48**) with high levels of enantioselectivity (see SI, Table S5, entries 7 and 8). Using this enzyme we were able to obtain the desired *syn* product (**48**) in 75% yield, with a 4 : 1 diastereoselectivity and greater than 95% ee (Table S5, entry 8).³⁸

Scheme 9. Synthesis of Diol 60



The synthesis of fragment **11** was advanced using the synthetic sequence depicted in Scheme 9. It began with the silyl protection of the C9 alcohol, which was subsequently followed by a 2-step Weinreb amide synthesis and Grignard addition to obtain ynone **52/53** in excellent yield.³⁹ (*S*)-CBS reduction under standard conditions ((*S*)-CBS, BH₃•DMS, THF) delivered the desired propargyl alcohols (**54/55**) in good yields (50-70%) as a 6.7 : 1 mixture of diastereomers. Yu has reported that the use of nitroethane as solvent has a profound effect on both reaction rate and stereoselectivity.⁴⁰ Pleasingly, we found that when we made this switch and used freshly distilled catechol borane the diastereoselectivity improved to >20:1.

Propargyl alcohol **54/55** was then subjected to a Ru-catalyzed hydrosilylation to generate trisubstituted (*Z*)-vinylsilane **56/57**.⁴¹ In each case (R=TBS, or R=TIPS) high selectivities

for a single geometric isomer (>15:1) were obtained. In preparation for tetrahydropyran formation via oxy-Michael addition, the enoate acceptor was installed through cross metathesis with ethyl acrylate.⁴² Finally, the C13 allyl moiety was introduced by a copper mediated cross coupling to form diol **60**.⁴³



Figure 4. THP Cyclizations of 60 and 62

1

2

3

4

5 6 7

8

9

10

11

12

13 14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 60

Deprotonation of diol 60 with sodium hydride generated tetrahydropyrans 61a/61b as a 1:1 mixture of diastereomers. Unfortunately, no improvement in diastereoselectvity was observed after screening various bases (t-BuOK and EtONa). solvents (DMF, THF, and DMSO) and reaction temperatures (-78°C, 0°C, rt and 60°C). Furthermore, equilibration of the diastereomeric mixture under basic conditions did not improve diastereoselectivity. We were aware that a similar approach to the southern tetrahydropyran ring was utilized in Kang's total synthesis of (+)-lasonolide A;^{7d} during their studies, an enhancement in diastereoselectivity was observed upon switching the protecting group on the C9 hydroxyl from TBS to TIPS. Likewise, we observed a modest increase in diastereoselectivity (1:1 to 2.4:1) for the cyclization when a TIPS protecting group was incorporated (62). This experiment supported the notion that this structural feature was important for improving selectivity in the cyclization.

Scheme 10. Further Studies Investigation THP Cyclizations



Several reports in the literature have indicated that cyclizations of secondary alcohols onto a proximal α , β -unsaturated aldehyde could enhance diastereoselectivity.⁴⁴ It is conceivable that the lower pKa of the proton adjacent to the aldehyde would render the reaction more reversible, and thereby favor the formation of the thermodynamically preferred 2,6-*cis* tetrahydropyran ring. To test this hypothesis, α , β -unsaturated aldehyde **65** was prepared (Scheme 10) in two steps from ester **60** and subjected to cyclization using DBU. We found that the diastereoselectivity was improved to ~4–5 : 1. Recalling our previous experience, we suspected that the incorporation of a protecting group at C9 could further enhance distereoselectivity. Upon preparing the TIPS protected substrate, spontaneous cyclization was observed upon oxidizing allylic alcohol 67 with MnO₂, and the desired tetrahydropyran (68) was isolated as a single diastereomer in an un-optimized 47% yield.

Scheme 11. Revised Synthetic Approach to Access THP 68



While encouraged by these results, access to allylic alcohol **67** was problematic due to several low yielding steps that were not amenable to scale-up. To circumvent these issues we decided to pursue an alternative reaction sequence that would feature a THP cyclization, bearing the necessary TIPS protection on the C9 hydroxyl, before installation of the C13 allyl segment (Scheme 11). However, in this alternate sequence, the alcohol required for the copper-mediated cross-coupling reaction would no longer be available. Therefore we decided to investigate a Hiyama cross coupling⁴⁵ for the late stage installation of the allyl moiety. At the time this work was executed, the use of allyl acetate in the Hiyama cross coupling had not been described.

Scheme 12. Synthesis of Vinyl Silane 71



The revised synthesis of fragment **11** began with the DIBAl-H reduction of ester **59** (Scheme 12). Allyic oxidation of alcohol **69** with MnO₂ afforded tetrahydropyran **70** directly in nearly quantitative yield as a single diastereomer.⁴⁶ Attempts to perform the Hiyama cross coupling with aldehyde **70** and allyl acetate were unsuccessful, presumably due to the insufficient stability of the starting aldehyde. To eliminate this problematic functionality, we first installed the dienoate moiety, before reexploring the Hiyama coupling. Elaboration of the aldehyde into *E*,*E*-dienoate **71** was accomplished via Horner-Wadsworth-Emmons olefination using 4 Å MS and LiOH.

Our investigation of the Hiyama coupling¹⁶ between allyl acetate⁴⁷ and vinyl silane **71** began with utilizing 2 mol% Pd₂dba₃•CHCl₃ and 4.2 equivalents of TBAF in a solution of THF (see Scheme 13 and SI Table S6). After 19 hours, a mixture of the desired product (**72**) along with an equimolar amount of silanol **73** (entry 1) was obtained in 75% combined yield. Extending the reaction time from 19 hours to 3 days also generated a 1:1 mixture of **72** : **73** but diminished the yield to 44% (entry 2). Increasing both the catalyst loading (10 mol%) and amount of TBAF (6.4 equiv.) was found to decrease the proportion of the remaining intermediate silanol **73**, and alkene **72** could be isolated in 78% yield (entry 3). It is important to note that, when we repeated the experiment, we noticed that the highest yields for this reaction (85%) were

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49 50 51

52

53

54

55

56

57 58 59

60

obtained when fresh TBAF solutions were used (entry 4). This modest increase in yield may be related to the water concentration present in TBAF solutions.⁴⁸ Alkene **75** was completed in two steps from **72**; saponification using lithium hydroxide, followed by TBS protection and *in situ* hydrolysis of the resulting silyl ester (Scheme 13).

Scheme 13. Synthesis of Alkene 75



Having achieved a synthesis of both fragments, we were poised to examine our key Ru-catalyzed alkene-alkyne coupling. Initial efforts were dedicated to exploring an intramolecular reaction to forge the macrocycle, since to date only three examples of intramolecular Ru-catalyzed alkene-alkyne couplings have been reported. These have appeared in the total syntheses of (+)-amphidinolide A, (+)-pinnatoxin $A_{,}^{49}$ and laulimalide. However, in each of these examples only branched products were generated. Figuring we possessed the necessary coordinating groups to favor the generation of a linear isomer, we were excited by the prospect of investigating a linear selective macrocyclization. Two substrates were examined in the macrocyclization reaction (76 and 78), each of which could easily be accessed using a Yamaguchi esterification.⁵⁰ Unfortunately, after extensive investigation we were never able to form the desired macrocycle as rapid decomposition was observed in all experiments that were attempted (Scheme 14).

We then decided to turn our attention to the exploration of an intermolecular alkene-alkyne coupling (Figure 5, eq 3).¹⁶ We were concerned that the π systems of electron deficient ester, present in alkene 11, could be coordinating to the ruthenium catalyst and inhibiting the coupling. As a result, we chose to use alkyne 28 in excess in an attempt to diminish this potential interaction (Table S7, entries 1-6). We observed 50% conversion of the alkene to the desired coupled products (80a and **80b**) as an inseparable 2:1 mixture of linear : branched isomers (entry 1). Unfortunately, the excess alkyne that was used could not be recovered as it seems to decompose over the course of the reaction, likely to the hydrated dimer.⁹ Heating the reaction to 50 °C had little effect on the overall conversion (entry 2). However, when the alkyne was added to the reaction drop-wise over 15 minutes and then heated at 50°C we noticed an increase in conversion to 70% (entry 3). Increasing

the concentration from 0.02 to 0.04 M (entry 4) also had no effect on conversion.

Scheme 14. Attempted Macrocyclization via Ru-Catalyzed Alkene-Alkyne Coupling



Content with our reaction conditions (entry 3), we increased the scale of the reaction from 5 μ mol to 60 μ mol in an attempt to obtain an isolated yield for the coupling. To our surprise, this seemingly minor modification was not tolerated and the conversion diminished significantly to ~20%. In this case, increasing the concentration to 0.12 M (entry 5) rescued the conversion slightly (35%) but not to the levels that were seen previously. Increasing both the catalyst loading and concentration had a deleterious effect on the overall conversion (entry 6).

Based on the experimental evidence presented in Table S7, entries 1-5 (see SI), we realized that our initial concerns related to catalyst deactivation via coordination to alkene 11 were largely attenuated due to the fact that in each of these entries, which successfully generated product, a 5-fold excess of alkene 11 to catalyst was present. Therefore we decided to screen reactions that used the alkene in excess. This change resulted in the complete consumption of alkyne 28 on a 68 µmol scale (Table S7, entry 7). Furthermore, after brief optimization, we determined that only 3 equiv. of alkene 11 were needed (using 10 mol% catalyst, in a 0.094 M solution of acetone) to achieve a 66% yield (entry 8).



Figure 5. Intermolecular Ru-Catalyzed Alkene-Alkyne Coupling





With a reliable synthetic route, we advanced our synthesis adhering to the current synthesis plan (Scheme 15). Saponification of the ester and lactone, followed by re-lactonization delivered seco acid **81**. Macrolactonization was attempted with a variety of reagents including 2,4,6-trichlorobenzoyl chloride, 2-methyl-6-nitrobenzoic anhydride,⁵¹ and dibutyltin oxide.⁵² Unfortunately, in all cases the macrocycle was never observed and the substrate decomposed under each of the reaction conditions. We speculated that the restricted conformational mobility of the C21 alcohol could be unable to access its optimal conformation for macrolactonization. Therefore, opening the lactone could enable further flexibility of the C21 alcohol and could allow for the desired cyclization to occur. As a result we decided to incorporate this idea into two modified substrates, **83** and **84**.

Scheme 16. Preparation of Wittig Salt 90 and Attempted Olefination with Lactol 91



We had hoped to access the elaborated seco acid (83) from lactone 80a via reduction to the lactol, subsequent Wittig olefination, and saponification. Disappointingly, after completing the synthesis of Wittig salt 90, the proposed reaction sequence was unsuccessful in our model system (Scheme 16). Warming the reaction above -15°C decomposed the unstabilized Wittig reagent rapidly.

 Table 2. Probing at Structure and Solvent Effects in Ru-Catalyzed Alkene-Alkyne Coupling



Entry ^a	R_1	R_2	R ₃	Solvent	Yield ^b	95a:95b ^c
1	Н	Н	in the second second	acetone	32%	3.2:1
2	Н	Н	in the second	cyclo- pentanone	39%	3.8 : 1
3	Н	in the second	in the second	acetone	41%	3.7:1
4	Н	it the	ž×ři	cyclo- pentanone	62%	3:1
5	TBS	in the	in the second se	cyclo- pentanone	36%	3.7 : 1

^aAll reactions were run using 3.0 equiv. **94**, 15 mol% [CpRu(MeCN)₃]PF₆, in 0.047 M solvent, at 55°C for 2 h 10 min. ^b Mixture of **95a**:**95b**. ^c Determined by ¹H NMR.

At this point, we decided to take a step back and, in turn, we began investigating a synthesis of seco acid **84**. Ideally, an alkene-alkyne coupling between triol **93** and alkene **94** could provide rapid access to seco acid **84**. Accordingly, we began screening conditions similar to those found to be optimal previously (Table S7). When acetone was used as solvent, we were able to obtain the desired products as their acetonides in 32% yield, as a 3.2:1 mixture of linear:branched isomers (Table 2, entry 1). Upon switching the solvent to cyclopentanone, the product ratio was enhanced to 3.8:1 (entry 2) – this was the highest ratio we had ever observed in our studies. We specu-

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

58 59

60

late that the enhanced selectivity could be related to several factors. Because the ruthenium catalyst has one open coordination site, the size of the solvent could have an effect on the product ratios. Alternatively, triol **93** might have been protected as its acetonide or cyclopentanone ketal before coupling occurs and the linear to branched ratios may be substrate specific. Finally, it is known that coordinating substituents in the substrates have dramatic effects on product distributions,⁵³ and the presence of the C25 primary alcohol may be contributing to selectivity.

We designed a series of experiments to probe these hypotheses. To determine the extent of contribution from solvent, we tested the cyclopentanone-protected substrate (Table 2, entry 3) in the coupling reaction using acetone as solvent. Only a slight decrease in the linear:branched ratio from 3.8:1 to 3.7:1 was observed. When the acetonide-protected substrate (entry 4) was tested, this time using cyclopentanone as solvent, we saw a decrease in linear:branched ratio to from 3.8:1 to 3:1. Finally, to probe the role of the C25 alcohol, we incorporated a TBS protecting group onto the cyclopentanone protected substrate (entry 5), expecting the linear:branched ratios to decrease if coordination was no longer possible. An identical 3.7:1 ratio of linear:branched isomers was obtained. The fact that the TBS protecting group was cleaved during the reaction invalidates any conclusions about the role of the hydroxyl group since the desilvlation could have occurred prior to or after the alkene-alkyne coupling.

Scheme 17. Formal Synthesis of (-)-Lasonolide A



Several proposed ruthenacyclopentene intermediates for both the linear and branched products are depicted in Figure 5. After oxidative coupling, ruthenium has one open coordination site and can be occupied by solvent (**S**) or by a coordinating atom present in the substrate. For the linear isomer, the oxygen in the tetrahydropyran ring or the primary alcohol could occupy this site. For the branched isomer only coordination of the primary alcohol may be possible but, as depicted in Figure 6, seems to be rather unfavorable. The facile desilylation that was observed in the reaction (Table 2, entry 5) suggests that the alcohol may be coordinated to the Lewis acidic ruthenium rendering it more susceptible towards desilylation. Taken together, these results begin to provide evidence that the product ratio is *highly* dependent on the identity of the substrate and is less influenced by the solvent (i.e. cyclopentanone vs. acetone).

From diene **95a**, protection of the primary alcohol as its TBDPS ether and hydrolysis of the cyclopentanone ketal provided diol **96**, an intermediate that was used in Shishido's total synthesis of (+)-lasonolide A (Scheme 17).^{7e} The physical data for diol **96** was in complete agreement with the reported data. The optical rotation of **96** was +11.4 (c 0.75, CHCl₃), opposite to that reported in the Shishido synthesis ($[\alpha]_D^{23}$ –19.3 (c 1.04, CHCl₃), and consistent with our absolute stereochemical assignment for (–)-lasonolide A.

Scheme 18. Synthesis of Alkyne 92



At this point, we decided to further streamline our synthesis of lasonolide A to help fulfill the need for material that will help advance biological investigations. We turned our attention back to the Ru-catalyzed coupling reaction and questioned if a more elaborated coupling partner could be tolerated under the reaction conditions. Alkyne **92**, which contains the fully elaborated side chain could be obtained in 4-steps from lactol **38** (Scheme 18). Although alkyne **92** contains two alkenes, disubstituted olefins are typically unreactive in the Ru-catalyzed coupling.⁵³

The alkene-alkyne coupling between **92** and **75** (Scheme 19), run under our standard set of conditions, took place in 43% yield with a linear:branched ratio of 3:1. The coupling between **92** and **74** also successfully occurred albeit with diminished linear:branched ratios (2:1) but in 69% yield. Reactions run in cyclopentanone led to poor conversion and slightly diminished linear:branched ratios in each case. Additionally, for the coupling between **92** and **74**, decreasing the amount of catalyst (from 15 mol% to 5 mol%) while also increasing the concentration (from 0.047 M to 0.14M) decreased the yield to 28% (51% of **92** as its acetonide was recovered) without affecting the linear:branched ratio. It is also important to note, that in each of the examples described above, >80% of the alkene **74/75** was recovered.



Figure 6. Possible Ruthenacyclopentene Intermediates for Linear and Branched Products

Scheme 19. Final Reaction Sequence Completing the Synthesis of (-)-Lasonolide A and Lasonolide Analogs



Completion of the synthesis was accomplished from both intermediates **98a** and **99a**.⁵⁴ Removal of the acetonide with CSA followed by protection of the most accessible alcohols in both **100** and **101** provided seco acid **102**, a common intermediate in both routes. Macrolactonization using the Yamaguchi reagent occurred without incident to provide the TBS protected lasonolide (**103**) in yields ranging from 40-62%. At this point, the undesired branched isomer that was generated during the coupling could be separated from the linear isomer. A final desilylation, using HF•Pyr, provided the target molecule, (–)-lasonolide A in 75% yield.

In addition to synthesizing the natural product we have also generated three analogs (Scheme 19). Compound **104** came from macrolactonization of the fully deprotected precursor hydroxyl acid. Compounds **105** and **106** were straightforward-ly derived from the branched by-products of the alkene-alkyne couplings. The synthetic lasonolide A and the 3 analogs were submitted to *in vitro* testing in an attempt to explore their activity against various cell lines.¹⁶ Each analog tested was essentially inactive compared to the synthetic (–)-lasonolide A in all assays except for the HCT116.

SUMMARY

1

6

7

8

9

10

11 12 13

14

15

16

17 18

19

20 21

22

23

24

25 26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

In conclusion, a synthesis of (–)-lasonolide A has been described. The synthesis has been accomplished in 16 linear steps and 34 total steps from commercially available starting materials. A formal intermediate was also prepared and matches the physical data that was reported in Shishido's synthesis of (+)-lasonolide A. Biological studies, utilizing the synthetic material generated from this work, are currently underway in an effort to further understand the mechanism of action. Importantly also, these studies verify that the Ru catalyzed alkene-alkyne coupling is amenable for making linear as well as branched 1,4-diene motifs en route to natural products, notably biologically active macrocycles. This success of the intermolecular coupling between fully unprotected polyhydroxy dieneyne 92 and tetraene 74/75 demonstrates a remarkable chemoselectivity.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures, analytical data (¹H NMR, ¹³C NMR, MS, IR, and $[a]_D$) for all new compounds as well as additional reaction optimization tables.

AUTHOR INFORMATION

Corresponding Author

*bmtrost@stanford.edu

Funding Sources

We thank the NIH (GM33049) for generous support to our programs. C.E.S., K.L.H., and A.H. are the recipients of NIH Ruth L. Kirschstein National Research Service Awards (GM101747, GM086093, and GM077840, respectively). C.P. is the recipient of a DAAD ISAP fellowship.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

We thank Johnson-Matthey for generous gifts of palladium and ruthenium salts and Codexis, Inc. for the generous donation of enzymes. Finally we would like to thank Genentech (Gail Phillips, Vishal Verma and John Flygare) for screening the synthetic (–)-lasonolide A and lasonolide analogs.

REFERENCES

(1) Horton, P. A.; Koehn, F. E.; Longley, R. E.; McConnell, O. J. J. Am. Chem. Soc. **1994**, *116*, 6015–6016.

(2) (a) Lee, E.; Song, H. Y.; Kang, J. W.; Kim, D.-S.; Jung, C.-K.; Joo, J. M. *J. Am. Chem. Soc.* **2002**, *124*, 384–385. (b) Song, H. Y.; Joo, J.M.; Kang, J.W.; Kim, D.-S.; Jung, C.-K.; Kwak, H.-S.; Park, J. H.; Lee, E.; Hong, C. Y.; Jeong, S. W.;

Jeon, K.; Park, J.H. J. Org. Chem. 2003, 68, 8080–8087.

(3) (a) Alley, M. C.; Scudiero, D. A.; Monks, A.; Hursey,
M. L.; Czerwinski, M. J.; Fine, D. L.; Abbott, B. J.; Mayo, J.
G.; Shoemaker, R. H.; Boyd, M. R. *Cancer Res.* 1988, 48, 589–601. (b) Zhang, Y.-W.; Ghosh, A. K.; Pommier, Y. *Cell Cycle* 2012, 11, 4424-4435. (c) Jossé, R.; Zhang, Y-W.;
Giroux, V.; Ghosh, A. K.; Luo, J.; Pommier, Y. *Mar. Drugs* 2015, 13, 3625–3639.

(4) (a) Lee, E.; Song, H. Y.; Joo, J.M.; Kang, J. W.; Kim, D.
S.; Jung, C. K.; Hong, C. Y.; Jeong, S. W.; Jeon, K. *Bioorg. Med. Chem. Lett.* 2002, *12*, 3519–3520. (b) Joo, J.M.; Kwak,
H. S.; Park, J. H.; Song, H.Y.; Lee, E. *Bioorg. Med. Chem. Lett.* 2004, *14*, 1905–1908.

(5) Additional lasonolides isolated from their natural source: Wright, A. E.; Chen, Y.; Winder, P. L.; Pitts, T. P.; Pomponi, S. A.; Longley, R. E. *J. Nat. Prod.* **2004**, *67*, 1351–1355.

(6) For synthetic studies towards the lasonolides see: (a) Ghosh, A. K.; Ren, G.-B. J. Org. Chem. 2012, 77, 2559–2565.
(b) Deba, T.; Yakushiji, F.; Mitsuru, S.; Shishido, K. Synlett, 2003, 10, 1500–1502. (c) Yoshimura, T.; Bando, T.; Shindo, M.; Shishido, K. Tetrahedron Lett. 2004, 45, 9241–9244. (d) Gurjar, M. K.; Kumar, P.; Rao, B. V. Tetrahedron Lett. 1996, 37, 8617–8620. (e) Gurjar, M. K.; Chakrabarti, A.; Rao, B. V.; Kumar, P. Tetrahedron Lett. 1997, 38, 6885–6888. (f) Nowakowski, M.; Hoffmann, H. M. R. Tetrahedron Lett. 1997, 38, 1001–1004. (g) Beck, H.; Hoffmann, H. M. R. Eur. J. Org. Chem. 1999, 11, 2991–2995. (h) Hart, D. J.; Patterson, S.; Unch, J. P. Synlett 2003, 9, 1334–1338. (i) Dalgrad, J. E.; Rychnovsky, S. D. Org. Lett. 2005, 7, 1589–1591. (j) Sawant, K. B.; Ding, F.; Jenings, M. P. Tetrahedron Lett. 2006, 47, 939–942.

(7) For total syntheses of lasonolide A see: (a) Trost, B. M.;
Stivala, C. E.; Hull, K. L.; Huang, A.; Fandrick, D. R. J. Am. Chem. Soc. 2014, 136, 88 – 91. (b) Ghosh, A. K.; Gong, G. Org. Lett. 2007, 9, 1437–1440. (c) Ghosh, A. K.; Gong, G. Chem. Asian J. 2008, 3, 1811–1823. (d) Kang, S. H.; Kang, S. Y.; Kim, C. M.; Choi, H.-w.; Jun, H. S.; Lee, B. M.; Park, C. M.; Jeong, J. W. Angew. Chem. Int. Ed. 2003, 42, 4779–4782.
(e) Yoshimura, T.; Yakushiji, F.; Kondo, S.; Wu, X.; Shindo, M.; Shishido, K. Org. Lett. 2006, 8, 475–478.

(8) (a) Trost, B. M.; Toste, F. D.; Pinkerton, A. B. *Chem. Rev.* **2005**, *101*, 2067–2096. (b) Trost, B. M.; Frederiksen, M. U.; Rudd, M. T. *Angew. Chem. Int. Ed.* **2005**, *44*, 6630–6666.

(9) Trost, B. M.; Probst, G.; Schoop, A. J. Am. Chem. Soc. **1998**, *120*, 9228 – 9236.

(10) (a) Trost, B. M.; Harrington, P. E.; Chisholm, J. D.; Wrobleski, S. T. J. Am. Chem. Soc. 2005, 127, 13598–13610.
(b) Trost, B. M.; Wrobleski, S. T.; Chisholm, J. D.; Harrington, P. E.; Jung, M. J. Am. Chem. Soc. 2005, 127, 13589–13597. (c) Trost, B. M.; Chisholm, J. D.; Wrobleski, S. J.; Jung, M. J. Am. Chem. Soc. 2002, 124, 12420–12421.

(11) (a) Trost, B. M.; Papillon, J. P. N. J. Am. Chem. Soc. **2004**, *126*, 13618-13619. (b) Trost, B. M.; Papillon, J. P. N.; Nussbaumer, T. J. Am. Chem. Soc. **2005**, *127*, 17921-17937.

(12) (a) Trost, B. M.; Amans, D.; Seganish W. M.; Chung, C. K. J. Am. Chem. Soc., 2009, 131, 17087–17089. (b) Trost, B.M.; Amans, D.; Seganish, W.M.; Chung, C.K. Chem. Eur. J. 2012, 18, 2961–2971. (c) Trost, B. M.; Seganish, W. M.; Chung, C. K.; Amans, D. Chem. Eur. J. 2012, 18, 2948–2960.

(13) Trost, B. M.; Yang, H.; Brindle, C. S.; Dong, G. *Chem-Eur. J.* **2011**, *17*, 9777–9788.

(14) It is possible that alkyne **20** can dimerize under the reaction conditions. For examples of Ru-catalyzed alkyne dimerizations see: (a) Trost, B. M.; Rudd, M. T. *J. Am. Chem. Soc.* **2001**, *123*, 8862–8863. (b) Trost, B. M.; Rudd, M. T. *J. Am. Chem. Soc.* **2002**, *124*, 4178–4179.

(15) Trost, B. M.; Fettes, A.; Shireman, B. T. J. Am. Chem. Soc. 2004, 126, 2660–2661.

(16) See Supporting Information for details.

(17) Chen, K.-M.; Hardtmann, G.E.; Pradad, K.; Repic, O.; Shapiro, M. J. *Tetrahedorn Lett.* **1987**, *28*, 155–158.

(18) (a) Kiyooka, S,-j.; Kuroda, H.; Shimasaki, Y. *Tetrahedron Lett.* **1986**, *27*, 3009–3012. (b) Mohr, P. *Tetrahedron Lett.* **1991**, *32*, 2219–2222.

(19) Ley, S. V.; Norman, J.; Griffith, W. P.; Marsden, S. P. Synthesis **1994**, *7*, 639–666.

(20) Smith, A. B.; Chen, S. S.-Y.; Nelson, F. C.; Reichert, J. M.; Salvatore, B. A. *J. Am. Chem. Soc.* **1995**, *117*, 12013–12014.

(21) HWE olefination and in situ cyclizations of lactols have been reported by: Allevi, P; Ciuffreda, P.; Colombo, D.; Mont, D.; Speranza, G.; Manitto, P. J. Chem. Soc. Perkin Trans 1 1989, 1281–1283.

(22) Figueroa-Perez, S.; Schmidt, R. R. Carbohydr. Res. 2000, 328, 95–102.

(23) Bennett, S. M.; Nguyen-Ba, N.; Ogilvie, K. K. J. Med. Chem. **1990**, *33*, 2162–2173.

(24) Bonner, T. G.; Bourne, E. J.; McNally, S. J. Chem. Soc. 1960, 2929–2934.

(25) Takano, S.; Hatakeyama, S.; Ogasawara, K. J. Chem. Soc., Perkin Trans 1 **1980**, 2, 457–461.

(26) Liu, Z.-Y.; Ji, J-X.; Li, B.-G. J. Chem. Soc., Perkin Trans 1 2000, 3519–3521.

(27) Chen, L.-Y.; Zaks, A.; Chackalamannil, S.; Dugar, S. J. Org. Chem. **1996**, *61*, 8341–8343.

(28) (a) Lipshutz, B. H.; Pegram, J. J.; Morey, M. C. Synth. Commun. **1982**, *12*, 267–277. (b) For a recent example of benzylidene acetal hydrolysis using LiBF₄ in a complex setting see: Araoz, R.; Servent, D.; Molgo, J.; Iorga, B. H.; Fruchart-Gaillard, C.; Benoit, E.; Gu, Z.; Stivala, C. E.; Zakarian, A. J. Am. Chem. Soc. **2011**, *133*, 10499–10511.

(29) Suarez, A.; Fu, G. C. Angew. Chem. Int. Ed. 2004, 43, 3580–3582.

(30) Trost, B. M.; Kazmaier, U. J. Am. Chem. Soc. 1992, 114, 7933–7935. (b) Trost, B. M.; Li, C.-J. J. Am. Chem. Soc. 1994, 116, 10819–10820. (c) Rychnovsky, S. D.; Kim, J. J. Org. Chem. 1994, 59, 2959–2660.

(31) (a) Noyori, R.; Ohkuma, T.; Kitamura, M. J. Am. Chem. Soc. **1987**, 109, 5856–5858. (b) Mashima, K.; Kusano, K.-h.; Sato, N.; Matsumura, Y.; Kyoko, N.; Kumobayashi, H.; Sayo, N.; Hori, Y.; Ishizaki, T.; Akutagawa, S.; Takaya, H. J. Org. Chem. **1994**, 59, 3064–3076. (c) Everaere, K.; Carpentier, J.-F.; Mortreux, A.; Bulliard, M. Tetrahedron Asym. **1999**, 10, 4083–4086.

(32) Trost, B. M. Science 1991, 254, 1471-1477.

(33) For examples of attempted diastereoselective asymmetric hydrogenation of α -methyl- β -keto-esters: (a) Kitamura,

M.; Ohkuma, T.; Tokunaga, M.; Noyori, R. *Tetrahedron Asym.* **1990**, *1*, 1–4. (b) Rychnovsky, S.D.; Hoye, R. C. *J. Am. Chem. Soc.* **1994**, *116*, 1753–1765. (c) Burk, M. J.; Harper, G.P.; Kalberg, C. S. *J. Am. Chem. Soc.* **1995**, *117*, 4423–4424.
(d) Yamano, T.; Taya, N.; Kawada, M.; Huang, T.; Imamoto, T. *Tetrahedron Lett.* **1999**, *40*, 2577–2580. (e) Zhang, Z.; Qian, H.; Longmire, J.; Zhang, X. J. Org. Chem. **2000**, *65*, 6223– 6226. (f) Ireland, T.; Tappe, K.; Grossheimann, G.; Knochel, P. *Chem. Eur. J.* , *8*, 843–852, (g) Evans, D. A.; Michael, F. E.; Tedrow, J. S.; Campos, K. R. J. Am. Chem. Soc. **2003**, *125*, 3539–3543.

(34) (a) Nakamura, K.; Kawai, Y.; Miyai, T.; Ohno, A. *Tetrahedron Lett.* **1990**, *31*, 3631-3632. (b) Nakamura, K.; Kawai, Y.; Ohno, A. *Tetrahedron Lett.* **1991**, *32*, 2927–2928. (c) Hamdani, M.; De Jeso, B.; Deleuze, H.; Maillard, B. *Tetrahedron Asym.* **1993**, *4*, 1229–1232. (d) Shieh, W.-R.; Sih, C. J. *Tetrahedron Asym.* **1993**, *4*, 1259–1269. (e) North, M. *Tetrahedron Lett.* **1996**, *37*, 1699–1702. (f) Ji, A.; Wolberg, M.; Hummel, W.; Wandrey, C.; Muller, M. *Chem. Commun.* **2001**, 57–58.

(35) Pedersen, O. S.; Peterson, L.; Brandt, M.; Nielsen, C.; Pedersen, E. B *Monatsh. Chem.* **1999**, *130*, 1499–1512.

(36) Stereochemistry also confirmed by comparing physical data to known β -hydroxyester: Hareau, G. P.-J.; Koiwa, M.; Hikichi, S.; Sato, F. *J. Am. Chem. Soc.* **1999**, *121*, 3640–3650.

(37) (a) Kalaitzakis, D.; Rozzell, J. D.; Kambourakis, S.; Smonou, I. *Org. Lett.* **2005**, *7*, 4799-4801. (b) Kalaitzakis, D.; Rozzell, J. D.; Kambourakis, S.; Smonou, I. *Eur. J. Org. Chem.* **2006**, 2309–2313.

(38) The absolute stereochemistry was established by conversion of a 4 : 1 (*syn:anti*) mixture of diastereomers to known saturated hydroxyl esters. See Supporting Information for details (a) Oppolzer, W.; Blagg, J.; Walther, E. J. Am. Chem. Soc. **1990**, *112*, 2767–2772. (b) Ghosh, A. K.; Onishi, M. J. Am. Chem. Soc. **1996**, *118*, 2527–2528.

(39) Williams, J. M.; Jobson, R. B.; Yasuda, N.; Marchesini, G.; Dolling, U.-H.; Grabowski, E. J. J. *Tetrahedron Lett.* **1995**, *36*, 5461–5464.

(40) Yu, C.-M.; Kim, C.; Kweon, J. H. Chem. Commun. 2004, 21, 2494–2495.

(41) Trost, B. M.; Ball, Z. T.; Joge, T. Angew. Chem. Int. Ed. 2003, 42, 3415–1418.

(42) Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. J. Am. Soc. Chem. **2000**, 122, 8168.

(43) Taguchi, H.; Ghoroku, K.; Tadaki, M.; Tsubouchi, A.; Takeda, T. Org. Lett. 2001, 3, 3811. b) Taguchi, H.; Ghoroku,

K.; Tadaki, M.; Tsubouchi, A.; Takeda, T. J. Org. Chem.
2002, 67, 8450. c) Tsubouchi, A.; Itoh, M.; Onishi, K.; Takeda, T. Synthesis 2004, 1504. d) Taguchi, H.; Miyashita, H.; Tsubouchi, A.; Takeda, T. Chem. Commun. 2002, 2218.
(e) Denmark, S.; Wehrli, D.; Choi, J. Org. Lett. 2000, 2, 2491.

(44) Paterson, I.; Arnott, E. A. *Tetrahedron Lett.* **1998**, *39*, 7185–7188.

(45) (a) Hiyama, T.; Hatanaka, Y. *Pure Appl. Chem.* **1994**, 66, 1471–1478. (b) Trost, B. M.; Machacek, M. R.; Ball, Z. T. *Org. Lett.* **2003**, *5*, 1895–1898. (c) Dey, R.; Chattopadhyay, K.; Ranu, B. C. *J. Org. Chem.* **2008**, *73*, 9461–9464. (d) For a review see: Denmark, S. E.; Sweis, R. F. *Chem. Pharm. Bull.* **2002**, *50*, 1531–1541.

(46) In our initial communication (Ref 7a) we described the formation of the substituted tetrahydropyran ring through a simultaneous Hoveyda-Grubbs 2nd generation catalyzed cross metathesis/oxa-Michael sequence: Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2000**, *122*, 8168–8179.

(47) Allyl chloride and allyl methyl carbonate could also be utilized in the Hiyama coupling.

(48) Hogrefe, R. I.; McCaffrey, A. P.; Corozdina, L. U.; McCampbell, E. S.; Vaghefi, M. W. *Nucleic Acids Res.* **1993**, *21*, 4739-4741.

(49) Nakamura, S.; Kikuchi, F.; Hashimoto, S. Angew. Chem. Int. Ed. 2008, 47, 7091–7094.

(50) (a) Yoshimura, T.; Yakushiji, F.; Kondo, S.; Wu, X.; Shindo, M.; Shishido, K. *Org. Lett.* **2006**, *8*, 475–478. (b) Inanaga, J.; Hirata, K.; Saeki, H.; Katsuki, T.; Yamaguchi, M. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 1989–1993.

(51) Shiina, I.; Kubota, M.; Oshiumi, H.; Hashizume, M. J. Org. Chem. **2004**, *69*, 1822–1830.

(52) Steliou, K.; Szczygielska-Nowosielska, A.; Favre, A.; Poupart, M. A.; Hanessian, S. J. Am. Chem. Soc. **1980**, 102, 7579–7581.

(53) Trost, B. M.; Cregg, J. J. J. Am. Chem. Soc. 2015, 137, 620–623.

(54) For clarity only the major (linear) isomer, from the alkene-alkyne coupling, is drawn. In the case where R=H a 2:1 ratio of products is advanced and in the case of R=TBS a 3:1 ratio of products is advanced. The linear and branched isomers become separable after macrolactonization.

Journal of the American Chemical Society

