

Role of Solvent and Effect of Substituent on Azobenzene Isomerization by Using Room-Temperature Ionic Liquids as Reaction Media

Guido Angelini,[†] Nadia Canilho,[‡] Mélanie Emo,[§] Molly Kingsley,^{||} and Carla Gasbarri^{*,†}

[†]Dipartimento di Farmacia, Università "G. d'Annunzio" di Chieti-Pescara, via dei Vestini, 66100 Chieti, Italy


[‡]Université de Lorraine, SRS MC, UMR7565, Vandœuvre-lès-Nancy F-54506 Cedex, France

[§]Institut J. Barriol, FR 2843, CNRS, Université de Lorraine, BP 70239, Vandœuvre-lès-Nancy F-54506 Cedex, France

^{||}University of Colorado Denver, Anschutz Medical Campus, 13001 East 17th Place, Aurora, Colorado 80045, United States

Supporting Information

ABSTRACT: The effects of a *para* substituent, as the electron-donating $-\text{OCH}_3$ and $-\text{OtBu}$ groups and the electron-withdrawing $-\text{Br}$ and $-\text{F}$ atoms, on azobenzene isomerization have been investigated in a series of imidazolium ionic liquids (BMIM PF₆, BMIM BF₄, BMIM Tf₂N, EMIM Tf₂N, BM₂IM Tf₂N, and HMIM Tf₂N). The thermal *cis*–*trans* conversion tends to be improved in the presence of the substituent, as pointed out by the first-order rate constants measured at 25 °C. Both the rotation and the inversion mechanisms occur in BMIM Tf₂N, EMIM Tf₂N, and HMIM Tf₂N, as highlighted by typical V-shape Hammett plots, but only rotation takes place in BMIM PF₆, BMIM BF₄, and BM₂IM Tf₂N. The possible interactions between the cation and the anion of the solvent and both the isomers of the azobenzene derivatives have been studied by small-wide-angle X-ray scattering (SWAXS). The calculated *cis* population in the photostationary state and the hardness parameter η of the *trans* isomer show that azobenzene and F-azobenzene are the less reactive molecules for the *trans*–*cis* conversion in all the investigated ionic liquids.



1. INTRODUCTION

Since Hartley's identification¹ of the *cis* isomer in 1937, the azobenzene molecule has been widely studied and successfully employed as a component of a large variety of systems, such as molecular devices,^{2,3} surface-modified materials,^{4,5} and light-sensitive biomolecules.^{6,7} The *trans*–*cis* isomerization can be induced by UV irradiation⁸ and electrostatic⁹ and mechanical stress¹⁰ leading to typical changes in the spectroscopic properties of azobenzene.¹¹ The thermal *cis*–*trans* conversion takes place spontaneously in the dark due to the higher stability of the *trans* isomer.¹² The mechanism by which the reaction occurs is not yet unequivocally established since competition between rotation and inversion has been proposed.^{13–15} Generally, the solvent properties, including polarity and viscosity, and the presence of substituents could promote one mechanism over the other^{16–19} so that azobenzene and derivatives find many applications as kinetic or spectroscopic probes in organic solvents^{20–22} and aqueous solutions.^{23–25} Room-temperature ionic liquids (RTILs) have attracted much attention in the last years for their unique properties²⁶ and their potential high recyclability.^{27,28} RTILs can be considered as designer solvents due to the possibility to change the cation or the anion in a wide range of combinations to induce strong changes in density, viscosity, melting point, and water miscibility.²⁹ Furthermore, different solute–solvent interactions

take place by using RTILs as reaction media since both the cation and the anion can be involved.³⁰ The effect of the solvent on the kinetics and the dynamics of azobenzene isomerization has been widely investigated,^{31–33} but to the best of our knowledge, the most recent studies in ionic liquids have concerned push–pull derivatives or a functionalized azobenzene moiety.^{34–36} The aim of this work has been the evaluation of the effect induced by one electron-donating or -withdrawing substituent ($-\text{OCH}_3$, $-\text{OtBu}$, $-\text{Br}$, and $-\text{F}$) in the *para* position on the azobenzene isomerization by using a BMIM⁺/Tf₂N[−]-based RTIL as solvent to highlight the role of the cation and the anion in the reaction. The structure of the investigated azobenzenes and RTILs are reported in Figure 1.

2. RESULTS AND DISCUSSION

It was assumed that the thermodynamically stable *trans* was the dominant isomer in the azobenzene solution before irradiation.^{37,38} The spectral changes induced by the exposition to UV light in BMIM Tf₂N are shown in Figure 2 as an example.

The kinetic profile of the thermal *cis*–*trans* isomerization follows a first-order decay, and the first-order rate constants k_{obs} determined at 25 ± 0.1 °C for the investigated azobenzenes in

Received: April 22, 2015




Figure 3. Hammett plots in EMIM Tf_2N (a) and BMIM BF_4 (b).




Figure 4. SWAXS spectra for Br- and OCH_3 -azobenzene isomers at 3×10^{-5} M in EMIM Tf_2N (a) and BMIM BF_4 (b).

Table 2. Hardness Parameter (eV) of the *Trans* Isomer and *Cis* Population after UV Irradiation in the Different Ionic Liquids

η_{trans}	<i>para</i> -substituent	<i>cis</i> isomer population (%)					
		HMIM Tf_2N	BM ₂ IM Tf_2N	EMIM Tf_2N	BMIM Tf_2N	BMIM PF_6	BMIM BF_4
1.98	H	49.4 ± 2	41.2 ± 4	55.1 ± 2	53.5 ± 1	60.7 ± 4	78.8 ± 2
1.86	OCH_3	93.8 ± 1	67.8 ± 6	96.1 ± 3	84.9 ± 9	97.4 ± 2	94.8 ± 2
1.93	OtBu	62.8 ± 4	81.8 ± 1	84.6 ± 1	91.2 ± 1	91.3 ± 5	85.9 ± 2
1.93	Br	80.8 ± 1	77.4 ± 2	81.4 ± 1	76.9 ± 1	86.8 ± 1	98.1 ± 2
1.97	F	58.4 ± 2	65.3 ± 1	63.4 ± 1	60.9 ± 2	61.7 ± 2	69.9 ± 1

to be not affected by the azobenzenes at the investigated concentrations, suggesting that the structural network of cations and anions of the RTILs is retained (Figure 4).

The rate of the thermal isomerization is independent of the initial concentration of the *cis* isomer in solution,⁶⁰ but the solvent and the nature of the substituents could influence the amount of the *cis* azobenzene formed by irradiation.^{19,61} The reactivity of the *trans* isomer to form the less stable *cis* by UV irradiation can be related to the η value. Generally, a higher η corresponds to a less reactive system.^{62,63} The *cis* population in the photostationary state obtained by using eq 1 and the hardness parameter η of the *trans* isomer for the investigated azobenzenes are reported in Table 2.

The formation of the *cis* isomer is strongly favored in the presence of the substituent in all the RTILs. The amount of the *cis* isomer of azobenzene increases from 49.4% in HMIM Tf_2N and 60.7% in BMIM PF_6 up to 93.8% and 97.4%, respectively, in the presence of the methoxy group.

The azobenzene molecule ($X = \text{H}$) is the less reactive system, as confirmed by the highest η , while the OCH_3 -azobenzene derivative is the most reactive between the

derivatives, as confirmed by the lowest η . Interestingly, F-azobenzene is the system closer to azobenzene for *cis* population and η value, suggesting a similar low reactivity in the *trans*–*cis* conversion.

3. CONCLUSIONS

Both rotation and inversion take place for the *cis*–*trans* azobenzene isomerization in BMIM Tf_2N , EMIM Tf_2N , and HMIM Tf_2N , according to the electronic nature of the substituent. A change of mechanism from inversion to rotation has been observed for Br- and F-azobenzene in BMIM BF_4 , BMIM PF_6 , and BM₂IM Tf_2N , according to the viscosity of the solvent. The structural organization of cations and anions seems to be not affected by the presence of the azobenzene isomers as suggested by SWAXS measurements. The *cis* population of the azobenzene derivatives increases in the photostationary state and the calculated η values of the *trans* isomer suggest that OCH_3 -azobenzene is the most reactive system of the investigated molecules for the *trans*–*cis* conversion. The azobenzene isomerization still represents a model reaction and the results presented herein could

contribute to the understanding and the improving of RTILs as reaction media.

4. EXPERIMENTAL SECTION

General Information. Azobenzene and ethanol (99% spectroscopy grade) were purchased from Fluka and used without further purification; *p*-OCH₃-azobenzene, *p*-OtBu-azobenzene, *p*-Br-azobenzene and *p*-F-azobenzene were obtained as previously described.⁶⁴ BMIM Tf₂N (1-butyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide), BMIM PF₆ (1-butyl-3-methylimidazolium hexafluorophosphate), BMIM BF₄ (1-butyl-3-methylimidazolium tetrafluoroborate), EMIM Tf₂N (1-ethyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide), BM₂IM Tf₂N (1-butyl-2,3-dimethylimidazolium bis(trifluoromethylsulfonyl)imide) and HMIM Tf₂N (1-hexyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide) were purchased from IoLiTec and Solvionics and stirred under vacuum at 60 °C overnight before the preparation of the samples.

Sample Preparation for the Kinetics Measurements. Stock ethanol solutions from the investigated azobenzenes were prepared at a concentration of 2.6×10^{-3} M and kept in the dark at room temperature at least for 4 days before use. Then, an appropriate amount was transferred into a 1 cm light pass quartz cuvette containing 200 μL of the proper ionic liquid to obtain a final concentration of 2.6×10^{-5} M. The sample was irradiated for 45 min by using a Hg–Xe arc lamp (150 W) equipped with a band-pass interference filter centered at 365.0 ± 0.2 nm wavelength and 10.0 ± 2.0 nm bandwidth: the spectral changes, as the decreasing of the high-intensity absorption band centered in the wavelength range 315–349 nm (depending both on the substituent and the solvent) and the increasing of the low-intensity band at about 440 nm (due to an n → π* transition) were used as evidence for the *trans*–*cis* isomerization. The first-order rate constants k_{obs} were spectrophotometrically calculated by monitoring in the dark the absorption change at or near the absorption maximum of the *trans* isomer over a period of about 24 h.

Determination of the *Cis* Population and the Hardness Parameter. The *cis* population in the photostationary state formed by UV irradiation was calculated as follows⁶¹

$$\% \text{ cis} = 100(A_{\text{trans}} - A_{\text{trans}(i)})/A_{\text{trans}} \quad (1)$$

where A_{trans} and $A_{\text{trans}(i)}$ are the absorbance of the *trans* isomer before and after the irradiation, respectively. The hardness parameter η values were calculated from the HOMO and LUMO energies⁶² of the *trans* isomers by using DFT (B3LYP/6-31G**).

SWAXS Analysis. Small-wide-angle X-ray scattering (SWAXS) experiments were performed by using an instrument with a line collimation setup. The system used a Cu K α radiation source in a sealed tube ($\lambda = 0.1542$ nm) working at 40 kV and 50 mA. The system is equipped with a translucent beam stop, which allows the determination of the $q = 0$ position by the direct measurement of an attenuated primary beam. The scattering signals were acquired on a sensitive SWAXS imaging plate slide as a detector (q range 0.28–28 nm⁻¹). In that configuration, the distance sample–detector is of 261.2 nm. The samples at 3×10^{-5} and 3×10^{-2} M azobenzene concentrations were introduced into a quartz capillary with a diameter of 1 mm and a sealed special glass capillary with a diameter of 1.5 mm, respectively. The scattering signals were collected over 30 min under vacuum at room temperature and treated with SAXSquant software. All data were corrected for the background scattering from the empty cell and for slit-smearing effects by a desmearing procedure using the Lake method.

■ ASSOCIATED CONTENT

● Supporting Information

Hammett plots in the RTILs, HOMO and LUMO energies of the *trans* isomers used for the determination of η , additional SWAXS spectra, and computational data. The Supporting

Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.joc.5b00898.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: c.gasbarri@unich.it. Tel: +39-0871-3554786.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work has been supported by MIUR (PRIN 2010-2011, prot. 2010N3T9M4) and University “G. d’Annunzio” of Chieti-Pescara.

■ REFERENCES

- (1) Hartley, G. S. *Nature* **1937**, *140*, 281–281.
- (2) Murakami, H.; Kawabuchi, A.; Kotoo, K.; Kunitake, M.; Nakashima, N. *J. Am. Chem. Soc.* **1997**, *119*, 7605–7606.
- (3) Balzani, V.; Credi, A.; Venturi, M. *Molecular Devices and Machines - A Journey into the Nano World*; Wiley-VCH: Weinheim, 2003.
- (4) Wen, Y.; Yi, W.; Meng, L.; Feng, M.; Jiang, G.; Yuan, W.; Zhang, Y.; Gao, H.; Jiang, L.; Song, Y. *J. Phys. Chem. B* **2005**, *109*, 14465–14468.
- (5) Ferri, V.; Elbing, M.; Pace, G.; Dickey, M. D.; Zharnikov, M.; Samori, P.; Mayor, M.; Rampi, M. A. *Angew. Chem., Int. Ed.* **2008**, *47*, 3407–3409.
- (6) Willner, I.; Rubin, S. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 367–385.
- (7) El Halabieh, R. H.; Mermut, O.; Barrett, C. J. *Pure Appl. Chem.* **2004**, *76*, 1445–1465.
- (8) Zacharias, P. S.; Ameerunisha, S.; Korupoju, S. R. *J. Chem. Soc., Perkin Trans. 2* **1998**, 2055–2059.
- (9) Tong, X.; Pelletier, M.; Lasia, A.; Zhao, Y. *Angew. Chem., Int. Ed.* **2008**, *47*, 3596–3599.
- (10) Turansky, R.; Konopka, M.; Doltsinis, N. L.; Stich, I.; Marx, D. *Phys. Chem. Chem. Phys.* **2010**, *12*, 13922–13932.
- (11) Knoll, H. In *CRC Handbook of Organic Photochemistry and Photobiology*, 2nd ed.; Horspool, W., Lenci, F., Eds.; CRC Press: Boca Raton, 2004; Vol. 89, 1–16.
- (12) Rau, H.; Lueddecke, E. *J. Am. Chem. Soc.* **1982**, *104*, 1616–1620.
- (13) Cimiraglia, R.; Asano, T.; Hofmann, H. *J. Gazz. Chim. Ital.* **1996**, *126*, 679–684.
- (14) Cattaneo, P.; Persico, M. *Phys. Chem. Chem. Phys.* **1999**, *1*, 4739–4743.
- (15) Cembran, A.; Bernardi, F.; Garavelli, M.; Gagliardi, L.; Orlandi, G. *J. Am. Chem. Soc.* **2004**, *126*, 3234–3243.
- (16) Bortolus, P.; Monti, S. *J. Phys. Chem.* **1979**, *83*, 648–652.
- (17) Malkin, S.; Fischer, E. *J. Phys. Chem.* **1962**, *66*, 2482–2486.
- (18) Asano, T.; Yano, T.; Okada, T. *J. Am. Chem. Soc.* **1982**, *104*, 4900–4904.
- (19) Bandara, H. M. D.; Burdette, S. C. *Chem. Soc. Rev.* **2012**, *41*, 1809–1825.
- (20) Whitten, D. G.; Wildes, P. D.; Pacifici, J. G.; Irick, G. *J. Am. Chem. Soc.* **1971**, *93*, 2004–2008.
- (21) Asano, T.; Okada, T. *J. Phys. Chem.* **1984**, *88*, 238–243.
- (22) Shin, D. M.; Whitten, D. G. *J. Am. Chem. Soc.* **1988**, *110*, 5206–5208.
- (23) Gille, K.; Knoll, H.; Quitzsch, K. *Int. J. Chem. Kinet.* **1999**, *31*, 337–350.
- (24) Wang, R.; Knoll, H.; Quitzsch, K. *Langmuir* **2001**, *17*, 2907–2912.
- (25) De Maria, P.; Fontana, A.; Gasbarri, C.; Siani, G.; Zanirato, P. *Arkivoc* **2009**, No. viii, 16–29.
- (26) Rogers, R. D.; Voth, G. A. *Acc. Chem. Res.* **2007**, *40*, 1077–1078.
- (27) Earle, M. J.; Seddon, K. R. *Pure Appl. Chem.* **2000**, *72*, 1391–1398.

- (28) Seddon, K. R. *J. Chem. Technol. Biotechnol.* **1997**, *68*, 351–356.
- (29) Hagiwara, R.; Ito, Y. *J. Fluorine Chem.* **2000**, *105*, 221–227.
- (30) Angelini, G.; Gasbarri, C. *Curr. Drug. Targets* **2015**, DOI: 10.2174/1389450115666141114152340.
- (31) Asano, T.; Okada, T. *J. Org. Chem.* **1984**, *49*, 4387–4391.
- (32) Wazzan, N. A.; Richardson, P. R.; Jones, A. C. *Photochem. Photobiol. Sci.* **2010**, *9*, 968–974.
- (33) Schanze, K. S.; Mattox, T. F.; Whitten, D. G. *J. Org. Chem.* **1983**, *48*, 2808–2813.
- (34) Baba, K.; Ono, H.; Itoh, E.; Itoh, S.; Noda, K.; Usui, T.; Ishihara, K.; Inamo, M.; Takagi, H. D.; Asano, T. *Chem. - Eur. J.* **2006**, *12*, 5328–5333.
- (35) Zhang, S.; Liu, S.; Zhang, Q.; Deng, Y. *Chem. Commun.* **2011**, *47*, 6641–6643.
- (36) Asaka, T.; Akai, N.; Kawai, A.; Shibuya, K. *J. Photochem. Photobiol., A* **2010**, *209*, 12–18.
- (37) Duarte, L.; Fausto, R.; Reva, I. *Phys. Chem. Chem. Phys.* **2014**, *16*, 16919–16930.
- (38) Sekkat, Z.; Wood, J.; Knoll, W. *J. Phys. Chem.* **1995**, *99*, 17226–17234.
- (39) Crecca, C. R.; Roitberg, A. E. *J. Phys. Chem. A* **2006**, *110*, 8188–8203.
- (40) Gasbarri, C.; Angelini, G. *RSC Adv.* **2014**, *4*, 17840–17845.
- (41) Gasbarri, C.; Angelini, G.; Fontana, A.; De Maria, P.; Siani, G.; Giannicchi, I.; Dalla Cort, A. *Biochim. Biophys. Acta, Biomembr.* **2012**, *1818*, 747–752.
- (42) Angelini, G.; Pisani, M.; Mobbili, G.; Marini, M.; Gasbarri, C. *Biochim. Biophys. Acta, Biomembr.* **2013**, *1828*, 2506–2512.
- (43) Alunni, S.; Del Giacco, T.; De Maria, P.; Fontana, A.; Gasbarri, C.; Ottavi, L. *J. Org. Chem.* **2004**, *69*, 6121–6123.
- (44) De Maria, P.; Fontana, A.; Siani, G.; D'Aurizio, E.; Cerichelli, G.; Chiarini, M.; Angelini, G.; Gasbarri, C. *Colloids Surf., B* **2011**, *87*, 73–78.
- (45) Fontana, A.; Viale, M.; Guernelli, S.; Gasbarri, C.; Rizzato, E.; Maccagno, M.; Petrillo, G.; Aiello, C.; Ferrini, S.; Spinelli, D. *Org. Biomol. Chem.* **2010**, *8*, 5674–5681.
- (46) Hansch, C.; Leo, A.; Taft, R. W. *Chem. Rev.* **1991**, *91*, 165–195.
- (47) Herkstroeter, W. *J. Am. Chem. Soc.* **1973**, *95*, 8686–8691.
- (48) McDaniel, D. H.; Brown, H. C. *J. Org. Chem.* **1958**, *23*, 420–427.
- (49) Talaty, E. R.; Fargo, J. C. *Chem. Commun.* **1967**, *2*, 65–66.
- (50) Nishiyama, Y.; Fukuda, M.; Terazima, M.; Kimura, Y. *J. Chem. Phys.* **2008**, *128*, 164514–164522.
- (51) Zhang, S.; Sun, N.; He, X.; Lu, X.; Zhang, X. *J. Phys. Chem. Ref. Data* **2006**, *35*, 1475–1517.
- (52) Asano, T.; Okada, T.; Shinkai, S.; Shigematsu, K.; Kusano, Y.; Manabe, O. *J. Am. Chem. Soc.* **1981**, *103*, 5161–5165.
- (53) Schanze, K. S.; Mattox, T. F.; Whitten, D. G. *J. Am. Chem. Soc.* **1982**, *104*, 1733–1735.
- (54) Asano, T. *J. Am. Chem. Soc.* **1980**, *102*, 1205–1207.
- (55) Gegiou, D.; Muszkat, K. A.; Fischer, E. *J. Am. Chem. Soc.* **1968**, *90*, 12–18.
- (56) Binenboym, J.; Burcat, A.; Lifshitz, A.; Shamir, J. *J. Am. Chem. Soc.* **1966**, *88*, 5039–5041.
- (57) Ab Rani, M. A.; Brant, A.; Crowhurst, L.; Dolan, A.; Lui, M.; Hassan, N. H.; Hallett, J. P.; Hunt, P. A.; Niedermeyer, H.; Perez-Arlandis, J. M.; Schrems, M.; Welton, T.; Wilding, R. *Phys. Chem. Chem. Phys.* **2011**, *13*, 16831–16840.
- (58) Ue, M. *J. Electrochem. Soc.* **1994**, *141*, 3336–3342.
- (59) Dupont, J. *J. Braz. Chem. Soc.* **2004**, *15*, 341–350.
- (60) Ortruba, J. P.; Weiss, R. G. *J. Org. Chem.* **1983**, *48*, 3448–3453.
- (61) Joshi, N. K.; Fuyuki, M.; Wada, A. *J. Phys. Chem. B* **2014**, *118*, 1891–1899.
- (62) Piyanzina, I.; Minisini, B.; Tayurskii, D.; Bardeau, J. F. *J. Mol. Model.* **2015**, *21*, 34.
- (63) Parr, R. G.; Pearson, R. G. *J. Am. Chem. Soc.* **1983**, *105*, 7512–7516.
- (64) Vecchi, I.; Arcioni, A.; Bacchiocchi, C.; Tiberio, G.; Zanirato, P.; Zannoni, C. *J. Phys. Chem. B* **2007**, *111*, 3355–3362.