Chiral Silver Phosphate Catalyzed Transformation of *ortho*-Alkynylaryl Ketones into 1*H*-Isochromene Derivatives through an Intramolecular-Cyclization/Enantioselective-Reduction Sequence**

Masahiro Terada,* Feng Li, and Yasunori Toda

Abstract: The transformation of ortho-alkynylaryl ketones through a cyclization/enantioselective-reduction sequence in the presence of a chiral silver phosphate catalyst afforded 1Hisochromene derivatives in high yield with fairly good to high enantioselectivity. An asymmetric synthesis of the 9oxabicyclo[3.3.1]nona-2,6-diene framework, which has been found in some biologically active molecules, is presented as a demonstration of the synthetic utility of this method.

sobenzopyrylium ions are stable oxonium cations owing to their aromaticity as a 10π -electron system. In general, these cationic species have been generated in situ by various methods for further manipulation, although the corresponding salts can be isolated as air-stable solids with less nucleophilic anions, such as BF₄⁻, and are utilized both for the characterization of these distinctive species and in electrophilic transformations.^[1–5] Their high and unique reactivity has attracted great attention during the past decade, because these oxonium species undergo a diverse array of transformations, including Diels-Alder reactions and nucleophilic addition reactions,^[2-4] and can be applied to the synthesis of natural products and biologically relevant molecules.^[5] However, catalytic enantioselective reactions of the isobenzopyrylium ions remain largely unexploited despite their unique reactivity and synthetic utility, presumably owing to the planar structure of the isobenzopyrylium ion, which lacks apparent coordination sites that can interact with chiral metal catalysts or organocatalysts.^[6,7] Indeed, successful examples are guite limited and have been based on the use

[*] Prof. Dr. M. Terada, F. Li, Dr. Y. Toda Department of Chemistry, Graduate School of Science Tohoku University Aramaki, Aoba-ku, Sendai 980-8578 (Japan)
E-mail: mterada@m.tohoku.ac.jp Homepage: http://www.orgreact.sakura.ne.jp/index.html
Prof. Dr. M. Terada Research and Analytical Center for Giant Molecules Graduate School of Science, Tohoku University Aramaki, Aoba-ku, Sendai 980-8578 (Japan)

[**] This research was partially supported by a Grant-in-Aid for Scientific Research on Innovative Areas "Advanced Molecular Transformations by Organocatalysts" from MEXT (Japan) and a Grant-in-Aid for Scientific Research (Grant No. 23655077) from the JSPS. We gratefully acknowledge Prof. T. Iwamoto and Prof. S. Ishida (Graduate School of Science, Tohoku University) for their kind assistance with X-ray crystallographic analysis. We also thank the JSPS for a research fellowship for Young Scientists (Y.T.).

Supporting information for this article is available on the WWW under http://dx.doi.org/10.1002/anie.201307371.

of metal catalysts with electronically neutral chiral ligand(s).^[6] Hence, the development of enantioselective transformations involving the isobenzopyrylium ion as a reactive intermediate is still a challenging topic in synthetic organic chemistry.^[8,9] To develop such a catalytic enantioselective transformation, we took advantage of the chemical properties of the isobenzopyrylium ion; notably, the formation of ion-pair salts between the isobenzopyrylium ion and less nucleophilic counteranions. In principle, the use of a chiral instead of an achiral counteranion should enable reactions to take place in an enantioselective fashion.^[9]

Chiral-counteranion-induced enantioselective transformations in metal-catalyzed reactions, namely, chiral-anion catalysis,^[10] has emerged as a powerful and attractive tool in asymmetric synthesis in recent years owing to its distinct advantages over conventional methods that rely on metal complexes coupled with electronically neutral chiral ligand(s). Chiral-anion catalysis offers another approach to the enantioselective reaction of isobenzopyrylium ions with a controlled stereochemical outcome.^[8a,9] In this context, we envisioned an enantioselective sequential transformation based on chiral-anion catalysis in which the key intermediate, the isobenzopyrylium ion, is generated in situ through metal-



Scheme 1. Enantioselective transformation involving an isobenzopyrylium cation as the reactive intermediate.

Angew. Chem. Int. Ed. 2014, 53, 235-239

complex-catalyzed intramolecular cyclization of *ortho*-alkynylaryl ketones **2**. The proposed sequence involves a two-step transformation (Scheme 1): 1) intramolecular cyclization of **2** catalyzed by a π -Lewis acidic metal complex with a binolderived phosphate **1** (binol = 1,1'-binaphthalene-2,2'-diol) as the chiral counteranion^[11] to generate an ion pair comprised of the isobenzopyrylium intermediate **A** and chiral phosphate **1**⁻, and 2) enantioselective reduction of this key intermediate **A** with a Hantzsch ester **3**^[12] to afford the final product **4** in an enantioselective manner under the influence of **1**⁻. Herein, we report the enantioselective transformation of *ortho*alkynylaryl ketones **2** into optically active 1*H*-isochromene derivatives **4** through a cyclization/enantioselective reduction sequence with a chiral silver phosphate catalyst and **3** as the reducing agent.

We began our investigation by exploring achiral π -Lewis acidic metal complexes with the ortho-alkynylaryl ketone 2a $(R^1 = Me, R^2 = phenyl, R^3 = H)$ as the primary substrate for the precursor of the isobenzopyrylium ion. The sequential transformation was performed with 10 mol% of the metal catalyst and 1.1 equivalents of Hantszch ester 3a in THF at room temperature. The reactivity and mode of cyclization were highly dependent on the metal catalyst employed (Table 1). Ag^I and Cu^{II} catalysts afforded the desired 6-endo cyclization product 4a exclusively in excellent yield (Table 1, entries 1 and 2, respectively). In contrast, the reactions with Pt^{II}, Au^{III}, and Au^I catalysts afforded not only the desired 6endo product 4a, but also the 5-exo product 5a (Table 1, entries 3, 4, and 6, respectively), although the Au^{III} and Au^I catalysts exhibited higher catalytic activity than that of the Ag^I catalyst (Table 1, entries 4 and 6 versus entry 1). To verify the participation of the π -Lewis acidic metal species in this reaction, trifluoromethanesulfonic acid (TfOH) was employed as a metal-free Brønsted acid catalyst (Table 1,

Table 1: Initial screening of π -Lewis acidic metal catalysts for the cyclization/reduction of 2a.^[a]



Entry	Metal catalyst	<i>t</i> [h]	Conversion of 2 a [%] ^[b]	Yield of 4a [%] ^[b]	Yield of 5 a [%] ^[b]
1	AgOTf	3	100	96	_
2	Cu(OTf)₂	24	96	96	-
3	PtCl ₂	24	35	18	5
4	AuCl₃	1	100	38	50
5	AuCl/AgOTf	24	trace	_	-
6	[AuCl(PPh ₃)]/AgOTf	1	100	24	58
7	TfOH	24	trace	_	-

[a] Reaction conditions: transition-metal catalyst (0.02 mmol, 10 mol%), **2a** (0.2 mmol), **3** (0.22 mmol, 1.1 equiv), THF (1 mL), room temperature. [b] Determined by ¹H NMR spectroscopic analysis of the crude product. Tf=trifluoromethanesulfonyl. entry 7). As expected, this strong Brønsted acid did not afford any product, and the starting material 2a was almost completely recovered. As the catalyst AgOTf exhibited exclusive formation of the 6-*endo* product 4a and higher reactivity than Cu(OTf)₂, Ag^I was chosen as the metal species for further studies on an enantioselective variant of the present sequential transformation.

The enantioselective variant of the reaction was conducted with the chiral silver phosphate Ag[(R)-1] (10 mol%), **2a**, and **3a** (1.1 equiv) in THF at room temperature. Ag[(R)-1] was preprepared from Ag_2CO_3 and the corresponding binol-derived phosphoric acid (*R*)-1 (2 equiv).^[11a,13] We screened a range of chiral silver phosphates by changing the substituents at the 3- and 3'-positions of the binaphthol backbone (Table 2, entries 1–4). The pentafluorophenyl-sub-

Table 2: Screening of chiral silver catalysts for the cyclization/enantioselective reduction of 2a to 4a and control experiments.^[a]

Ph	catalyst (10 mol%) 3a (1.1 equiv)	Ph
< <u><</u>	THF (0.2 м), RT	\sim
2a	(<i>R</i>)- 1a : G = C ₆ F ₅ - b : G = 9-anthryl	4a
	c : G = 2,4,6- <i>i</i> Pr ₃ C ₆ H ₂ -	
	d : G = C ₆ H ₅ -	

Entry	Catalyst	<i>t</i> [h]	Additive	Yield [%] ^[b]	ee [%] ^[c]
1	Ag[(R)- 1 a]	4	-	94	85
2	Ag[(R)- 1 b]	9	-	84	37
3	Ag[(R)- 1c]	7	-	94	66
4	Ag[(R)-1 d]	5	-	77	41
5	$Ag_2CO_3^{[d]}$	24	-	trace ^[e]	-
6	(R)-1 a	24	-	trace ^[e]	-
7 ^[f]	$Ag_2CO_3/(R)-1a$	5	_	94	83
8 ^[g]	Ag[(R)- 1 a]	48	3 Å MS	80 ^[h]	86
9 ^[g]	Ag[(R)- 1 a]	54	4 Å MS	32 ^[]	77
10 ^[g]	Ag[(R)- 1 a]	6	5 Å MS	95	87

[a] Reaction conditions (unless otherwise noted): catalyst (0.02 mmol, 10 mol%), **2a** (0.2 mmol), **3a** (0.22 mmol, 1.1 equiv), THF (1 mL), room temperature. [b] Yield of isolated **4a**. The consumption of **2a** was monitored by TLC. [c] The *ee* value of **4a** was determined by chiral-stationary-phase HPLC analysis. [d] The reaction was carried out with Ag₂CO₃ (0.01 mmol, 5 mol%). [e] Ketone **2a** was recovered in nearly quantitative yield. [f] The catalyst was generated in situ from Ag₂CO₃ (0.01 mmol, 5 mol%) and (*R*)-**1a** (0.02 mmol, 10 mol%). [g] The reaction was carried out with MS (75 mg), Ag[(*R*)-**1a**] (0.015 mmol, 10 mol%), **2a** (0.15 mmol), and **3a** (0.165 mmol, 1.1 equiv) in THF (0.75 mL). [h] Ketone **2a** was recovered in 19% yield. [i] Ketone **2a** was recovered in 66% yield.

stituted chiral silver phosphate Ag[(R)-1a] was found to provide the highest chemical yield and enantioselectivity (Table 2, entry 1). Ag[(R)-1a] promoted the consecutive reactions smoothly to afford the desired product 4a in 94% yield with fairly good enantioselectivity (85% ee). The use of Ag_2CO_3 or the chiral phosphoric acid alone did not afford any product, and the starting ketone 2a was recovered in a nearly quantitative manner (Table 2, entries 5 and 6). These control experiments strongly suggest that Ag[(R)-1] is essential for this reaction sequence to proceed in an enantioselective

236 www.angewandte.org

manner. The chiral silver phosphate generated in situ from (R)-1**a** and Ag₂CO₃ was also examined (Table 2, entry 7). However, preprepared Ag[(R)-1**a**] displayed slightly higher enantioselectivity than that of the catalyst generated in situ. We then explored the use of other additives to improve the enantioselectivity of the reaction (Table 2, entries 8–10). The addition of 3 Å and 4 Å molecular sieves (MS) significantly retarded the reaction, and the starting ketone 2**a** was not consumed completely, even after 2 days (Table 2, entries 8 and 9, respectively). In particular, a considerable reduction in both the chemical yield and the *ee* value was observed in the presence of 4 Å MS. In contrast, when 5 Å MS were used as an additive, even higher enantioselectivity was observed without any detrimental effect on the chemical yield (Table 2, entry 10).

Next, we explored the mechanism of the silver phosphate catalyzed sequential transformation of **2** into **4** to confirm the participation of the isobenzopyrylium ion as the reactive intermediate. From a mechanistic viewpoint, it can be considered that there are two possible reaction pathways. One pathway involves intramolecular cyclization followed by enantioselective reduction of the isobenzopyrylium intermediate (Scheme 1). However, the same product would be obtained if the reactions occurred in the reverse order, that is, reduction to the alcohol, followed by cyclization.^[14] To verify the order of the reaction sequence, we conducted a control experiment with alcohol **6** as a possible intermediate (Scheme 2). When compound **6** was subjected to the catalytic



Scheme 2. Mechanistic study: control experiment.

reaction under the influence of Ag[(R)-1a] (10 mol%) without the Hantzsch ester, the reaction proceeded with very low conversion, even after 6 h, and product 5a of 5-exo cyclization was obtained in 10% yield, whereas almost none of the desired 6-endo cyclization product 4a was observed. This behavior is in contrast to that of the present reaction of the primary substrate 2a, which afforded the 6-endo cyclization product 4a exclusively in excellent yield within 4 h in the presence of the Hantzsch ester (Table 2, entry 1). Therefore, these results strongly suggest that alcohol 6 is not involved as an intermediate in the present sequential transformation, and that the isobenzopyrylium species participates as the key intermediate, as shown in Scheme 1. Thus, it was concluded that the reaction is initiated by the coordination of Ag[(R)-1]to the C-C triple bond of 2. This interaction induces intramolecular nucleophilic attack by the carbonyl oxygen atom to generate an ion pair consisting of the isobenzopyrylium ion and the chiral phosphate. Subsequent enantioselective reduction by Hantzsch ester 3, followed by metal-proton exchange, regenerates the catalytic species and releases the isochromene product $\mathbf{4}^{[15,16]}$

Having identified the sequential process involving an isobenzopyrylium reactive intermediate, the scope of the reaction was then investigated with a range of alkynylaryl ketones **2** under the optimized reaction conditions. A broad range of alkynylaryl ketones are applicable to the present sequential transformation, although reactivity and enantioselectivity are highly dependent on the nature of the substituents (Table 3). When alkyl and aryl groups were introduced at

Table 3: Scope of the chiral silver phosphate catalyzed transformation of **2** into $\mathbf{4}^{[a]}$



Entry	2 (R ¹ , R ² , R ³)	<i>t</i> [h]	4	Yield [%] ^[b]	ee [%] ^[c]
1	2b (Me, 4-MeC ₆ H₄-, H)	11	4b	98	81
2	2c (Me, 4-CF ₃ C ₆ H ₄ -, H)	10	4c	87	67
3	2d (Me, 4-MeOC ₆ H ₄ -, H)	5	4 d	92	80
4	2e (<i>n</i> Pr, Ph, H)	9	4e	95	81
5	2 f (<i>i</i> Bu, Ph, H)	6	4 f	89	82
6	2g (Me, Ph, F)	6	4g	89	87
7	2h (Me, <i>n</i> Bu, H)	2	4h	89	22
8	2i (Ph, Ph, H)	24	4i	85	92
9	2j (4-MeOC ₆ H₄-, Ph, H)	26	4j	81	90
10	2k (4-BrC ₆ H₄-, Ph, H)	26	4 k	81	91 ^[d]
11	2I (4-CF ₃ C ₆ H ₄ -, Ph, H)	30	41	84	91
12	2m (3-MeOC ₆ H ₄ -, Ph, H)	29	4m	81	91
13	2n (2-TBSOC ₆ H₄-, Ph, H)	24	4 n	90	88
14	20 (Ph, Ph, F)	28	4 o	68	90
15	2p (Ph, <i>n</i> Bu, H)	8	4 p	86	49

[a] Reactions were carried out with Ag[(R)-1 a] (0.015 mmol, 10 mol%), 2 (0.15 mmol), and 3 (0.165 mmol, 1.1 equiv). For 2b-h, reactions were carried out with 3 a (R = Et) in THF (0.75 mL) in the presence of 5 Å MS (75 mg). For 2i-p, reactions were carried out with 3 b (R = Me) in EtOAc (0.75 mL) in the absence of MS. [b] Yield of isolated 4. [c] The *ee* value of 4 was determined by chiral-stationary-phase HPLC analysis. [d] The absolute configuration of 4 k was determined to be *S* by X-ray crystallographic analysis.^[17] See the Supporting Information for details.

the R^1 and R^2 positions, respectively (Table 3, entries 1–6), these substrates exhibited relatively high reactivity and afforded isochromene derivatives **4** in high yields. However, in terms of the enantioselectivity, the introduction of a *para* substituent on the aromatic group at the R^2 position (substrates **2b–d**) led to a slight decrease in enantioselectivity as compared to that observed with the primary substrate **2a** ($R^2 = Ph$). Substrates bearing linear or branched alkyl chains at the R^1 position (**2e** and **2f**, respectively) also underwent the reaction with a slight reduction in enantioselectivity, whereas fluoro substitution on the tethering benzene ring (**2g**) did not compromise the stereochemical outcome. However, the introduction of an alkyl substituent at the R^2 position led to



a considerable decrease in enantioselectivity (Table 3, entry 7). The reactions of substrates 2i-o bearing aromatic substituents at the R^1 position and a phenyl group at the R^2 position were very sluggish under the optimized conditions and proceeded in low chemical yields, although the ee values of the products were around 90%. To our delight, however, these substrates were smoothly converted under modified reaction conditions [with 3b (R = Me) in ethyl acetate instead of THF and in the absence of 5 Å MS] to give the corresponding isochromene products 4i-n in good yields with high enantioselectivity (Table 3, entries 8-13). A variety of aryl groups at the R¹ position were tolerated in this reaction (substrates 2j-n; Table 3, entries 9–13). Even with a bulky tert-butyldimethylsilyloxy (OTBS) substituent in the ortho position, the corresponding product **4n** was obtained in 90% yield with 88% ee (Table 3, entry 13). The introduction of a fluoro substituent on the tethering benzene ring of a diarylsubstituted ketone (substrate 20) did, however, result in a slight decrease in chemical yield, but with little detrimental effect on the enantioselectivity (Table 3, entry 14). However, when an alkyl substituent was introduced at the R² position, a considerable reduction in enantioselectivity was observed (Table 3, entry 15).

Finally, the synthetic utility of the present method was demonstrated (Scheme 3). 1-Allyl 1*H*-isochromenes, such as 4q, have been utilized as key intermediates for the synthesis



Scheme 3. Derivatization of the reaction product. Ts = p-toluene-sulfonyl.

of the 9-oxabicyclo[3.3.1]nona-2,6-diene framework,^[18] and this bicyclic ring structure has been found in some biologically active molecules.^[19] We thus attempted to construct the fundamental structure of these compounds in an optically active form by utilizing our methodology followed by an acidcatalyzed annulation reaction. Compound **2q** was subjected to the optimized reaction conditions, and the 1*H*-isochromene product **4q** was obtained with fairly good enantioselectivity (84 % *ee*). In the presence of *p*-TsOH·H₂O (5 mol %), **4q** was then converted into the desired bicyclic product **7a** and the regioisomer **7b** in moderate overall yield for the two steps. During the course of the acid-catalyzed annulation, no loss of enantiomeric purity was observed. In conclusion, we have successfully developed a cyclization/enantioselective-reduction sequence for the transformation of *ortho*-alkynylaryl ketones under the catalysis of a chiral silver phosphate complex to afford 1*H*-isochromene derivatives in high yields with fairly good to high enantioselectivity. A mechanistic study strongly suggested that an isobenzopyrylium cation is involved as the key intermediate in this transformation. We also demonstrated the utility of this methodology for the asymmetric synthesis of the 9oxabicyclo[3.3.1]nona-2,6-diene framework, which has been found in some biologically active molecules. Further development of the present methodology for application not only to other types of nucleophiles, but also to other transformations involving cationic species that are difficult to perform by other conventional methods will be completed in due course.

Experimental Section

Typical procedure for the reaction of 2a-h: MS (5 Å; 75.0 mg), Hantzsch ester 3a (41.8 mg, 0.165 mmol), and Ag[(R)-1a] (10 mol%, 11.8 mg, 0.015 mmol) were placed in a dry test tube. The atmosphere was replaced with argon, THF (0.75 mL) was added to the mixture, and the resulting mixture was stirred vigorously for 10 min. Ketone 2a(32.8 mg, 0.15 mmol) was then added, and the reaction mixture was stirred for 6 h at room temperature and then filtered through a pad of Al₂O₃ (with EtOAc as the eluent). The filtrate was concentrated, and the crude product was purified by flash column chromatography on silica gel with hexane/EtOAc/Et₃N (100:1:1) as the eluent to afford 4a(31.5 mg, 95%) as a white solid. The *ee* value was determined by chiral-stationary-phase HPLC analysis. See the Supporting Information for the reaction of 2i-p.

Received: August 21, 2013 Revised: October 7, 2013 Published online: November 24, 2013

Keywords: asymmetric catalysis · cyclization · enantioselective reduction · isobenzopyrylium ions · silver salts

- For reviews, see: a) E.-V. Kuznetsov, I.-V. Shcherbakova, A.-T. Balaban, Adv. Heterocycl. Chem. 1990, 50, 157–254; b) M. Nogradi, Sci. Synth. 2003, 14, 201–273; c) J. Santamaría, C. Valdés, Mod. Heterocycl. Chem. 2011, 3, 1631–1682; for selected examples, see: d) J. D. Tovar, T. M. Swager, J. Org. Chem. 1999, 64, 6499–6504; e) I. Prauda, J. Reiter, J. Heterocycl. Chem. 2001, 38, 199–204; f) I. Prauda, I. Kövesdi, P. Trinka, J. Reiter, J. Heterocycl. Chem. 2001, 38, 403–414; g) T. Suzuki, C. Okada, K. Arai, A. Awaji, T. Shimizu, K. Tanemura, T. Horaguchi, J. Heterocycl. Chem. 2001, 38, 1409–1418; h) W.-G. Wei, Z.-J. Yao, J. Org. Chem. 2005, 70, 4585–4590.
- [2] For selected examples of metal-catalyzed transformations via isobenzopyrylium salts, see: a) N. Asao, K. Takahashi, S. Lee, T. Kasahara, Y. Yamamoto, J. Am. Chem. Soc. 2002, 124, 12650–12651; b) N. Asao, T. Kasahara, Y. Yamamoto, Angew. Chem. 2003, 115, 3628–3630; Angew. Chem. Int. Ed. 2003, 42, 3504–3506; c) N. Asao, T. Nogami, S. Lee, Y. Yamamoto, J. Am. Chem. Soc. 2003, 125, 10921–10925; d) J.-L. Zhu, A.-R. Germain, J. A. Porco, Jr., Angew. Chem. 2004, 116, 1259–1263; Angew. Chem. Int. Ed. 2004, 43, 1239–1243; e) J.-L. Zhu, N. P. Grigoriadis, J.-P. Lee, J. A. Porco, Jr., J. Am. Chem. Soc. 2005, 127, 9342–9343; f) N. Asao, K. Sato, Menggenbateer, Y. Yamamoto, J. Org. Chem. 2005, 70, 3682–3685; g) H. Kusama, H. Funami, M. Shido, Y. Hara, J. Takaya, N. Iwasawa, J. Am. Chem. Soc. 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 2005, 70, 3682–3685; g) H. Kusama, H. Funami, M. Shido, Y. Hara, J. Takaya, N. Iwasawa, J. Am. Chem. Soc. 2005, 200

127, 2709–2716; h) S. Shin, A.-K. Gupta, C.-Y. Rhim, C.-H. Oh, *Chem. Commun.* **2005**, 4429–4431; i) A.-B. Beeler, S. Su, C.-A. Singleton, J. A. Porco, Jr., *J. Am. Chem. Soc.* **2007**, *129*, 1413– 1419; j) Y.-C. Hsu, C.-M. Ting, R.-S. Liu, *J. Am. Chem. Soc.* **2009**, *131*, 2090–2091.

- [3] For selected examples of iodonium-mediated transformations via isobenzopyrylium salts, see: a) J. Barluenga, H. Vázquez-Villa, A. Ballesteros, J.-M. González, J. Am. Chem. Soc. 2003, 125, 9028–9029; b) J. Barluenga, H. Vázquez-Villa, A. Ballesteros, J.-M. González, Org. Lett. 2003, 5, 4121–4123; c) D. Yue, N. Della Cà, R.-C. Larock, Org. Lett. 2004, 6, 1581–1584; d) J. Barluenga, H. Vázquez-Villa, I. Merino, A. Ballesteros, J. M. González, Chem. Eur. J. 2006, 12, 5790–5805; e) D. Yue, N. Della Cà, R. C. Larock, J. Org. Chem. 2006, 71, 3381–3388.
- [4] For reactions of isolated isobenzopyrylium salts, see: a) Z.-L.
 Hu, W.-J. Qian, S. Wang, S. Wang, Z.-J. Yao, *Org. Lett.* 2009, 11, 4676–4679; b) Z.-L. Hu, W.-J. Qian, S. Wang, S. Wang, Z.-J. Yao, *J. Org. Chem.* 2009, 74, 8787–8793.
- [5] a) G. Dyker, D. Hildebrandt, J. Org. Chem. 2005, 70, 6093-6096;
 b) K. Sato, N. Asao, Y. Yamamoto, J. Org. Chem. 2005, 70, 8977-8981; c) M. A. Marsini, K. M. Gowin, T. R. R. Pettus, Org. Lett. 2006, 8, 3481-3483; d) W.-J. Qian, W.-G. Wei, Y.-X. Zhang, Z.-J. Yao, J. Am. Chem. Soc. 2007, 129, 6400-6401; e) Y.-S. Yao, Z.-J. Yao, J. Org. Chem. 2008, 73, 5221-5225.
- [6] a) D. Hojo, K. Noguchi, K. Tanaka, Angew. Chem. 2009, 121, 8273-8276; Angew. Chem. Int. Ed. 2009, 48, 8129-8132; b) S. Handa, L.-M. Slaughter, Angew. Chem. 2012, 124, 2966-2969; Angew. Chem. Int. Ed. 2012, 51, 2912-2915; for a catalytic enantioselective reaction of 2-oxonium-1,3-dienes, see: c) K. Ishida, H. Kusama, N. Iwasawa, J. Am. Chem. Soc. 2010, 132, 8842-8843.
- [7] For catalytic enantioselective transformations of benzopyrylium ions, see: a) F. Benfatti, E. Benedetto, P. G. Cozzi, *Chem. Asian J.* 2010, *5*, 2047–2052; b) P. N. Moquist, T. Kodama, S. E. Schaus, *Angew. Chem.* 2010, *122*, 7250–7254; *Angew. Chem. Int. Ed.* 2010, *49*, 7096–7100; c) M. Rueping, C. M. R. Volla, I. Atodiresei, *Org. Lett.* 2012, *14*, 4642–4645; d) C.-C. Hsiao, H.-H. Liao, E. Sugiono, I. Atodiresei, M. Rueping, *Chem. Eur. J.* 2013, *19*, 9775–9779; e) M. Terada, T. Yamanaka, Y. Toda, *Chem. Eur. J.* 2013, *19*, 13658–13662.
- [8] For an enantioselective transformation of an isobenzopyrylium ion containing a hydroxy group at the 4-position with a chiral phosphoric acid, see: a) M. Terada, Y. Toda, Angew. Chem. 2012, 124, 2135–2139; Angew. Chem. Int. Ed. 2012, 51, 2093–2097; for an enantioselective transformation of isobenzopyrylium-4-olates with a binary catalytic system composed of a Rh^{II} catalyst and a chiral Lewis acid catalyst, see: b) H. Suga, K. Inoue, S. Inoue, A. Kakehi, J. Am. Chem. Soc. 2002, 124, 14836–14937; c) H. Suga, K. Inoue, S. Inoue, A. Kakehi, M. Shiro, J. Org. Chem. 2005, 70, 47–56; d) H. Suga, T. Suzuki, K. Inoue, A. Kakehi, Tetrahedron 2006, 62, 9218–9225; e) H. Suga, S. Higuchi, M. Ohtsuka, D. Ishimoto, T. Arikawa, Y. Hashimoto, S. Misawa, T. Tsuchida, A. Kakehi, T. Baba, Tetrahedron 2010, 66, 3070–3089.
- [9] During the preparation of our manuscript, an enantioselective transformation of isobenzopyrylium ions was reported: S.-Y. Yu, H. Zhang, Y. Gao, L. Mo, S. Wang, Z.-J. Yao, *J. Am. Chem. Soc.* 2013, *135*, 11402–11407.
- [10] For recent reviews on chiral-anion catalysis, see: a) J. Lacour, D. Moraleda, *Chem. Commun.* 2009, 7073-7089; b) R. J. Phipps, G. L. Hamilton, F. D. Toste, *Nat. Chem.* 2012, *4*, 603-614; c) M. Mahlau, B. List, *Angew. Chem.* 2013, *125*, 540-556; *Angew. Chem. Int. Ed.* 2013, *52*, 518-533; d) K. Brak, E. N. Jacobsen, *Angew. Chem.* 2013, *125*, 558-588; *Angew. Chem. Int. Ed.* 2013,

52, 534-561; see also: e) A. Parra, S. Reboredo, A. M. M. Castro, J. Alemán, Org. Biomol. Chem. 2012, 10, 5001-5020.

- [11] For selected examples of chiral-anion catalysis by chiral phosphates in transition-metal-catalyzed reactions, see: a) G. L. Hamilton, E. J. Kang, M. Mba, F. D. Toste, Science 2007, 317, 496-499; b) S. Mukherjee, B. List, J. Am. Chem. Soc. 2007, 129, 11336-11337; c) M. Rueping, A. P. Antonchick, C. Brinkmann, Angew. Chem. 2007, 119, 7027-7030; Angew. Chem. Int. Ed. 2007, 46, 6903-6906; d) C. Li, C. Wang, B. Villa-Marcos, J. Xiao, J. Am. Chem. Soc. 2008, 130, 14450-14451; e) S. Liao, B. List, Angew. Chem. 2010, 122, 638-641; Angew. Chem. Int. Ed. 2010, 49, 628-631; f) G. Jiang, R. Halder, Y. Fang, B. List, Angew. Chem. 2011, 123, 9926-9929; Angew. Chem. Int. Ed. 2011, 50, 9752-9755; g) V. Rauniyar, Z. J. Wang, H. E. Burks, F. D. Toste, J. Am. Chem. Soc. 2011, 133, 8486-8489; h) J. R. Zbieg, E. Yamaguchi, E. L. McInturff, M. J. Krische, Science 2012, 336, 324-327; i) Z. Chai, T. J. Rainey, J. Am. Chem. Soc. 2012, 134, 3615-3618; j) Z.-L. Tao, W.-Q. Zhang, D.-F. Chen, A. Adele, L.-Z. Gong, J. Am. Chem. Soc. 2013, 135, 9255-9258.
- [12] For reviews on chiral Brønsted acid catalyzed enantioselective transfer hydrogenation with Hantzsch esters as the reducing agent, see: a) S.-L. You, *Chem. Asian J.* 2007, *2*, 820–827; b) M. Rueping, E. Sugiono, F. R. Schoepke, *Synlett* 2010, 852–865; c) C. Zheng, S.-L. You, *Chem. Soc. Rev.* 2012, *41*, 2498–2518.
- [13] For the preparation of silver phosphates from Ag₂CO₃ and a chiral phosphoric acid, see: K. Aikawa, M. Kojima, K. Mikami, *Angew. Chem.* **2009**, *121*, 6189–6193; *Angew. Chem. Int. Ed.* **2009**, *48*, 6073–6077.
- [14] a) R. Yanada, S. Obika, H. Kono, Y. Takemoto, *Angew. Chem.* **2006**, *118*, 3906–3909; *Angew. Chem. Int. Ed.* **2006**, *45*, 3822–3825; b) S. Obika, H. Kono, Y. Yasui, R. Yanada, Y. Takemoto, *J. Org. Chem.* **2007**, *72*, 4462–4468.
- [15] As shown in Scheme 1, we propose that the metal species is attached to the isobenzopyrylium ion. However, during the reduction step, the actual structure of the cationic species has not been confirmed yet. Before enantioselective reduction, an exchange from the silver atom to a hydrogen atom would also be possible, and the silver metal would then not be involved in the stereodetermining step. We also conducted a reaction with $Cu[(R)-1a]_2$ as the chiral catalyst, and product 4a was obtained with 86% *ee*, albeit in moderate yield. This enantioselectivity is nearly equal to that observed upon Ag[(R)-1a] catalysis (Table 2, entry 1: 85% *ee*). These results imply that the metal species may not participate in the enantioselective reduction step.
- [16] When the 4,4-dideuterium-substituted Hantzsch ester was employed, deuterium was introduced at the C1 position exclusively. This result strongly suggests that the reaction proceeded through the formation of an isobenzopyrylium ion as the reactive intermediate. See the Supporting Information for details.
- [17] CCDC 954470 [(S)-4k] contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [18] S. Bhunia, K.-C. Wang, R.-S. Liu, Angew. Chem. 2008, 120, 5141-5144; Angew. Chem. Int. Ed. 2008, 47, 5063-5066.
- [19] a) B. Wünsch, M. Zott, G. Höfner, Arch. Pharm. 1992, 325, 733–739; b) B. Wünsch, M. Zott, G. Höfner, Arch. Pharm. 1993, 326, 823–830; c) R. Dupont, L. Jeanson, J.-F. Mouscadet, P. Cotelle, Bioorg. Med. Chem. Lett. 2001, 11, 3175–3178; d) C. Maurin, F. Bailly, P. Cortelle, Curr. Med. Chem. 2003, 10, 1795–1810.

Angew. Chem. Int. Ed. 2014, 53, 235-239